

Araştırma

THE EFFECTS OF RESUSCITATION FLUIDS ON HEMODYNAMIC PARAMETERS AND BLOOD BIOCHEMISTRY OF RATS BLED TO HYPOVOLEMIA

Fethi GÜL¹, Osman EKİNCİ¹, Tuba PELİT¹, Berna TERZİOĞLU², Gülşen BOSNA¹, Mehmet ERŞAHİN³, Neşe AYDIN¹, M. Zafer GÖREN⁴

THE EFFECTS OF RESUSCITATION FLUIDS ON HEMODYNAMIC PARAMETERS AND BLOOD BIOCHEMISTRY OF RATS BLED TO HYPOVOLEMIA

SUMMARY

Purpose: Hemorrhagic shock is a rare but serious complication which may lead to hemodynamic instability, decrease in oxygen delivery and decreased tissue perfusion, cellular hypoxia, organ damage and death. Current study was designed to study the efficiency of whole blood, low molecular weight starch solution (LWS) and hypertonic saline with dextrane (HSD) on hemodynamics parameters, coagulation and tissue perfusion of rats bled to hypovolemia.

Methods: Experiments were performed with Sprague Dawley rats (n=7 per group), weighing 200–250 g. The rats were bled through left iliac artery for 20 min in three successive steps until the mean arterial pressure fell to and stabilized at approximately 20 mmHg. Then animals in each group was treated either with HSD (4 ml/kg; i.v.) or LWS solution (4 ml/kg; i.v.) or stored whole blood transfusion (2 ml/100 g; i.v.) for 30 minutes.

Results: In rats treated either with whole blood or HSD, the mean arterial pressure values were found to be higher than the rats treated with LWS solution. There was no significant difference in heart rate values among all three groups. LWS solution treatment adversely affected coagulation ($p<0.01$) whereas no effect was recorded with HSD and whole blood treatments. The partial pressure of blood gases and oxygen saturation did not show any significant difference in all three groups. Base excess has not been found to be different in terms of pre- and post-treatment values. Metabolic acidosis was observed in HSD treatment group. Lactate was detected to be statistically lower in rats treated with whole blood ($p=0.03$).

Conclusion: Use of combination of small volume of HSD and whole blood as fluid resuscitation may exert a favorable extended profile of hemodynamic effects, better hemostasis and tissue perfusion in hemorrhagic shock.

Key words: coagulation, asid-base disorders, mean arterial pressure, whole blood, hydroxethyl starch , hypertonic saline

¹Haydarpaşa Numune Research and Education Hospital, Department of Anesthesiology Haydarpaşa, 34668 Istanbul, Turkey

²Haydarpaşa Numune Research and Education Hospital, Biochemistry Clinics Haydarpaşa, 34668 Istanbul, Turkey

³Haydarpaşa Numune Research and Education Hospital, Department of Neurosurgery Haydarpaşa, 34668 Istanbul, Turkey

⁴ Marmara University, Department of Pharmacology and Clinical Pharmacology, School of Medicine, Haydarpaşa, 34668 Istanbul, Turkey

Kanama ile Hipovolemi Oluşturulan Sıçanlarda Resüsitasyon Sıvılarının Hemodinamik Parametreler ve Kan Biyokimyası Üzerine Etkileri

Amaç: Hemorajik şok nadir rastlanılan ancak hemodinamik bozukluk, oksijen taşınımında ve doku perfüzyonunda azalma, hücrel hipoksi, organ hasarı ve ölüme neden olan ciddi bir komplikasyondur. Bu çalışma, kanama ile hipovolemi oluşturulan sıçanlarda tam kan, düşük moleküler ağırlıklı nişasta solüsyonu ve dekstranlı hipertonic saline solüsyonlarının hemodinamik parametreler, koagülasyon ve doku perfüzyonu üzerine olan etkilerini araştırmak amacıyla tasarlanmıştır.

Metodlar: Deneylerde 200–250 g Sprague Dawley sıçanlar (her grupta n=7) kullanılmıştır. Sıçanlar, sol iliak arterden 20 dakikalık üç birbirini takip eden basamakla kanama yapılarak ortalama kan basıncı yaklaşık 20 mmHg da stabilize edilene kadar düşürülmüştür. Gruplardaki sıçanlara sonrasında dekstranlı hipertonic saline solüsyon (4 ml/kg; i.v.), düşük moleküler ağırlıklı nişasta solüsyonu (4 ml/kg; i.v.) ya da saklanmış tam kan transfüzyonu (2 ml/100 g; i.v.) 30 dakika süresince uygulanmıştır.

Bulgular: Tam kan ya da dekstranlı hipertonic saline solüsyonu uygulanan sıçanlardaki ortalama kan basıncı düşük moleküler ağırlıklı nişasta solüsyonu uygulanan sıçanların ortalama kan basıncından yüksek bulunmuştur. Gruplar arasında kalp hızı değerleri arasında anlamlı bir farklılık görülmedi. Düşük moleküler ağırlıklı nişasta solüsyonu koagülasyonu olumsuz yönde etkilerken ($p<0.01$), diğerlerinde böyle bir etki görülmedi. Kan gazlarını parsiyel basıncı ve oksijen saturasyonu gruplar arasında farklılık göstermedi. Baz fazlalığı tedavi öncesi ve sonrasında farklı bulunmadı. Dekstranlı hipertonic saline solüsyonu grubunda metabolik asidoz görüldü. Tam kan uygulanan sıçanlarda laktat düzeyi anlamlı olarak düşük bulundu ($p=0.03$).

Sonuç: Dekstranlı hipertonic saline solüsyonu ve tam kan kombinasyonu daha olumlu geniş profilli hemodinamik etkiler sunmakta, hemorajik şokta daha iyi hemostaz ve daha iyi doku perfüzyonu sağlamaktadır.

yonu sağlamaktadır.

Anahtar kelimeler: Koagülasyon, asit-baz bozuklukları, ortalama kan basıncı, tam kan, nişasta solüsyonu, hipertonic saline

INTRODUCTION

Hemorrhagic shock is inadequate tissue perfusion and inadequate removal of cellular waste products causing failure of oxidative metabolism that can involve decreases in oxygen delivery and usage¹. The type of fluid, the dose and the infusion rate are critical issues while treating hemorrhagic shock that define the therapeutic endpoints². The efficacy of fluid resuscitation after injury can be assessed by many criteria including recovery of cardiovascular instability, restoration of organ perfusion and duration of hemodynamic effects³. Crystalloid solutions and blood transfusion are the mainstays of pre-hospital and in-hospital treatment of hemorrhagic shock where in the pre-hospital setting, four types of fluid are recommended: crystalloid solutions, colloid solutions, hypertonic saline and oxygen-carrying blood substitutes^{4,5}.

The concept of resuscitation with the use of a small volume (4 to 6 mL/kg) of very hypertonic (7.5%) sodium chloride solution has been described in experimental and clinical conditions involving hypovolemia. Small volume hypertonic saline infusions rapidly increase cardiovascular and metabolic functions by producing plasma volume expansion through the displacement of intracellular and interstitial fluid to the vascular compartment⁶. The use of hypertonic saline (6%) with dextran-70 (HSD), which is hypertonic and hyperoncotic, is a new approach for intravascular volume replacement since it prolongs volume expansion through an endogenous fluid redistribution⁶. Colloids considerably reduce total fluid requirement, prolong volume expansion and obtain hemodilution with increased microvascular perfusion⁷.

It has been reviewed that whole fresh blood is preferred to resuscitate the patients with acidosis, hypothermia or coagulopathy⁵. Massive blood transfusion is also recommended in hemody-

dynamic resuscitation without reaching a particular hemoglobin or hematocrit target level⁸.

In this current work, we aimed to investigate the effects of low molecular weight starch solution (LWS), hypertonic saline with dextrane (HSD) or transfusion on systemic hemodynamic recovery, coagulation, acid-base status and arterial blood gas parameters in rats bled to hypovolemia.

MATERIALS AND METHODS

Animals

Experiments were performed with male albino Sprague Dawley rats, weighing 200–250 g. An approval of Institutional Animal Care and Use Committee was taken before the experiments. The animals were kept in a temperature-controlled room with 12-h light and dark cycle and fed with standard animal food and water ad libitum. Experiments were performed under urethane (1.2 g/kg, i.p.) anesthesia. Normal body temperature was maintained by continuous monitoring via a rectal thermometer and a heating pad during the experiments.

Direct measurement of blood pressure

Bilateral iliac arteries were catheterized with a PE-10 catheter attached to PE-50 polyethylene tubing, one for direct measurement of blood pressure and the other for bleeding. Arterial blood pressure was recorded on a polygraph (Grass Model 7, USA) via a pressure transducer (Grass). The heart rate was monitored via a tachograph (Grass Model 7P44D, USA). The left iliac vein was also catheterized for intravenous administration of drug solutions. All catheters were filled with 1% heparin–saline solution. Heparin-saline solutions were discarded before collection of blood samples.

Experimental protocol

Upon completion of catheterisation the rats were connected to polygraph and blood pressure and heart rate was monitored for 10 min. Then, the rats were bled through left iliac artery for 20 min in three subsequent steps until the mean arterial

pressure fell to and stabilized at approximately 20 mm Hg. A total of 1.99 ± 0.1 ml blood/100 g was withdrawn that is approximately equivalent to 50% of the total blood volume in rats⁹.

Either hypertonic saline with dextrane (4 ml/kg; i.v.) or low molecular weight starch solution (4 ml/kg; i.v.) was injected or stored whole blood transfusion (2 ml/100 g; i.v.) was performed after the stabilization of mean arterial pressure (MAP) at around 20 mm Hg. The blood withdrawn for inducing hypovolemia was used for biochemical assays or collected in citrated tubes (containing 1 mg citrate) for transfusion. The rats were monitored continuously 30 min and final blood samples (3 ml) were taken before sacrifice. The coagulation tests were measured with MDA[®], microbial analyzing system (Trinity Biotech, USA) and partial pressures of gases in blood are measured by Radiometer model ABL825 (Radiometer Medical APS, Copenhagen).

Drugs

Low molecular weight starch solution (LWS solution; %6 HydroxyEthylStarch 130/0.4 in 0.9% sodium chloride solution, Voluven[®], Fresenius Kabi, Canada) and hypertonic saline with dextrane (HSD; 7.5% hypertonic saline + 6% dextrane70, RescueFlow[®], BioPhausia AB, Sweden) were used in the study. The blood withdrawn for inducing hypovolemia was stored and used for transfusion.

Data analysis

The results were expressed as “mean \pm standard error of mean (sem)”. Mean arterial pressure was calculated as “1/3 pulse pressure+diastolic blood pressure”. Analysis of variance for repeated measures (two-way) and Bonferroni post test were used to analyse the effect of solutions on mean arterial blood pressure or heart rate. Paired Student's t-test was used to compare pre- and post-bleeding biochemical values. One-way analysis of variance followed by Dunnet's post test was used for comparing the post-treatment biochemical values in different groups receiving different treatments. The level of statistical significance was accepted as $P < 0.05$.

RESULTS

The effect of whole blood, HSD and LWS treatments on MAP has been found to produce a significant time course difference ($p < 0.0001$; $F = 35.75$, $df = 6$) as shown in Fig.1 where the treatment blocks were found to be different ($p = 0.0007$; $F = 11.24$, $df = 2$) and a significant interaction was detected ($p < 0.0001$; $F = 6.32$, $df = 1$). In rats treated either with whole blood or HSD, the MAP values were found to be higher than the rats treated with LWS. Neither of the solutions displayed any quick restoration of MAP values within 10 min like whole blood. There was no significant difference in heart rate values among all three groups (Fig. 2).

When the blood chemistry was investigated, the groups did not produce a significant difference in terms of pre-treatment values of three treatment approaches. LWS treatment prolonged international normalized ratio (INR; $p = 0.0062$) and LWS treatment was found to be different HSD and whole blood treatments in terms of coagulation

($p < 0.01$; Table1).

Base excess has not been found to be different in terms of pre- and post-treatment values or treatment groups (Table.1). According to the pH values listed in Table.1, we observed metabolic acidosis in HSD treatment group ($p = 0.0292$). Lactate was detected to be lower in rats treated with whole blood ($p = 0.03$). Whole blood increased hemoglobin values when compared to HSD ($p < 0.01$) and LWS ($p < 0.05$).

The comparison of partial pressure of blood gases and oxygen saturation did not produce any significant difference both in the pre-treatment and post-treatment values in three resuscitation fluids (Table 2).

Hemoglobin values increased from 14.44 ± 0.59 to 15.99 ± 0.76 in rats treated with whole blood ($p = 0.0014$) but decreased in HSD or LWS treated rats ($p = 0.0734$ and $p = 0.0095$, respectively).

DISCUSSION

In hemorrhagic shock, the main management strategies are the control of bleeding and the




Figure Legends.

Fig. 1 The effect of whole blood, hypertonic saline-dextrane 70 (HSD) and low molecular weight starch solution (LSW) on mean arterial pressure in rats bled to hemorrhagia ($n = 7$ per group). Treatments were given at time 0.

*, $p < 0.05$ (Bonferroni post-test, compared to LWS).

** , $p < 0.01$ (Bonferroni post-test, compared to LSW group).




Fig. 2 The effect of whole blood, hypertonic saline-dextrane 70 (HSD) and low molecular weight starch solution (LSW) on heart rate in rats bled to hemorrhagia ($n = 7$ per group). Treatments were given at time 0.

replacement of circulating volume¹⁰. The critical problems of hemorrhagic shock are decreased tissue nutrition and drainage of metabolites into tissues thus producing subsequent hypoxic cellular damage¹. It was accepted that trauma victims in hypotensive hemorrhage should receive large volumes of fluids as early as possible for increasing cardiac output and oxygen delivery to maintain microvascular perfusion and oxygenation⁴. Correction of the deficit in blood volume with crystalloid volume expanders will generally maintain hemodynamic stability, while transfusion of red cells is used to improve and maintain tissue oxygenation. The main target of blood transfusions in hemorrhagic shock is to restore the capacity of intravascular volume for oxygen transport¹¹.

The use of hypertonic saline was introduced by Velasco and colleagues in 1980¹². Using small-volume hypertonic solutions for resuscitation have some advantages in improving microvascular flow, controlling intracranial pressure, stabilizing arterial pressure and cardiac output^{4,13}. However, its potent and rapid hemodynamic effects only last for 30 to 60 min. Combining dextran or hydroxyethyl starch (HES) with hypertonic saline increases the duration of its effects up to 3-4 h and expands plasma volume 3-4 times^{14,15}. Previously, it was demonstrated that these solution dramatically restore the blood pressure and the cardiac output within 2 min in experimental animals^{15,16}. Colloids are often preferred for correcting hypovolemia, improving systemic and microcirculatory hemodynamics^{17,18}.

In this current study, we demonstrated that the groups receiving whole blood or HSD treatments had more marked elevations in MAP values than the group received LWS solution treatment.

Hemorrhagic coagulopathy is also another major problem that occurs following traumatic injury and underlying mechanism still remains unknown¹⁹. Intravenous fluids can produce adverse effects on hemostatic mechanisms while stabilizing hemodynamics. Dilutional coagulopathy and secondary clot disruption due to increased blood flow, increased perfusion pressure and

decreased blood viscosity are the main reasons for hemorrhagic coagulopathy^{7,10}. Dilution of plasma clotting proteins results in prolongation of the prothrombin time (PT) and the activated partial thromboplastin time (aPTT) and this requires the monitorisation of PT, aPTT, and fibrinogen in patients receiving massive blood transfusions²⁰. The use of artificial colloids may be associated with untoward effects on coagulation as well. In some studies colloids have been found effective and safe substitutes for blood loss without relevant adverse effects on coagulation, but some showed that it should be avoided especially in patients at risk of increased bleeding²¹. Hydroxyethyl starches (HES) is believed to alter coagulation and platelet function leading to increased bleeding tendency. This effect is caused by physicochemical characteristics of the HES preparation and the electrolyte composition of the solvent^{22,23}. Bleeding tendency is also associated with reduction in circulating concentrations of factor VIII and von Willebrand factor (vWF) by the hydroxyethyl starches²⁴. The effects on blood coagulation closely depends on physicochemical properties of different types of HES. Whereas some studies have showed lower molar substitution of HES (130/0.4) causes lower perioperative blood loss and has minimal coagulation impairment, other studies suggest that all molecular substitutes of HES can alter coagulation^{24,25}.

Fluid resuscitation with hypertonic saline in case of hemorrhagic shock causes delay in clot formation and greater bleeding results²⁷. Hypertonic saline solutions have been investigated in various clinical settings. As a replacement fluid, it is cheaper, free of risk of infection or allergic reactions compared with colloids. In current study, HES statistically prolonged INR when compared to HSD and whole blood ($p < 0.05$). Fluid resuscitation with HES caused coagulation impairment. Large amounts of fluid replacement for hemorrhagic shock especially associated with untoward changes of acid-base status²⁵. pH is usually used as an index for tissue perfusion and it has been verified in a variety of patients and in experimental hemorrhagic shock models²⁸. Base ex-

cess is an important marker to identify patients with under-perfused tissues²⁹. During hemorrhagic shock, metabolic acidosis is common and hyperlactatemia is the main cause. The most sensitive marker which predicts complications in trauma patients is arterial base deficit and is correlated with arterial lactate concentration. Elevated base deficit and lactate concentrations after shock are related to oxygen transport imbalance at the cellular level³⁰. Lactate and base deficit are all more sensitive endpoints of cellular resuscitation³¹. In our study, we demonstrated that lactate decreased significantly in rats treated with whole blood ($p < 0.05$). HSD didn't recover metabolic acidosis when compared to LWS and whole blood administration due to its hypernatremic and hyperchloremic effects. P02, PCO2 and SPO2 parameters didn't show statistically significant difference among all three group ($p > 0.05$).

CONCLUSION

The primary treatment of hemorrhagic shock is control of the source of bleeding as soon as possible and fluid replacement. There are some concerns regarding the best method for resuscitation including the choice of fluid type, its amount, its application time and rate. The results of our study demonstrate that hypertonic saline solution improves hemodynamic status and has beneficial effects on microcirculation in hemorrhagic shock without causing significant side effects. LWS solution may have negative effects on hemostasis. Thus, HSD and whole blood combination is a good approach for fluid resuscitation in massive hemorrhagic shock.

REFERENCES

1. Andrew P, Geoffrey F, David EL. Fluid Resuscitation. In: State of the Science for Treating Combat Casualties and Civilian Injuries, National Academy Press, Washington DC; 1999. pp 19.
2. Guillermo G, David R, Marian EW. Clinical review: Hemorrhagic shock. *Crit Care Med* 2004; 8:373-381.
3. Dubick MA, Atkins JL. Small-volume fluid resuscitation for the far-forward combat environment: current concepts. *J Trauma* 2003; 54:43-45.
4. Michael MK. Initial resuscitation of hemorrhagic shock. *World J Emerg Surg* 2006; 1:14.
5. Repine TB, Perkins JG, Kauvar DS, Blackburne L. The use of fresh whole blood in massive transfusion. *J Trauma* 2006;60:59-69.
6. Braz JR, do Nascimento PJr, Paiva Filho O et al. The Early Systemic and Gastrointestinal Oxygenation Effects of Hemorrhagic Shock Resuscitation with Hypertonic Saline and Hypertonic Saline 6% Dextran-70: A Comparative Study in Dogs. *Anesth Analg* 2004; 99:536-546.
7. Antonino Gullo Philip DL, José B, Ged FW. Intensive and Critical Care Medicine. In: Plasma Volume Expansion: The Current Controversy, Springer, Milan; 2009. pp 151-161.
8. Riskin DJ, Tsai TC, Riskin L et al. Massive transfusion protocols: the role of aggressive resuscitation versus product ratio in mortality reduction. *J of the Am Coll of Surg* 2009; 209:198-205.
9. Butcher HR Jr. Measured blood volumes as the basis for the extent of hemorrhage. Joseph RS, Amanda BA, Jessica LZ et al. Fluid Resuscitation Therapy for Hemorrhagic Shock. *J Trauma Nurs* 2007; 14:152-160.
11. Ribeiro MA, Jr Epstein MG, Alves LD. Volume replacement in trauma. *Ulus Travma Acil Cerrahi Derg* 2009; 15:311-316.
12. Velasco IT, Pontieri V, Rocha e Silva MJr. Hyperosmotic NaCl and severe hemorrhagic shock. *Am J Physiol* 1980; 239:664-673.
13. Nakayama S, Sibley L, Gunther RA et al. Small-volume resuscitation with hypertonic saline (2,400 mOsm/liter) during hemorrhagic shock. *Circ Shock* 1984; 13:149-159.
14. Kaainen T, Alaoja H, Heikkinen J et al. Hypertonic Saline Dextran Improves Outcome After Hypothermic Circulatory Arrest: A Study in a Surviving Porcine Model. *Ann Thorac Surg* 2006; 81:183-190.
15. Chiara O, Pelosi P, Brazzi L et al. Resuscitation from hemorrhagic shock: Experimental model comparing normal saline, dextran, and hypertonic saline solutions. *Crit Care Med* 2003; 31:1915-1922.
16. Smith GJ, Kramer GC, Perron P et al. A comparison of several hypertonic solutions for resuscitation of bled sheep. *J Surg Res* 1985; 39:517-528.
17. Martel MJ, MacKinnon KJ, Arsenault MY et al. Clinical Practice Obstetrics Committee and Executive and Council, Society of Obstetricians and Gynaecologists of Canada. Hemorrhagic Shock *J Obstet Gynaecol Can* 2002; 24:504-520.
18. Lang K, Boldt J, Suttner S, Haisch G. Colloids versus crystalloids and tissue oxygen tension in patients undergoing major abdominal surgery. *Anesth Analg* 2001; 93:405-409.
19. Wenjun Z, David L, Michael A. Effects of hemorrhage and lactated ringer's resuscitation on coagulation and fib-

- rinogen metabolism in swine. *Shock* 2006; 26:396-401.
20. Hardy JF, De Moerloose P, Samama CM. Members of the Groupe d'Intérêt en Hémostase Périopératoire. Massive transfusion and coagulopathy: pathophysiology and implications for clinical management. *Can J Anaesth* 2006; 53:40-58.
21. Hüttner I, Boldt J, Haisch G et al. Influence of different colloids on molecular markers of haemostasis and platelet function in patients undergoing major abdominal surgery. *Br J Anaesth* 2000; 85:417-423.
22. Boldt J. The balanced concept of fluid resuscitation. *Br J Anaesth* 2007; 99:312-315.
23. Treib J, Haass A, Pindur G et al. All medium starches are not the same: influence of the degree of hydroxyethyl substitution of hydroxyethyl starch on plasma volume, hemorrheologic conditions, and coagulation. *Transfusion* 1996; 36:450-455.
24. Madjdpour N, Dettori N, Frascarolo P et al. Molecular weight of hydroxyethyl starch: is there an effect on blood coagulation and pharmacokinetics? *Br J Anaesth* 2005; 94:569-576.
25. Sibylle A, Kozek L, Cornelius J et al. The Effects of Hydroxyethyl Starch 130/0.4 (6%) on Blood Loss and Use of Blood Products in Major Surgery: A Pooled Analysis of Randomized Clinical Trials. *Anesth Analg* 2008; 107:382-390.
26. Cabrales P, Tsai AG, Intaglietta M. Resuscitation from hemorrhagic shock with hydroxyethyl starch and coagulation changes. *Shock* 2007; 28:461-467.
27. Tan TS, Tan KH, Ng HP, Loh MW. The effects of hypertonic saline solution (7.5%) on coagulation and fibrinolysis: an in vitro assessment using thromboelastography. *Anaesthesia* 2002; 57:644-648.
28. Braz JR, do Nascimento P Jr, Paiva Filho O et al. The Early Systemic and Gastrointestinal Oxygenation Effects of Hemorrhagic Shock Resuscitation with Hypertonic Saline and Hypertonic Saline 6% Dextran-70: A Comparative Study in Dogs. *Anesth. Analg* 2004; 99:536-546.
29. Brill SA, Stewart TR, Brundage SI, Schreiber MA. Base deficit does not predict mortality when secondary to hyperchloremic acidosis. *Shock* 2002; 17:459-462.
30. Kincaid E. Elevated arterial base deficit in trauma patients: a marker of impaired oxygen utilization. *J Am Coll Surg* 2009; 187:384-392.
31. Gutierrez G, Taylor D. Gastrointestinal tonometry: basic principles and recent advances in monitoring regional CO₂ metabolism. *Sem. Respir Crit Care Med* 1999; 20:17-27.