

Poster Sessions – Abstract P063

Transmission route and reasons for HIV testing among recently diagnosed HIV patients in HIV-TR cohort, 2011–2012

Dokuzoguz, Basak¹; Korten, Volkan²; Gökengin, Deniz³; Fincanci, Muzaffer⁴; Yildirmak, Taner⁵; Nuray Kes, Uzun⁶; Tasdelen Fisgin, Nuriye⁷; Inan, Dilara⁸; Eraksoy, Haluk⁹ and Akalin, Halis¹⁰

¹Infectious Diseases and Clinical Microbiology, Ankara Numune Education Training and Research Hospital, Ankara, Turkey. ²Infectious Diseases, Marmara University Hospital, Istanbul, Turkey. ³Infectious Diseases, Ege University Hospital, Izmir, Turkey. ⁴Infectious Diseases, Istanbul Education and Research Hospital, Istanbul, Turkey. ⁵Infectious Diseases, Okmeydanı Education and Research Hospital, Istanbul, Turkey. ⁶Infectious Diseases, Bıbli Education and Research Hospital, Istanbul, Turkey. ⁷Infectious Diseases, 19 Mayıs University Hospital, Samsun, Turkey. ⁸Infectious Diseases, Akdeniz University Hospital, Antalya, Turkey. ⁹Infectious Diseases, Istanbul Medical School, Istanbul University, Istanbul, Turkey. ¹⁰Infectious Diseases, Uludağ University Hospital, Bursa, Turkey.

Introduction: Routes of transmission and reasons for HIV testing are important epidemiologic data to analyze the epidemic and to tailor the response to AIDS. The aim of this study was to analyze reasons for testing and transmission ways of HIV among recently diagnosed HIV patients registered in the multicenter HIV-TR cohort in Turkey.

Methods: Transmission ways and reasons for testing of all patients diagnosed in 2011 and 2012 were recorded on a web-based data collection system and were analyzed retrospectively.

Results: The study included 693 patients (561 male, 132 female) from 24 sites. Reason for HIV testing was available in 640 patients (92%). The most common reason for HIV testing was diagnostic workout for other conditions or illness followed by patient-initiated testing. The reasons for testing were listed in Table 1. The most common routes of HIV transmission were heterosexual intercourse (62.7%) and sex among men who have sex with men (MSM) (22.6%). At the time of HIV diagnosis, the mean CD4 lymphocyte cell count was 355/mm³ (3–1433/mm³). Primary HIV infection was determined in 42/693 (6%) patients and 9/693 (% 1, 2) cases were considered “probable primary HIV infection.” The majority of the cases presented to a clinic for follow-up right after the diagnosis. On the other hand 32/616 (5.2%) patients delayed their presentation for more than 3 months. The longest delay was 11 months.

Conclusions: The results of the database suggest that targeted testing is lacking in the country. The shift toward homosexual transmission during the last 2 years emphasizes the need for targeted interventions. Patients present relatively late and HIV infection could only be diagnosed when immunosuppression related findings appeared. Patient-initiated testing, an indicator of awareness, was very low suggesting a need to scale-up awareness raising interventions.

Table 1. Reasons for HIV testing

Reason for testing	N	%
Blood/organ donation	63	9.8
Prior to surgical/parenteral intervention	63	9.8
Premarital testing	20	3.1
Screening for pregnancy	11	1.7
Patient-initiated	98	15.3
Diagnostic workout for other medical condition/illness	312	48.8
Job application	8	1.3
Other	65	10.2
Total	640	100

Published 2 November 2014

Copyright: © 2014 Dokuzoguz B et al; licensee International AIDS Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution 3.0 Unported (CC BY 3.0) License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.