



# MIT Open Access Articles

## *Search for New Phenomena in Monophoton Final States in Proton–proton Collisions at $\sqrt{s} = 8$ TeV*

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

<b>Citation</b>	Khachatryan, V., A.M. Sirunyan, A. Tumasyan, W. Adam, T. Bergauer, M. Dragicevic, J. Erö, et al. "Search for New Phenomena in Monophoton Final States in Proton–proton Collisions at $\sqrt{s} = 8$ TeV." Physics Letters B 755 (April 2016): 102–124 © 2016 CERN for the benefit of the CMS Collaboration
<b>As Published</b>	<a href="http://dx.doi.org/10.1016/J.PHYSLETB.2016.01.057">http://dx.doi.org/10.1016/J.PHYSLETB.2016.01.057</a>
<b>Publisher</b>	Elsevier
<b>Version</b>	Author's final manuscript
<b>Citable link</b>	<a href="http://hdl.handle.net/1721.1/116633">http://hdl.handle.net/1721.1/116633</a>
<b>Terms of Use</b>	Creative Commons Attribution 4.0 International License
<b>Detailed Terms</b>	<a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a>

# Accepted Manuscript

Search for new phenomena in monophoton final states in proton–proton collisions at  $\sqrt{s} = 8$  TeV

CMS Collaboration

PII: S0370-2693(16)00076-9  
DOI: <http://dx.doi.org/10.1016/j.physletb.2016.01.057>  
Reference: PLB 31691

To appear in: *Physics Letters B*

Received date: 31 October 2014  
Revised date: 24 January 2016  
Accepted date: 26 January 2016

Please cite this article in press as: CMS Collaboration, Search for new phenomena in monophoton final states in proton–proton collisions at  $\sqrt{s} = 8$  TeV, *Phys. Lett. B* (2016), <http://dx.doi.org/10.1016/j.physletb.2016.01.057>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



# Search for new phenomena in monophoton final states in proton-proton collisions at $\sqrt{s} = 8$ TeV

The CMS Collaboration<sup>a</sup>

<sup>a</sup>CERN

## Abstract

Results are presented from a search for new physics in final states containing a photon and missing transverse momentum. The data correspond to an integrated luminosity of  $19.6 \text{ fb}^{-1}$  collected in proton-proton collisions at  $\sqrt{s} = 8$  TeV with the CMS experiment at the LHC. No deviation from the standard model predictions is observed for these final states. New, improved limits are set on dark matter production and on parameters of models with large extra dimensions. In particular, the first limits from the LHC on branon production are found and significantly extend previous limits from LEP and the Tevatron. A cross section upper limit of  $14.0 \text{ fb}$  is set at the 95% confidence level for events with a monophoton final state with photon transverse momentum greater than 145 GeV and missing transverse momentum greater than 140 GeV.

**Keywords:** CMS, physics, monophoton, dark matter

## 1. Introduction

The production of events containing photons with large transverse momentum and having large missing transverse momentum at the CERN LHC is sensitive to physics beyond the standard model (SM). In this Letter we investigate three possible extensions of the SM: a model incorporating pair production of dark matter (DM) particles, and two models with extra spatial dimensions, as described below.

At the LHC, DM particles ( $\chi$ ) [1] can be produced in the process  $q\bar{q} \rightarrow \gamma\chi\bar{\chi}$ , where the photon is radiated by one of the incoming quarks. With a photon in the final state, we gain sensitivity to the production of invisible particles. The SM-DM interaction is assumed to be mediated by a virtual particle (“mediator”) with a mass  $M$  much heavier than the fermionic DM particle mass ( $M_\chi$ ). Various processes are contracted into an effective field theory (EFT) [2–5], assuming  $M$  much larger than the momentum transfer scale  $Q$  (i.e.  $M \gg Q$ ) and a contact interaction scale  $\Lambda$  given by  $\Lambda^{-2} = g_\chi g_q M^{-2}$ , where  $g_\chi$  and  $g_q$  are the mediator couplings to  $\chi$  and to quarks, respectively. Using this formalism, results from searches at the LHC can be related to limits for direct searches sensitive to  $\chi$ -nucleon scattering [5].

The ADD model [6, 7] of large extra dimensions is postulated to have  $n$  extra compactified spatial dimensions at a characteristic scale  $R$  that reflects an effective Planck scale  $M_D$  through  $M_{\text{Pl}}^2 \approx M_D^{n+2} R^n$ , where  $M_{\text{Pl}}$  is the Planck scale. If  $M_D$  is of the same order as the electroweak scale ( $M_{\text{EW}} \sim 10^2 \text{ GeV}$ ), the large value of  $M_{\text{Pl}}$  can be interpreted as being a consequence of large-volume ( $\sim R^n$ ) suppression from extra dimensional space. This model predicts a sizable cross section for the process  $q\bar{q} \rightarrow \gamma G$ , where  $G$  is a graviton that escapes detection, and motivates the search for events with a single  $\gamma$  and missing transverse momentum.

In both the ADD and branon models, the SM particles are constrained to live on a 3+1 dimensional 3-brane surface. In the branon family of models [8–11], it is assumed that the brane fluctuates in the extra dimensions, in contrast to the ADD model, where the brane is rigid. In this alternative scheme, the brane tension scale  $f$  is expected to be much smaller than other relevant scales such as  $M_D$ . The particles associated with such fluctuations are scalar particles called branons. Branons are stable and massive scalar particles of mass  $M_B$ , and are natural candidates for dark matter [12]. They can be pair-produced in association with SM particles at the LHC, giving rise to  $\gamma$ +missing transverse momentum final states [13]. If  $N$  extra dimensions are considered, then  $N$  branons are expected and their production cross

<sup>a</sup>Email address:

cms-publication-committee-chair@cern.ch (The CMS Collaboration)

section scales with  $N$ . In the following, only the  $N = 1$  case is considered.

The primary background to the  $\gamma$ +missing transverse momentum signal is the irreducible SM background from  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$  production. Other backgrounds include  $W\gamma \rightarrow \ell\nu\gamma$  (where  $\ell$  is an undetected charged lepton),  $W \rightarrow e\nu$  (where the electron is misidentified as a photon),  $\gamma$ +jet, QCD multijet (with a jet misidentified as a photon),  $Z\gamma \rightarrow \ell\ell\gamma$ , and diphoton events, as well as backgrounds from beam halo.

## 2. The CMS detector

The CMS experiment uses a right-handed coordinate system, with the origin at the nominal interaction point, the  $x$  axis pointing to the center of the LHC, the  $y$  axis pointing up (perpendicular to the LHC plane), and the  $z$  axis along the anticlockwise-beam direction. The azimuthal angle  $\phi$  is measured from the  $x$ -axis in the  $x$ - $y$  plane and the polar angle  $\theta$  is measured from the  $z$ -axis. Pseudorapidity is defined as  $\eta = -\ln[\tan(\theta/2)]$ .

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the superconducting solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel ( $|\eta| < 1.479$ ) and two endcap ( $1.479 < |\eta| < 3.0$ ) sections. Electrons are found by associating clusters of ECAL energy with adjacent tracker hits. Muons are detected in the pseudorapidity range  $|\eta| < 2.4$ , using gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid, and reconstructed from tracks in these detectors combined with those from the silicon tracker. Extensive forward calorimetry ( $3.15 < |\eta| < 4.9$ ) complements the coverage provided by the barrel and endcap detectors. The energy resolution for photons with transverse momentum  $\geq 60$  GeV varies between 1.1% and 2.6% over the solid angle of the ECAL barrel, and from 2.2% to 5.0% in the endcaps [14]. The timing measurement of the ECAL has a resolution better than 200 ps for energy deposits larger than 10 GeV [14]. In the  $\eta$ - $\phi$  plane, and for  $|\eta| < 1.48$ , the HCAL cells map onto  $5 \times 5$  arrays of ECAL crystals to form calorimeter towers projecting radially outward from the nominal interaction point. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Ref [15].

## 3. Event selection

In the following, it is convenient to refer to the missing transverse momentum vector,  $\vec{E}_T^{\text{miss}}$ , defined as the projection on the plane perpendicular to the beams of the negative vector sum of the momenta of all reconstructed particles in an event. Its magnitude is referred to as  $E_T^{\text{miss}}$ .

Events are selected from a data sample corresponding to an integrated luminosity of  $19.6 \text{ fb}^{-1}$  collected in proton-proton collisions at  $\sqrt{s} = 8 \text{ TeV}$  with the CMS experiment at the LHC. Triggers requiring at least one electromagnetic cluster or a cluster along with large  $E_T^{\text{miss}}$  are used. For the selected signal region of transverse energy  $E_T^{\gamma} > 145 \text{ GeV}$ , pseudorapidity  $|\eta^{\gamma}| < 1.44$ , and  $E_T^{\text{miss}} > 140 \text{ GeV}$ , these triggers are  $\approx 96\%$  efficient for  $E_T^{\gamma}$  in the 145–160 GeV range, and fully efficient for  $E_T^{\gamma} > 160 \text{ GeV}$ . Events are required to have at least one primary vertex reconstructed within a longitudinal distance of  $|z| < 24 \text{ cm}$  of the center of the detector and at a distance  $< 2 \text{ cm}$  from the  $z$ -axis. The primary vertex is chosen to be the vertex with the highest sum in  $p_T^2$  of its associated tracks, where  $p_T$  is the transverse momentum.

Candidate electromagnetic (EM) showers are restricted to the barrel region of the ECAL, where their purity is highest [16]. Photon candidates [17] are selected by requiring the ratio of the energy deposited in the closest HCAL tower to the energy of the EM showers in the ECAL to be less than 0.05 and the spatial distribution of energy in the EM shower to be consistent with that expected for a photon. In order to reject hadronic activity, photon candidates are required to be isolated, using the sum of the transverse energy of additional particles within a cone of  $\Delta R < 0.3$  centered on the shower axis, where  $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ , reconstructed using a particle-flow algorithm [18, 19]. In this isolation cone, the sum of the transverse energy (in GeV) of additional photons is required to be less than  $(0.7 + 0.005E_T^{\gamma})$ , of neutral hadrons is required to be less than  $(1.0 + 0.04E_T^{\gamma})$ , and of charged hadrons is required to be less than 1.5. The charged hadron contribution includes that calculated from the other interaction vertices in the event (pileup), arising from the uncertainty in assigning the photon candidate to a particular vertex. The effect of pileup on the isolation variables is mitigated using the scheme presented in Ref. [20].

The ECAL crystal containing the highest energy within the cluster of the photon candidate is required to have a time of deposition within  $\pm 3 \text{ ns}$  of particles arriving from the collision. This selection suppresses contributions from noncollision backgrounds. To reduce contamination from beam halo, the crystals (excluding

those associated with the photon candidate) are examined for evidence of the passage of a minimum-ionizing particle roughly parallel to the beam axis (beam halo tag). If sufficient energy is found along such a trajectory, the event is rejected. Highly ionizing particles traversing the sensitive volume of the readout photodiodes can give rise to spurious signals within the EM shower [21]. These EM showers are eliminated by requiring consistency among the timings of energy depositions in all crystals within the shower. Photon candidates are rejected if they are likely to be electrons, as inferred from characteristic patterns of hits in the pixel detector, called “pixel seeds”, that are matched to candidate EM showers [22].

Jets are reconstructed with the anti- $k_T$  algorithm [23] using a radius parameter of  $R = 0.5$ . Jets that are identified as arising from pileup are rejected [24]. In order to reduce QCD multijet backgrounds, events are rejected if there is more than one jet with  $p_T > 30$  GeV at  $\Delta R > 0.5$  relative to the photon. Events with isolated leptons (electron or muon) with  $p_T > 10$  GeV,  $|\eta| < 2.4$  (2.5) for muons (electrons) and  $\Delta R > 0.5$  relative to the photon, are also rejected to suppress  $W\gamma \rightarrow \ell\nu\gamma$  and  $Z\gamma \rightarrow \ell\ell\gamma$  backgrounds. Lepton isolation is computed using the sum of transverse energies of tracks, ECAL, and HCAL depositions within a surrounding cone of  $\Delta R < 0.3$ . For electron isolation, each contributing component of transverse energy (tracker, ECAL, and HCAL) is required to be less than 20% of the electron  $p_T$ , while for muons only the tracker component is considered and is required to be less than 10% of the muon  $p_T$ .

The candidate events are required to have  $E_T > 140$  GeV. A topological requirement of  $\Delta\phi(\vec{E}_T, \gamma) > 2$  rad is applied to suppress the contribution from the  $\gamma$ +jet background.

A major source of background comes from events with mismeasured  $E_T$  due to finite detector resolution, mainly associated with jets. In order to reduce the contribution from events with mismeasured  $E_T$ , for each event a  $\chi^2$  function is constructed and minimized:

$$\chi^2 = \sum_i \left( \frac{(p_T^{\text{reco}})_i - (\bar{p}_T)_i}{(\sigma_{p_T})_i} \right)^2 + \left( \frac{\tilde{E}_x}{\sigma_{\tilde{E}_x}} \right)^2 + \left( \frac{\tilde{E}_y}{\sigma_{\tilde{E}_y}} \right)^2, \quad (1)$$

where the summation is over the reconstructed particles, i.e., the photon and the jets. In the above equation,  $(p_T^{\text{reco}})_i$  are the transverse momenta, and the  $(\sigma_{p_T})_i$ , the expected momentum resolutions of the reconstructed particles. The  $(\bar{p}_T)_i$  are the free parameters allowed to vary in order to minimize the function. The resolution parametrization associated with the  $E_T$  is obtained from

Ref [25]. Lastly,  $\tilde{E}_x$  and  $\tilde{E}_y$  can be expressed as:

$$\begin{aligned} \tilde{E}_{x,y} &= E_{x,y}^{\text{reco}} + \sum_{i=\text{objects}} (p_{x,y}^{\text{reco}})_i - (\bar{p}_{x,y})_i \\ &= - \sum_{i=\text{objects}} (\bar{p}_{x,y})_i \end{aligned} \quad (2)$$

In events with no genuine  $E_T$ , the mismeasured quantities will be more readily re-distributed back into the particle momenta, which will result in a low  $\chi^2$  value. On the other hand, in events with genuine  $E_T$  from undetected particles, minimization of the  $\chi^2$  function will be more difficult and generally will result in larger  $\chi^2$  values. To reduce the contribution of events with mismeasured  $E_T$ , the probability value obtained from the  $\chi^2$  minimization is required to be smaller than  $10^{-6}$  and  $\tilde{E}_T = \sqrt{\tilde{E}_x^2 + \tilde{E}_y^2}$ , in which the original reconstructed particle momenta are replaced with those obtained with the  $\chi^2$  minimization, is required to be greater than 120 GeV. These requirements are optimized using the significance estimator  $S/\sqrt{S+B}$  and remove 80% (35%) of  $\gamma$ +jet (QCD multijet) events, while keeping 99.5% of signal events.

After applying all selection criteria, 630 candidate events remain in the sample.

#### 4. Background determination

Backgrounds from  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$ ,  $W\gamma \rightarrow \ell\nu\gamma$ ,  $\gamma$ +jet,  $Z\gamma \rightarrow \ell\ell\gamma$ , and diphoton production are estimated from simulated samples processed through the full GEANT4-based simulation of the CMS detector [26, 27], trigger emulation, and the same event reconstruction programs as used for data. The  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$  and  $W\gamma \rightarrow \ell\nu\gamma$  samples are generated with MADGRAPH 5v1.3.30 [28], and the cross section is corrected to include next-to-leading-order (NLO) effects through an  $E_T^\gamma$  dependent correction factor calculated with MCFM 6.1 [29]. The central values of the NLO cross section and the prediction for the photon  $E_T$  spectrum are calculated following the prescriptions of the PDF4LHC Working Group [30–32]. This prescription is also used to calculate the systematic uncertainties due to the parton distribution functions (PDF), and the strong coupling  $\alpha_s$  and its dependence on the factorization scale and renormalization scale. The systematic uncertainties in the NLO cross sections are found to be in the range 8% to 48% and 16% to 82% for  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$  and  $W\gamma \rightarrow \ell\nu\gamma$ , respectively, over the  $E_T^\gamma$  spectrum from 145 GeV to 1000 GeV. The strong correlation in the uncertainties of the two channels is

propagated to the final result. The  $Z\gamma \rightarrow \ell\ell\gamma$  sample is obtained using the MADGRAPH 5v1.3.30 generator [28]. The  $\gamma$ +jet and diphoton samples are obtained using the PYTHIA 6.426 generator [33] at leading order (LO), with the CTEQ6L1 [34] PDF. The  $\gamma$ +jet cross section is corrected to include NLO effects.

The backgrounds estimated from simulations are scaled by a factor  $F$  to correct for observed differences in efficiency between data and simulation. This overall data/simulation correction factor receives contributions from four sources as follows: the photon reconstruction efficiency ratio, estimated to be  $0.97 \pm 0.02$  using  $Z \rightarrow ee$  decays; the ratio of probabilities for satisfying a crystal timing requirement, estimated to be  $0.99 \pm 0.03$  from a sample of electron data; the lepton veto efficiency ratio, estimated to be  $0.99 \pm 0.02$  using  $W \rightarrow e\nu$  decays; and the jet veto efficiency ratio, estimated to be  $0.99 \pm 0.05$  using  $W \rightarrow e\nu$  decays, and confirmed using  $Z\gamma \rightarrow ee\gamma$  data samples. The total correction factor obtained by combining these contributions is  $F = 0.94 \pm 0.06$ .

The total uncertainty in the backgrounds estimated through simulation includes contributions from the theoretical cross section, data-simulation factor  $F$ , pileup modeling, and the accuracy of energy calibration and resolution for photons [14], jets [35], and  $\cancel{E}_T$  [36]. The estimated contribution from the  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$  and  $W\gamma \rightarrow \ell\nu\gamma$  processes to the background are, respectively,  $345 \pm 43$  and  $103 \pm 21$  events, where the dominant uncertainty is from the theoretical cross section calculations. To gain confidence in the estimates from simulation, control regions, which are dominated by these backgrounds and have negligible contributions from a signal, are defined in the data. As a crosscheck, the total contribution from  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$  is estimated in data using a sample of  $Z\gamma \rightarrow \mu\mu\gamma$  candidates, where the muons from the decay of the  $Z$  boson are considered as invisible particles hence contributing to  $\cancel{E}_T$  [37]. The normalization is corrected both for the ratio of the branching fractions of  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$  and  $Z\gamma \rightarrow \mu\mu\gamma$ , and for differences in the acceptance and selection efficiencies. This crosscheck provides an estimate of  $341 \pm 50$  events, where the uncertainty is dominated by the size of the sample. A control region dominated by the  $W\gamma$  process is also studied by using the signal selection but inverting the lepton veto i.e., the final state is required to contain a reconstructed charged lepton. After this selection, 104 events are observed and  $126 \pm 23$  are expected.

Electrons misidentified as photons arise mainly from highly off-shell  $W$  boson ( $W^* \rightarrow e\nu$ ) events. These backgrounds are inclusively estimated from data. The efficiency,  $\epsilon_{\text{pix}}$ , of matching electron showers in the calorimeter to pixel seeds is estimated using a tag-and-

probe technique [38] on  $Z \rightarrow ee$  events in data, verified with simulated events. The efficiency is found to be  $\epsilon_{\text{pix}} = 0.984 \pm 0.002$  for electrons with  $E_T > 100$  GeV. A control sample of  $W^* \rightarrow e\nu$  events is also obtained from data through use of all the standard candidate selections, with the exception of the pixel seed, which is inverted. The number of events in this sample is scaled by the value of  $(1 - \epsilon_{\text{pix}})/\epsilon_{\text{pix}}$  resulting in an inclusive estimate of  $60 \pm 6$   $W^* \rightarrow e\nu$  events in the signal region.

The contamination from jets misidentified as photons is estimated in data using a control sample with  $\cancel{E}_T < 30$  GeV, dominated by QCD events. This sample is used to measure the ratio of the number of objects that pass photon identification criteria to the number that fail at least one of the isolation requirements. The control sample also contains objects from QCD direct photon production that must be removed from the numerator of the ratio. This contribution is estimated by fitting the shower shape distribution with template distributions. For true photons, a template for the shower width is formed using simulated  $\gamma$ +jets events. For jets misidentified as photons, the template is formed using a separate control sample, where the objects are required to fail charged hadron isolation. This corrected ratio is used to scale a set of data events that pass the denominator selection of the fake ratio and all other candidate requirements, providing an inclusive estimate for all backgrounds in which jets are misidentified as photons of  $45 \pm 14$  events.

Noncollision backgrounds are estimated from data by examining the shower width of the EM cluster and the time-of-arrival of the signal in the crystal containing the largest deposition of energy. Templates for anomalous signals, cosmic ray muons, and beam halo events are obtained by inverting the shower shape and beam halo tag requirements, and are fitted to the timing distribution of the candidate sample. The only nonnegligible residual contribution to the candidate sample is found to arise from the beam halo, with an estimated  $25 \pm 6$  events.

## 5. Results

Table 1 shows the estimated number of events and associated uncertainty from each background process along with the total number of events observed in the data, for the entire data set, which corresponds to  $19.6 \text{ fb}^{-1}$ . The number of events observed in data agrees with the expectation from SM background. The photon  $E_T$  and  $\cancel{E}_T$  distributions for the selected candidates and estimated backgrounds are shown in Fig. 1. The spectra expected from the ADD model for  $M_D = 2$  TeV and

$n = 3$  are also shown for comparison. Limits are set for the DM, ADD, and branon models using the  $E_T^\gamma$  spectrum.

Table 1: Summary of estimated backgrounds and observed total number of candidates. Backgrounds listed as ‘‘Others’’ include the small contributions from  $W \rightarrow \mu\nu$ ,  $W \rightarrow \tau\nu$ ,  $Z\gamma \rightarrow \ell\ell\gamma$ ,  $\gamma\gamma$ , and  $\gamma$ +jet. Uncertainties include both statistical and systematic contributions, and the total systematic uncertainty includes the effect of correlations in the individual estimates.

Process	Estimate
$Z(\rightarrow \nu\bar{\nu}) + \gamma$	$345 \pm 43$
$W(\rightarrow \ell\nu) + \gamma$	$103 \pm 21$
electron $\rightarrow \gamma$ MisID	$60 \pm 6$
jet $\rightarrow \gamma$ MisID	$45 \pm 14$
Beam halo	$25 \pm 6$
Others	$36 \pm 3$
Total background	$614 \pm 63$
Data	630

Table 2: Observed (expected) 95% CL and 90% CL upper limits on  $\sigma A$  as a function of the cut on the  $E_T^\gamma$  for the photon and  $\cancel{E}_T$  final state. The  $\cancel{E}_T$  threshold is fixed at 140 GeV. In addition to 95% CL upper limits, 90% limits are also shown to allow direct comparison with results from astrophysics DM searches.

$E_T^\gamma$ Threshold [GeV]	$\sigma A$ [fb]	$\sigma A$ [fb]
	(95% CL)	(90% CL)
145	14 (13)	12 (11)
160	11 (10)	9.3 (8.8)
190	5.4 (6.4)	4.4 (5.4)
250	2.9 (3.2)	2.4 (2.7)
400	0.87 (1.0)	0.71 (0.83)
700	0.22 (0.32)	0.16 (0.25)

The product of the acceptance and the efficiency ( $A\epsilon$ ) is estimated by calculating  $A\epsilon_{MC}$  from the simulation, and multiplying it by the  $F$  to account for the difference in efficiency between simulation and data. The ADD, DM, and branon simulated samples are processed through the full GEANT4-based simulation of the CMS detector [26, 27], trigger emulation, and the same event reconstruction programs as used for data. For DM production, the simulated samples are produced using MADGRAPH 5v1.3.12 [39], and requiring  $E_T^\gamma > 130$  GeV and  $|\eta^\gamma| < 1.5$ . The estimated value of  $A\epsilon_{MC}$  for  $M_\chi$  in the range 1–1000 GeV varies over the range 41.6–44.4% for vector and 41.4–44.1% for axial-vector couplings, respectively. The  $E_T^\gamma$  spectra for ADD simulated events are generated using PYTHIA 8.153 [40], requiring  $E_T^\gamma > 130$  GeV. The  $A\epsilon_{MC}$  for the ADD model varies over the range 33.4–37.4% in the parameter space spanned by  $n = 3$ –6 and  $M_D = 1$ –3 TeV. The spectra for sim-




Figure 1: The photon  $E_T$  and  $\cancel{E}_T$  distributions for the candidate sample, compared with estimated contributions from SM backgrounds, and the predictions from the ADD model for  $M_D = 2$  TeV and  $n = 3$ . The horizontal bar on each data point indicates the width of the bin. The background uncertainty includes statistical and systematic components. The bottom panel shows the ratio of data and SM background predictions.

ulated branon events are generated using MADGRAPH 5v1.5.5 [39], requiring  $E_T^\gamma > 130$  GeV. The value of  $A\epsilon_{MC}$  for branon production varies over the range 41.3–48.9% in the parameter space spanned by the range of branon masses  $M_B = 100$ –3500 GeV and brane tensions  $f = 100$ –1000 GeV. The systematic uncertainty in  $A\epsilon_{MC}$  from the modeling of pileup, the energy calibration, and the resolution for photons, jets, and  $\cancel{E}_T$  is  $\pm 2.1\%$ . The systematic uncertainty from the scale fac-




Figure 2: Upper limits at 95% confidence level (CL) on the product of cross section and acceptance as a function of the  $E_T^\gamma$  threshold ( $>145$  GeV) for the photon and  $E_T$  final state.

tor is 6.4%, resulting in a total systematic uncertainty in  $A_{\text{MC}}$  of 6.7%. The systematic uncertainty in the measured integrated luminosity is  $\pm 2.6\%$  [41]. Theoretical uncertainties in the acceptance of the signal processes, based on the choice of PDF and scale, are found to be of order 1%, and thus have a negligible effect on the observed limits.

Upper limits on the signal cross section are calculated using the  $\text{CL}_s$  method [42, 43]. In the fit to the observed spectra, systematic uncertainties are represented by nuisance parameters with log-normal prior probability density functions. The changes in shape of the expected spectra that result from varying the photon energy scale and the theoretical differential cross section within their respective uncertainties are treated using a morphing technique [44]. The signal region studied in this analysis is defined with the requirement  $E_T^\gamma > 145$  GeV. The observed and expected upper limits on the product of cross section and acceptance ( $\sigma A$ ), plotted as a function of the  $E_T^\gamma$  threshold ( $> 145$  GeV), are shown in Fig. 2 and listed in Table 2. Results shown can be generally applied to any new physics that leads to the photon and  $E_T$  final state.

Tables 3 and 4 summarize the 90% CL upper limits on the production cross sections of the DM particles  $\chi\bar{\chi}$ , as a function of  $M_\chi$ . In general, the effective operator could be a mixture of vector and axial terms; for explicitness, the limiting cases of pure vector and pure axial vector operators have been chosen,





Figure 3: The 90% CL upper limits on the  $\chi$ -nucleon cross section as a function of the DM particle mass  $M_\chi$  for spin-independent couplings (top) and spin-dependent couplings (bottom). Results from the current search are shown as “CMS Monophoton, 8 TeV”. Shown are the limits from CMS using monojet [37] and monolepton [45] signatures (where  $\xi$  is the interference parameter addressing potentially different couplings to up- and down-type quarks and values of  $\xi = \pm 1$  maximize the effects of interference). Also shown are the limits from several published direct detection experiments [46–55]. The solid and hatched yellow contours show the 68% and 95% CL contours respectively for a possible signal from CDMS [56]. Limits similar to those from the current search are obtained by ATLAS [57].

corresponding to spin-independent and spin-dependent interactions, respectively. Following the procedures of Refs. [2] and [5], the upper limits on the DM production

Table 3: Dark Matter production cross sections as a function of the DM mass, assuming a vector interaction: theoretical DM production cross sections, where the generated photon transverse momentum is greater than 130 GeV and the contact interaction scale  $\Lambda$  is 10 TeV; observed (expected) 90% CL upper limits on the DM production cross section  $\sigma$ ; 90% CL lower limits on the contact interaction scale  $\Lambda$ ; and 90% CL upper limits on the  $\chi$ -nucleon cross section.

Mass [GeV]	$\sigma_{\text{theo}}$ [fb]	$\sigma$ [fb]	$\Lambda$ [GeV]	$\sigma_{\chi\text{-nucleon}}$ [cm <sup>2</sup> ]
1	$2.5 \times 10^{-4}$	7.8 (10.6)	750 (694)	$8.2 \times 10^{-40}$ ( $1.1 \times 10^{-39}$ )
10	$2.5 \times 10^{-4}$	8.0 (10.5)	745 (696)	$2.6 \times 10^{-39}$ ( $3.5 \times 10^{-39}$ )
100	$2.4 \times 10^{-4}$	8.0 (11.2)	742 (684)	$3.2 \times 10^{-39}$ ( $4.4 \times 10^{-39}$ )
200	$2.2 \times 10^{-4}$	7.6 (9.9)	729 (684)	$3.4 \times 10^{-39}$ ( $4.4 \times 10^{-39}$ )
300	$1.8 \times 10^{-4}$	6.9 (9.4)	714 (660)	$3.7 \times 10^{-39}$ ( $5.1 \times 10^{-39}$ )
500	$1.0 \times 10^{-4}$	5.2 (7.8)	666 (602)	$4.9 \times 10^{-39}$ ( $7.4 \times 10^{-39}$ )
1000	$1.5 \times 10^{-5}$	4.9 (7.2)	422 (382)	$3.1 \times 10^{-38}$ ( $4.6 \times 10^{-38}$ )

Table 4: Dark Matter production cross sections as a function of the DM mass, assuming an axial-vector interaction: theoretical DM production cross sections, where the generated photon transverse momentum is greater than 130 GeV and the contact interaction scale  $\Lambda$  is 10 TeV; observed (expected) 90% CL upper limits on the DM production cross section  $\sigma$ ; 90% CL lower limits on the contact interaction scale  $\Lambda$ ; and 90% CL upper limits on the  $\chi$ -nucleon cross section.

Mass [GeV]	$\sigma_{\text{theo}}$ [fb]	$\sigma$ [fb]	$\Lambda$ [GeV]	$\sigma_{\chi\text{-nucleon}}$ [cm <sup>2</sup> ]
1	$2.4 \times 10^{-4}$	7.9 (10.5)	746 (694)	$3.1 \times 10^{-41}$ ( $4.1 \times 10^{-41}$ )
10	$2.5 \times 10^{-4}$	7.9 (11.0)	748 (688)	$9.6 \times 10^{-41}$ ( $1.3 \times 10^{-40}$ )
100	$2.2 \times 10^{-4}$	8.2 (10.7)	718 (671)	$1.3 \times 10^{-40}$ ( $1.7 \times 10^{-40}$ )
200	$1.6 \times 10^{-4}$	6.7 (9.5)	702 (643)	$1.5 \times 10^{-40}$ ( $2.0 \times 10^{-40}$ )
300	$1.1 \times 10^{-4}$	5.8 (8.5)	663 (604)	$1.8 \times 10^{-40}$ ( $2.6 \times 10^{-40}$ )
500	$4.9 \times 10^{-5}$	5.5 (8.1)	544 (495)	$4.0 \times 10^{-40}$ ( $5.9 \times 10^{-40}$ )
1000	$4.2 \times 10^{-6}$	5.3 (7.7)	298 (272)	$4.5 \times 10^{-39}$ ( $6.5 \times 10^{-39}$ )




Figure 4: Observed limits on the SM-DM interaction mediator mass divided by coupling,  $M/\sqrt{g_\chi g_q}$ , as a function of the mediator mass  $M$ , assuming vector interactions, for DM particle masses of 50 GeV and 500 GeV. The width,  $\Gamma$ , of the mediator is varied between  $M/8\pi$  and  $M/3$ . The dotted lines show contours of constant coupling.

cross sections are converted into corresponding lower limits on the contact interaction scale  $\Lambda$ , which are then translated into upper limits on the  $\chi$ -nucleon scattering cross sections, calculated within the EFT framework. These results, as a function of  $M_\chi$ , are listed in Tables 3 and 4 and also displayed in Fig. 3. Superimposed are the results published by other experiments [46–56].

Table 5: Observed and expected 95% CL lower limits on ADD model parameters  $M_D$ , the effective Planck scale, as a function of  $n$ , the number of extra dimensions.

$n$	Obs. Limit [TeV]	Exp. Limit [TeV]
3	2.12	1.96
4	2.07	1.92
5	2.02	1.89
6	1.97	1.88

The validity of the EFT framework at the energy scale probed by the LHC has been recently explored in detail [2, 3, 5, 65–67]. These studies show that the condition  $M \gg Q$  may not always be satisfied because of the high momentum transfer scale at the LHC energies. Therefore, to interpret the data in a meaningful way where the EFT does not hold, following [3] we consider a simplified model predicting DM produc-




Figure 5: The 95% CL lower limits on the effective Planck scale,  $M_D$ , as a function of the number of extra dimensions in the ADD model, together with LO results from similar searches at the Tevatron [58, 59], LEP [60–63] and CMS [64].

tion via an  $s$ -channel vector mediator. For this simplified model, the simulated samples are produced using MADGRAPH 5v1.5.12 [39], and requiring  $E_T^\gamma > 130$  GeV and  $|\eta^\gamma| < 1.5$ . Limits on the SM-DM interaction mediator mass divided by coupling, for this model, are shown in Fig. 4. The mass of the mediator is varied for two fixed values of the mass of the DM particle: 50 GeV and 500 GeV, and the width of the mediator is varied from  $M/8\pi$  to  $M/3$  [3]. The contours for fixed values of  $\sqrt{g_\chi g_q}$  are also shown for comparison. For  $M_\chi = 500$  GeV the results for a mediator with a mass  $\gtrsim 5$  TeV are similar to those obtained from the EFT approach as listed in Table 3, while the limits are weaker for  $M \lesssim 100$  GeV. The limits are stronger than those of the EFT approach in the range of  $M$  from  $\sim 100$  GeV to  $\sim 4$  TeV, because of the resonance production enhancement in the cross section. In other words, the limits derived within the EFT framework are conservative in this region. For illustration purposes, similar distributions for  $M_\chi = 50$  GeV are also shown in Fig. 4.

Upper limits at 95% CL are also placed on the production cross section of the ADD and branon models, and translated into exclusions on the parameter space of the models. For the ADD model we follow the convention of Ref. [69] and only consider  $\hat{s} < M_D^2$  when calculating the cross sections. The limits on  $M_D$  for several values of  $n$ , the number of extra dimensions, are summarized in Table 5. These limits, along with existing





Figure 6: The 95% CL upper limits on the branon cross sections as a function of the branon mass  $M_B$  for  $N=1$ . Also shown are the theoretical cross sections in the branon model for the brane tension scale  $f = 100, 200, 300$ , and  $400$  GeV (top). Limits on  $f$  as a function of  $M_B$ , compared to results from similar searches at LEP [68] and the Tevatron [13] (bottom).

ADD limits from the Tevatron [58, 59] and LEP [60–63], are shown in Fig. 5 as a function of  $M_D$ . All these results are based on LO cross sections. Our results extend significantly the experimental limits on the ADD model in the single-photon channel [64, 70], and set limits of  $M_D > 2.12$ – $1.97$  TeV for  $n = 3$ – $6$ , at 95% CL. These results are comparable with the recent ATLAS limits [57].

Limits on  $f$  for branons are summarized in Table 6. For massless branons, the brane tension  $f$  is found to be greater than 410 GeV at 95% CL. These limits along

Table 6: Observed and expected 95% CL lower limits on the brane tension  $f$  as a function of the branon mass  $M_B$  for  $N=1$ .

	$M_B$ [GeV]									
	100	500	1000	1500	2000	2500	2800	3000	3200	3500
Obs. limit [GeV]	410	380	320	240	170	97	59	48	36	20
Exp. limit [GeV]	400	370	310	240	170	97	59	48	36	20

with the existing limits from LEP [68] and the Tevatron [13], are shown in Fig. 6. Branon masses  $M_B < 3.5$  TeV are excluded at 95% CL for low brane tension (20 GeV). These bounds are the most stringent published to date. These limits complement astrophysical constraints already set on the branon parameters [12].

## 6. Summary

Proton-proton collision events containing a photon and missing transverse momentum have been investigated to search for new phenomena. In the  $\sqrt{s} = 8$  TeV data set corresponding to  $19.6 \text{ fb}^{-1}$  of integrated luminosity, no deviations from the standard model predictions are observed. Bounds are placed on models predicting monophoton events; specifically, 95% confidence level upper limits for the cross section times acceptance for the selected final state are set and vary from  $14.0 \text{ fb}$  for  $E_T^\gamma > 145 \text{ GeV}$  to  $0.22 \text{ fb}$  for  $E_T^\gamma > 700 \text{ GeV}$ . Constraints are set on  $\chi$  production and translated into upper limits on vector and axial-vector contributions to the  $\chi$ -nucleon scattering cross section, assuming the validity of the EFT framework. For  $M_\chi = 10 \text{ GeV}$ , the  $\chi$ -nucleon cross section is constrained to be less than  $2.6 \times 10^{-39} \text{ cm}^2$  ( $9.6 \times 10^{-41} \text{ cm}^2$ ) for a spin-independent (spin-dependent) interaction at 90% confidence level. In addition the most stringent limits to date are obtained on the effective Planck scale in the ADD model with large spatial extra dimensions and on the brane tension scale in the branon model.

## Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction

and operation of the LHC and the CMS detector provided by the following funding agencies: BMWFW and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, and FAPESP (Brazil); MES (Bulgaria); CERN; CAS, MoST, and NSFC (China); COLCIENCIAS (Colombia); MSES and CSF (Croatia); RPF (Cyprus); MoER, ERC IUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece); OTKA and NIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); NRF and WCU (Republic of Korea); LAS (Lithuania); MOE and UM (Malaysia); CINVESTAV, CONACYT, SEP, and UASLP-FAI (Mexico); MBIE (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Dubna); MON, RosAtom, RAS and RFBR (Russia); MESTD (Serbia); SEIDI and CPAN (Spain); Swiss Funding Agencies (Switzerland); MST (Taipei); ThEPCenter, IPST, STAR and NSTDA (Thailand); TUBITAK and TAEK (Turkey); NASU and SFFR (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie programme and the European Research Council and EPLANET (European Union); the Leventis Foundation; the A. P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Council of Science and Industrial Research, India; the HOMING PLUS programme of Foundation for Polish Science, cofinanced from European Union, Regional Development Fund; the Compagnia di San Paolo (Torino); the Consorzio per la Fisica (Trieste); MIUR project 20108T4XTM (Italy); the Thalís and Aristeia programmes cofinanced by EU-ESF and the Greek NSRF; and the National Priorities Research Program by Qatar National Research Fund.

## References

- [1] R. Gaitskell, Direct detection of dark matter, *Annual Review of Nuclear and Particle Science* 54 (2004) 315. doi:10.1146/annurev.nucl.54.070103.181244.
- [2] Y. Bai, P. J. Fox, R. Harnik, The Tevatron at the frontier of dark matter direct detection, *JHEP* 12 (2010) 048. doi:10.1007/JHEP12(2010)048. arXiv:1005.3797v2.
- [3] P. J. Fox, R. Harnik, J. Kopp, Y. Tsai, Missing energy signatures of dark matter at the LHC, *Phys. Rev. D* 85 (2012) 056011. doi:10.1103/PhysRevD.85.056011. arXiv:1109.4398.
- [4] J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T. M. P. Tait, H.-B. Yu, Constraints on light Majorana dark matter from colliders, *Phys. Lett. B* 695 (2011) 185. doi:10.1016/j.physletb.2010.11.009. arXiv:1005.1286.
- [5] J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T. M. P. Tait, H.-B. Yu, Constraints on dark matter from colliders, *Phys. Rev. D* 82 (2010) 116010. doi:10.1103/PhysRevD.82.116010. arXiv:1008.1783.
- [6] N. Arkani-Hamed, S. Dimopoulos, G. R. Dvali, The hierarchy problem and new dimensions at a millimeter, *Phys. Lett. B* 429 (1998) 263. doi:10.1016/S0370-2693(98)00466-3. arXiv:hep-ph/9803315.
- [7] N. Arkani-Hamed, S. Dimopoulos, G. R. Dvali, Phenomenology, astrophysics and cosmology of theories with submillimeter dimensions and TeV scale quantum gravity, *Phys. Rev. D* 59 (1999) 086004. doi:10.1103/PhysRevD.59.086004. arXiv:hep-ph/9807344.
- [8] R. Sundrum, Effective field theory for a three-brane universe, *Phys. Rev. D* 59 (1999) 085009. doi:10.1103/PhysRevD.59.085009. arXiv:hep-ph/9805471.
- [9] A. Dobado, A. L. Maroto, The dynamics of the Goldstone bosons on the brane, *Nucl. Phys. B* 592 (2001) 203. doi:10.1016/S0550-3213(00)00574-5. arXiv:hep-ph/0007100.
- [10] J. A. R. Cembranos, A. Dobado, A. L. Maroto, Brane skyrmions and wrapped states, *Phys. Rev. D* 65 (2002) 026005. doi:10.1103/PhysRevD.65.026005. arXiv:hep-ph/0106322.
- [11] J. A. R. Cembranos, R. L. Delgado, A. Dobado, Brane worlds at the LHC: branons and KK gravitons, *Phys. Rev. D* 88 (2013) 075021. doi:10.1103/PhysRevD.88.075021. arXiv:1306.4900.
- [12] J. A. R. Cembranos, A. Dobado, A. L. Maroto, Cosmological and astrophysical limits on brane fluctuations, *Phys. Rev. D* 68 (2003) 103505. doi:10.1103/PhysRevD.68.103505. arXiv:hep-ph/0307062.
- [13] J. A. R. Cembranos, A. Dobado, A. L. Maroto, Branon search in hadronic colliders, *Phys. Rev. D* 70 (2004) 096001. doi:10.1103/PhysRevD.70.096001. arXiv:hep-ph/0405286.
- [14] CMS Collaboration, Energy calibration and resolution of the CMS electromagnetic calorimeter in pp collisions at  $\sqrt{s} = 7$  TeV, *JINST* 8 (2013) P09009. doi:10.1088/1748-0221/8/09/P09009. arXiv:1306.2016.
- [15] CMS Collaboration, The CMS experiment at the CERN LHC, *JINST* 3 (2008) S08004. doi:10.1088/1748-0221/3/08/S08004.
- [16] CMS Collaboration, Performance of photon reconstruction and identification with the CMS detector in proton-proton collisions at  $\sqrt{s} = 8$  TeV, 2015. arXiv:1502.02702, submitted to JINST.
- [17] CMS Collaboration, Isolated Photon Reconstruction and Identification at  $\sqrt{s} = 7$  TeV, CMS Physics Analysis Summary CMS-PAS-EGM-10-006, 2011. URL: <http://cdsweb.cern.ch/record/1324545>.
- [18] CMS Collaboration, Particle-Flow Event Reconstruction in CMS and Performance for Jets, Taus, and MET, CMS Physics Analysis Summary CMS-PAS-PFT-09-001, 2009. URL: <http://cdsweb.cern.ch/record/1194487>.
- [19] CMS Collaboration, Commissioning of the Particle-flow Event Reconstruction with the first LHC collisions recorded in the CMS detector, CMS Physics Analysis Summary CMS-PAS-PFT-10-001, 2010. URL: <http://cdsweb.cern.ch/record/1247373>.
- [20] M. Cacciari, G. P. Salam, G. Soyez, The catchment area of jets, *JHEP* 04 (2008) 005. doi:10.1088/1126-6708/2008/04/005. arXiv:0802.1188.
- [21] CMS Collaboration, Mitigation of anomalous APD signals in the CMS electromagnetic calorimeter, in: XVth International Conference on Calorimetry in High Energy Physics (CALOR2012), Santa Fe, USA, 2012. doi:10.1088/1742-6596/404/1/012043, [J. Phys. Conf. Ser. 404 (2012) 012043].
- [22] CMS Collaboration, Electron Reconstruction and Identification at  $\sqrt{s} = 7$  TeV, CMS Physics Analysis Summary CMS-PAS-EGM-10-004, 2010. URL: <http://cdsweb.cern.ch/record/1299116>.
- [23] M. Cacciari, G. P. Salam, G. Soyez, The anti- $k_r$  jet clustering algorithm, *JHEP* 04 (2008) 063. doi:10.1088/1126-6708/2008/04/063. arXiv:0802.1189.
- [24] CMS Collaboration, Pileup Jet Identification, CMS Physics Analysis Summary CMS-PAS-JME-13-005, 2013. URL: <http://cdsweb.cern.ch/record/1581583>.
- [25] CMS Collaboration, MET performance in 8 TeV data, CMS Physics Analysis Summary CMS-PAS-JME-12-002, 2013. URL: <http://cdsweb.cern.ch/record/1543527>.
- [26] S. Agostinelli, et al. (GEANT4), Geant4—a simulation toolkit, *Nucl. Instrum. Meth. A* 506 (2003) 250. doi:10.1016/S0168-9002(03)01368-8.
- [27] J. Allison, et al., Geant4 developments and applications, *IEEE Trans. Nucl. Sci.* 53 (2006) 270. doi:10.1109/TNS.2006.869826.
- [28] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torielli, M. Zaro, The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, *JHEP* 07 (2014) 079. doi:10.1007/JHEP07(2014)079. arXiv:1405.0301.
- [29] J. Campbell, R. Ellis, C. Williams, MCFM v6.1: A Monte Carlo for FeMtobarn processes at Hadron Colliders, 2011. URL: <http://mcfm.fnal.gov/mcfm.pdf>.
- [30] S. Alekhin, et al., The PDF4LHC Working Group Interim Report, 2011. arXiv:1101.0536.
- [31] M. Botje, J. Butterworth, A. Cooper-Sarkar, A. de Roeck, J. Feltesse, S. Forte, A. Glazov, J. Huston, R. McNulty, T. Sjöstrand, R. Thorne, The PDF4LHC Working Group Interim Recommendations, 2011. arXiv:1101.0538.
- [32] R. D. Ball, V. Bertone, F. Cerutti, L. Del Debbio, S. Forte, A. Guffanti, J. I. Latorre, J. Rojo, M. Ubiali (NNPDF), Impact of heavy quark masses on parton distributions and LHC phenomenology, *Nucl. Phys. B* 849 (2011) 296. doi:10.1016/j.nuclphysb.2011.03.021. arXiv:1101.1300.
- [33] T. Sjöstrand, S. Mrenna, P. Z. Skands, Pythia 6.4 physics and manual, *JHEP* 05 (2006) 26. doi:10.1088/1126-6708/2006/05/026. arXiv:hep-ph/0603175.
- [34] J. Pumplin, D. R. Stump, J. Huston, H.-L. Lai, P. Nadolsky, W.-K. Tung, New generation of parton distributions with uncertainties from global QCD analysis, *JHEP* 07 (2002) 012. doi:10.1088/1126-6708/2002/07/012. arXiv:hep-ph/0201195.
- [35] CMS Collaboration, Determination of jet energy calibration and transverse momentum resolution in CMS, *JINST* 6 (2011) P11002. doi:10.1088/1748-0221/6/11/P11002.

- arXiv:1107.4277.
- [36] CMS Collaboration, Missing transverse energy performance of the CMS detector, *J. Instrum.* 6 (2011) P09001. doi:10.1088/1748-0221/6/09/P09001.
- [37] CMS Collaboration, Search for dark matter, extra dimensions, and unparticles in monojet events in proton-proton collisions at  $\sqrt{s} = 8$  TeV, 2014. arXiv:1408.3583, submitted to *Eur. Phys. J. C*.
- [38] CMS Collaboration, Measurement of the inclusive W and Z production cross sections in pp collisions at  $\sqrt{s} = 7$  TeV with the cms experiment, *J. High Energy Phys.* 10 (2011) 132. doi:10.1007/JHEP10(2011)132.
- [39] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer, T. Stelzer, *MadGraph 5: going beyond*, *JHEP* 06 (2011) 128. doi:10.1007/JHEP06(2011)128. arXiv:1106.0522.
- [40] T. Sjöstrand, S. Mrenna, P. Skands, A brief introduction to PYTHIA 8.1, *Comput. Phys. Commun.* 178 (2008) 852. doi:10.1016/j.cpc.2008.01.036. arXiv:0710.3820.
- [41] CMS Collaboration, CMS Luminosity Based on Pixel Cluster Counting - Summer 2013 Update, CMS Physics Analysis Summary CMS-PAS-LUM-13-001, 2013. URL: <http://cdsweb.cern.ch/record/1598864>.
- [42] A. L. Read, Presentation of search results: the  $CL_s$  technique, *J. Phys. G* 28 (2002) 2693. doi:10.1088/0954-3889/28/10/313.
- [43] T. Junk, Confidence level computation for combining searches with small statistics, *Nucl. Instrum. Meth. A* 434 (1999) 435. doi:10.1016/S0168-9002(99)00498-2. arXiv:hep-ex/9902006.
- [44] J. S. Conway, Nuisance parameters in likelihoods for multi-source spectra, in: H. B. Propser, L. Lyons (Eds.), *Proceedings of PHYSTAT 2011 Workshop on Statistical Issues Related to Discovery Claims in Search Experiments and Unfolding*, CERN, Geneva, Switzerland, 2011, p. 115. doi:10.5170/CERN-2011-006.
- [45] CMS Collaboration, Search for physics beyond the standard model in final states with a lepton and missing transverse energy in proton-proton collisions at  $\sqrt{s} = 8$  TeV, 2014. arXiv:1408.2745, submitted to *Phys. Rev. D*.
- [46] E. Aprile, et al. (XENON100), Dark Matter Results from 225 Live Days of XENON100 Data, *Phys. Rev. Lett.* 109 (2012) 181301. doi:10.1103/PhysRevLett.109.181301. arXiv:1207.5988.
- [47] Z. Ahmed, et al. (CDMS), Results from a low-energy analysis of the CDMS II germanium data, *Phys. Rev. Lett.* 106 (2011) 131302. doi:10.1103/PhysRevLett.106.131302. arXiv:1011.2482v3.
- [48] Z. Ahmed, et al. (CDMS II), Dark matter search results from the CDMS II experiment, *Science* 327 (2010) 1619. doi:10.1126/science.1186112.
- [49] C. E. Aalseth, et al. (CoGeNT), Results from a search for light-mass dark matter with a  $p$ -type point contact germanium detector, *Phys. Rev. Lett.* 106 (2011) 131301. doi:10.1103/PhysRevLett.106.131301. arXiv:1002.4703.
- [50] M. Felizardo, et al. (SIMPLE), Final Analysis and Results of the Phase II SIMPLE Dark Matter Search, *Phys. Rev. Lett.* 108 (2012) 201302. doi:10.1103/PhysRevLett.108.201302. arXiv:1106.3014.
- [51] E. Behnke, et al. (COUPP), First Dark Matter Search Results from a 4-kg  $\text{CF}_3\text{I}$  Bubble Chamber Operated in a Deep Underground Site, *Phys. Rev. D* 86 (2012) 052001. doi:10.1103/PhysRevD.86.052001. arXiv:1204.3094, [Erratum: 10.1103/PhysRevD.90.079902].
- [52] M. G. Aartsen, et al. (IceCube), Search for dark matter annihilations in the Sun with the 79-string IceCube detector, *Phys. Rev. Lett.* 110 (2013) 131302. doi:10.1103/PhysRevLett.110.131302. arXiv:1212.4097.
- [53] T. Tanaka, et al. (Super-Kamiokande), An indirect search for weakly interacting massive particles in the sun using 3109.6 days of upward-going muons in Super-Kamiokande, *Astrophys. J.* 742 (2011) 78. doi:10.1088/0004-637X/742/2/78. arXiv:1108.3384.
- [54] D. S. Akerib, et al. (LUX), First results from the LUX dark matter experiment at the Sanford Underground Research Facility, *Phys. Rev. Lett.* 112 (2014) 091303. doi:10.1103/PhysRevLett.112.091303. arXiv:1310.8214.
- [55] R. Agnese, et al. (SuperCDMS Soudan), Search for Low-Mass Weakly Interacting Massive Particles Using Voltage-Assisted Calorimetric Ionization Detection in the SuperCDMS Experiment, *Phys. Rev. Lett.* 112 (2014) 041302. doi:10.1103/PhysRevLett.112.041302. arXiv:1309.3259.
- [56] R. Agnese, et al. (CDMS), Silicon Detector Dark Matter Results from the Final Exposure of CDMS II, *Phys. Rev. Lett.* 111 (2013) 251301. doi:10.1103/PhysRevLett.111.251301. arXiv:1304.4279.
- [57] ATLAS Collaboration, Search for new phenomena in events with a photon and missing transverse momentum in  $pp$  collisions at  $\sqrt{s} = 8$  TeV with the ATLAS detector, *Phys. Rev. D* 91 (2015) 012008. doi:10.1103/PhysRevD.91.012008. arXiv:1411.1559.
- [58] T. Aaltonen, et al. (CDF), Search for large extra dimensions in final states containing one photon or jet and large missing transverse energy produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV, *Phys. Rev. Lett.* 101 (2008) 181602. doi:10.1103/PhysRevLett.101.181602. arXiv:0807.3132.
- [59] V. M. Abazov, et al. (D0), Search for Large Extra Dimensions via Single Photon plus Missing Energy Final States at  $\sqrt{s} = 1.96$  TeV, *Phys. Rev. Lett.* 101 (2008) 011601. doi:10.1103/PhysRevLett.101.011601. arXiv:0803.2137.
- [60] J. Abdallah, et al. (DELPHI), Photon events with missing energy in  $e^+e^-$  collisions at  $\sqrt{s} = 130$  GeV to 209 GeV, *Eur. Phys. J. C* 38 (2005) 395. doi:10.1140/epjc/s2004-02051-8. arXiv:hep-ex/0406019.
- [61] P. Achard, et al. (L3), Single- and multi-photon events with missing energy in  $e^+e^-$  collisions at LEP, *Phys. Lett. B* 587 (2004) 16. doi:10.1016/j.physletb.2004.01.010. arXiv:hep-ex/0402002.
- [62] G. Abbiendi, et al. (OPAL), Photonic events with missing energy in  $e^+e^-$  collisions at  $\sqrt{s} = 189$  GeV, *Eur. Phys. J. C* 18 (2000) 253. doi:10.1007/s100520000522. arXiv:hep-ex/0005002.
- [63] A. Heister, et al. (ALEPH), Single photon and multiphoton production in  $e^+e^-$  collisions at  $\sqrt{s}$  up to 209 GeV, *Eur. Phys. J. C* 28 (2003) 1. doi:10.1140/epjc/s2002-01129-7.
- [64] CMS Collaboration, Search for Dark Matter and Large Extra Dimensions in  $pp$  Collisions Yielding a Photon and Missing Transverse Energy, *Phys. Rev. Lett.* 108 (2012) 261803. doi:10.1103/PhysRevLett.108.261803. arXiv:1204.0821.
- [65] H. An, X. Ji, L.-T. Wang, Light dark matter and  $Z'$  dark force at colliders, *JHEP* 07 (2012) 182. doi:10.1007/JHEP07(2012)182. arXiv:1202.2894.
- [66] A. Friedland, M. L. Graesser, I. M. Shoemaker, L. Vecchi, Probing nonstandard standard model backgrounds with LHC monojets, *Phys. Lett. B* 714 (2012) 267. doi:10.1016/j.physletb.2012.06.078. arXiv:1111.5331.
- [67] O. Buchmüller, M. J. Dolan, C. McCabe, Beyond effective field theory for dark matter searches at the LHC, *JHEP* 01 (2014) 025. doi:10.1007/JHEP01(2014)025. arXiv:1308.6799.
- [68] P. Achard, et al. (L3), Search for bransons at LEP, *Phys. Lett. B* 597 (2004) 145. doi:10.1016/j.physletb.2004.07.014.

- arXiv:hep-ex/0407017.
- [69] G. F. Giudice, R. Rattazzi, J. D. Wells, Quantum gravity and extra dimensions at high-energy colliders, *Nucl. Phys. B* 544 (1999) 3. doi:10.1016/S0550-3213(99)00044-9. arXiv:hep-ph/9811291.
- [70] ATLAS Collaboration, Search for dark matter candidates and large extra dimensions in events with a photon and missing transverse momentum in  $pp$  collision data at  $\sqrt{s} = 7$  TeV with the ATLAS detector, *Phys. Rev. Lett.* 110 (2013) 011802. doi:10.1103/PhysRevLett.110.011802. arXiv:1209.4625.

**Yerevan Physics Institute, Yerevan, Armenia**

V. Khachatryan, A.M. Sirunyan, A. Tumasyan

**Institut für Hochenergiephysik der OeAW, Wien, Austria**

W. Adam, T. Bergauer, M. Dragicevic, J. Erö, M. Friedl, R. Frühwirth<sup>1</sup>, V.M. Ghete, C. Hartl, N. Hörmann, J. Hrubec, M. Jeitl<sup>1</sup>, W. Kiesenhofer, V. Knünz, M. Krammer<sup>1</sup>, I. Krätschmer, D. Liko, I. Mikulec, D. Rabady<sup>2</sup>, B. Rabaran, H. Rohringer, R. Schöfbeck, J. Strauss, W. Treberer-Treberspurg, W. Waltenberger, C.-E. Wulz<sup>1</sup>

**National Centre for Particle and High Energy Physics, Minsk, Belarus**

V. Mossolov, N. Shumeiko, J. Suarez Gonzalez

**Universiteit Antwerpen, Antwerpen, Belgium**

S. Alderweireldt, M. Bansal, S. Bansal, T. Cornelis, E.A. De Wolf, X. Janssen, A. Knutsson, J. Lauwers, S. Luyckx, S. Ochesanu, R. Rougny, M. Van De Klundert, H. Van Haevermaet, P. Van Mechelen, N. Van Remortel, A. Van Spilbeek

**Vrije Universiteit Brussel, Brussel, Belgium**

F. Blekman, S. Blyweert, J. D'Hondt, N. Daci, N. Herculoux, J. Keaveney, S. Lowette, M. Maes, A. Olbrechts, Q. Python, D. Strom, S. Tavernier, W. Van Doninck, P. Van Mulders, G.P. Van Onsem, I. Villella

**Université Libre de Bruxelles, Bruxelles, Belgium**

C. Caillol, B. Clerbaux, G. De Lentdecker, D. Dobur, L. Favart, A.P.R. Gay, A. Grebenyuk, A. Léonard, A. Mohammadi, L. Perniè<sup>2</sup>, T. Reis, T. Seva, L. Thomas, C. Vander Velde, P. Vanlaer, J. Wang, F. Zenoni

**Ghent University, Ghent, Belgium**

V. Adler, K. Beernaert, L. Benucci, A. Cimmino, S. Costantini, S. Crucy, S. Dildick, A. Fagot, G. Garcia, J. Mccartin, A.A. Ocampo Rios, D. Ryckbosch, S. Salva Diblen, M. Sigamani, N. Strobbe, F. Thyssen, M. Tytgat, E. Yazgan, N. Zaganidis

**Université Catholique de Louvain, Louvain-la-Neuve, Belgium**

S. Basesmez, C. Beluffi<sup>3</sup>, G. Bruno, R. Castello, A. Caudron, L. Ceard, G.G. Da Silva, C. Delaere, T. du Pree, D. Favart, L. Forthomme, A. Giammancoti<sup>4</sup>, J. Hollar, A. Jafari, P. Jez, M. Komm, V. Lemaitre, C. Nuttens, D. Pagano, L. Perrini, A. Pin, K. Piotrkowski, A. Popov<sup>5</sup>, L. Quertenmont, M. Selvaggi, M. Vidal Marono, J.M. Vizan Garcia

**Université de Mons, Mons, Belgium**

N. Beliy, T. Caebergs, E. Daubie, G.H. Hammad

**Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil**

W.L. Aldá Júnior, G.A. Alves, L. Brito, M. Correa Martins Junior, T. Dos Reis Martins, C. Mora Herrera, M.E. Pol

**Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil**

W. Carvalho, J. Chinellato<sup>6</sup>, A. Custódio, E.M. Da Costa, D. De Jesus Damiao, C. De Oliveira Martins, S. Fonseca De Souza, H. Malbouisson, D. Matos Figueiredo, L. Mundim, H. Nogima, W.L. Prado Da Silva, J. Santaolalla, A. Santoro, A. Sznajder, E.J. Tonelli Manganot<sup>6</sup>, A. Vilela Pereira

**Universidade Estadual Paulista<sup>a</sup>, Universidade Federal do ABC<sup>b</sup>, São Paulo, Brazil**

C.A. Bernardes<sup>b</sup>, S. Dogra<sup>a</sup>, T.R. Fernandez Perez Tomei<sup>a</sup>, E.M. Gregores<sup>b</sup>, P.G. Mercadante<sup>b</sup>, S.F. Novaes<sup>a</sup>, Sandra S. Padula<sup>a</sup>

**Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria**

A. Aleksandrov, V. Genchev<sup>2</sup>, P. Iaydjiev, A. Marinov, S. Piperov, M. Rodozov, G. Sultanov, M. Vutova

**University of Sofia, Sofia, Bulgaria**

A. Dimitrov, I. Glushkov, R. Hadjiiska, L. Litov, B. Pavlov, P. Petkov

**Institute of High Energy Physics, Beijing, China**

J.G. Bian, G.M. Chen, H.S. Chen, M. Chen, T. Cheng, R. Du, C.H. Jiang, R. Plestina<sup>7</sup>, F. Romeo, J. Tao, Z. Wang

**State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China**

C. Asawatangtrakuldee, Y. Ban, Q. Li, S. Liu, Y. Mao, S.J. Qian, D. Wang, W. Zou

**Universidad de Los Andes, Bogota, Colombia**

C. Avila, L.F. Chaparro Sierra, C. Florez, J.P. Gomez, B. Gomez Moreno, J.C. Sanabria

**University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia**

N. Godinovic, D. Lelas, D. Polic, I. Puljak

**University of Split, Faculty of Science, Split, Croatia**

Z. Antunovic, M. Kovac

**Institute Rudjer Boskovic, Zagreb, Croatia**

V. Brigljevic, K. Kadija, J. Luetic, D. Mekterovic, L. Sudic

**University of Cyprus, Nicosia, Cyprus**

A. Attikis, G. Mavromanolakis, J. Mousa, C. Nicolaou, F. Ptochos, P.A. Razis

**Charles University, Prague, Czech Republic**

M. Bodlak, M. Finger, M. Finger Jr.<sup>8</sup>

**Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**

Y. Assran<sup>9</sup>, S. Elgammal<sup>10</sup>, M.A. Mahmoud<sup>11</sup>, A. Raditenskiy<sup>12</sup>

**National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**

M. Kadastik, M. Murumaa, M. Raidal, A. Tiko

**Department of Physics, University of Helsinki, Helsinki, Finland**

P. Eerola, G. Fedi, M. Voutilainen

**Helsinki Institute of Physics, Helsinki, Finland**

J. Härkönen, V. Karimäki, R. Kinnunen, M.J. Kortelainen, T. Lampén, K. Lassila-Perini, S. Lehti, T. Lindén, P. Luukka, T. Mäenpää, T. Peltola, E. Tuominen, J. Tuominiemi, E. Tuovinen, L. Wendland

**Lappeenranta University of Technology, Lappeenranta, Finland**

J. Talvitie, T. Tuuva

**DSM/IRFU, CEA/Saclay, Gif-sur-Yvette, France**

M. Besancon, F. Couderc, M. Dejardin, D. Denegri, B. Fabbro, J.L. Faure, C. Favaro, F. Ferri, S. Ganjour, A. Givernaud, P. Gras, G. Hamel de Monchenault, P. Jarry, E. Locci, J. Malcles, J. Neveu, J. Rander, A. Rosowsky, M. Titov

**Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France**

S. Baffioni, F. Beaudette, P. Busson, C. Charlot, T. Dahms, M. Dalchenko, L. Dobrzynski, N. Filipovic, A. Florent, R. Granier de Cassagnac, L. Mastrolorenzo, P. Miné, C. Mironov, I.N. Naranjo, M. Nguyen, C. Ochando, P. Paganini, S. Regnard, R. Salerno, J.B. Sauvan, Y. Sirois, C. Veelken, Y. Yilmaz, A. Zabi

**Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg, Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France**

J.-L. Agram<sup>14</sup>, J. Andrea, A. Aubin, D. Bloch, J.-M. Brom, E.C. Chabert, C. Collard, E. Contet<sup>14</sup>, J.-C. Fontainet<sup>14</sup>, D. Gelé, U. Goerlach, C. Goetzmann, A.-C. Le Bihan, P. Van Hove

**Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France**

S. Gadrat

**Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3, Institut de Physique**

**Nucléaire de Lyon, Villeurbanne, France**

S. Beauceron, N. Beaupere, G. Boudoult<sup>2</sup>, E. Bouvier, S. Brochet, C.A. Carrillo Montoya, J. Chasserat, R. Chierici, D. Contard<sup>2</sup>, P. Depasse, H. El Mamouni, J. Fan, J. Fay, S. Gascon, M. Gouzevitch, B. Ille, T. Kurca, M. Lethuillier, L. Mirabito, S. Perries, J.D. Ruiz Alvarez, D. Sabes, L. Sgandurra, V. Sordini, M. Vander Donckt, P. Verdier, S. Viret, H. Xiao

**E. Andronikashvili Institute of Physics, Academy of Science, Tbilisi, Georgia**

L. Rurua

**RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**

C. Autermann, S. Beranek, M. Bontenackels, M. Edelhoff, L. Feld, A. Heister, O. Hindrichs, K. Klein, A. Ostapchuk, F. Raupach, J. Sammet, S. Schael, H. Weber, B. Wittmer, V. Zhukov<sup>5</sup>

**RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany**

M. Ata, M. Brodski, E. Dietz-Laursonn, D. Duchardt, M. Erdmann, R. Fischer, A. Güth, T. Hebbeker, C. Heidemann, K. Hoepfner, D. Klingebiel, S. Knutzen, P. Kreuzer, M. Merschmeyer, A. Meyer, P. Millet, M. Olschewski, K. Padeken, P. Papacz, H. Reithler, S.A. Schmitz, L. Sonnenschein, D. Teyssier, S. Thüer, M. Weber

**RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany**

V. Cherepanov, Y. Erdogan, G. Flügge, H. Geenen, M. Geisler, W. Haj Ahmad, F. Hoehle, B. Kargoll, T. Kress, Y. Kuessel, A. Künsken, J. Lingemann<sup>2</sup>, A. Nowack, I.M. Nugent, L. Perchalla, O. Pooth, A. Stahl

**Deutsches Elektronen-Synchrotron, Hamburg, Germany**

I. Asin, N. Bartosik, J. Behr, W. Behrenhoff, U. Behrens, A.J. Bell, M. Bergholtz<sup>15</sup>, A. Bethani, K. Borras, A. Burgmeier, A. Cakir, L. Calligaris, A. Campbell, S. Choudhury, F. Costanza, C. Diez Pardos, G. Dolinska, S. Dooling, T. Dorland, G. Eckerlin, D. Eckstein, T. Eichhorn, G. Flucke, J. Garay Garcia, A. Geiser, P. Gunnellini, J. Hauk, M. Hempelt<sup>15</sup>, D. Horton, H. Jung, A. Kalogeropoulos, M. Kasemann, P. Katsas, J. Kieseler, C. Kleinwort, I. Korol, D. Krücker, W. Lange, J. Leonard, K. Lipka, A. Lobanov, W. Lohmann<sup>15</sup>, B. Lutz, R. Mankel, I. Marfint<sup>15</sup>, I.-A. Melzer-Pellmann, A.B. Meyer, G. Mittag, J. Mnich, A. Mussgiller, S. Naumann-Emme, A. Nayak, O. Novgorodova, E. Ntomari, H. Perrey, D. Pitzl, R. Placakyte,

A. Raspereza, P.M. Ribeiro Cipriano, B. Roland, E. Ron, M.Ö. Sahin, J. Salfeld-Nebgen, P. Saxena, R. Schmidt<sup>15</sup>, T. Schoerner-Sadenius, M. Schröder, C. Seitz, S. Spannagel, A.D.R. Vargas Trevino, R. Walsh, C. Wissing

**University of Hamburg, Hamburg, Germany**

M. Aldaya Martin, V. Blobel, M. Centis Vignali, A.R. Draeger, J. Erfle, E. Garutti, K. Goebel, M. Görner, J. Haller, M. Hoffmann, R.S. Höing, H. Kirschenmann, R. Klanner, R. Kogler, J. Lange, T. Lapsien, T. Lenz, I. Marchesini, J. Ott, T. Peiffer, A. Perieanu, N. Pietsch, J. Poehlsen, T. Poehlsen, D. Rathjens, C. Sander, H. Schettler, P. Schleper, E. Schlieckau, A. Schmidt, M. Seidel, V. Sola, H. Stadie, G. Steinbrück, D. Troendle, E. Usai, L. Vanelderen, A. Vanhoef

**Institut für Experimentelle Kernphysik, Karlsruhe, Germany**

C. Barth, C. Baus, J. Berger, C. Böser, E. Butz, T. Chwalek, W. De Boer, A. Descroix, A. Dierlamm, M. Feindt, F. Frensch, M. Giffels, A. Gilbert, F. Hartmann<sup>2</sup>, T. Haut<sup>2</sup>, U. Husemann, I. Katkov<sup>5</sup>, A. Kornmayert<sup>2</sup>, E. Kuznetsova, P. Lobbelle Pardo, M.U. Mozer, Th. Müller, A. Nürnberg, G. Quast, K. Rabbertz, S. Röcker, H.J. Simonis, F.M. Stober, R. Ulrich, J. Wagner-Kuhr, S. Wayand, T. Weiler, R. Wolf

**Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece**

G. Anagnostou, G. Daskalakis, T. Gerasimidis, V.A. Giakoumopoulou, A. Kyriakis, D. Loukas, A. Markou, C. Markou, A. Psallidas, I. Topsis-Giotis

**University of Athens, Athens, Greece**

A. Agapitos, S. Kesisoglou, A. Panagiotou, N. Saoulidou, E. Stiliaris

**University of Ioánnina, Ioánnina, Greece**

X. Aslanoglou, I. Evangelou, G. Flouris, C. Foudas, P. Kokkas, N. Manthos, I. Papadopoulos, E. Paradas, J. Strolagas

**Wigner Research Centre for Physics, Budapest, Hungary**

G. Bencze, C. Hajdu, P. Hidas, D. Horvath<sup>16</sup>, F. Sikler, V. Veszpremi, G. Vesztergombit<sup>17</sup>, A.J. Zsigmond

**Institute of Nuclear Research ATOMKI, Debrecen, Hungary**

N. Beni, S. Czellar, J. Karancsit<sup>18</sup>, J. Molnar, J. Palinkas, Z. Szillasi

**University of Debrecen, Debrecen, Hungary**

A. Makovec, P. Raics, Z.L. Trocsanyi, B. Ujvari

**National Institute of Science Education and Research, Bhubaneswar, India**

S.K. Swain

**Panjab University, Chandigarh, India**

S.B. Beri, V. Bhatnagar, R. Gupta, U.Bhawandeep, A.K. Kalsi, M. Kaur, R. Kumar, M. Mittal, N. Nishu, J.B. Singh

**University of Delhi, Delhi, India**

Ashok Kumar, Arun Kumar, S. Ahuja, A. Bhardwaj, B.C. Choudhary, A. Kumar, S. Malhotra, M. Naimuddin, K. Ranjan, V. Sharma

**Saha Institute of Nuclear Physics, Kolkata, India**

S. Banerjee, S. Bhattacharya, K. Chatterjee, S. Dutta, B. Gomber, Sa. Jain, Sh. Jain, R. Khurana, A. Modak, S. Mukherjee, D. Roy, S. Sarkar, M. Sharan

**Bhabha Atomic Research Centre, Mumbai, India**

A. Abdulsalam, D. Dutta, S. Kailas, V. Kumar, A.K. Mohanty<sup>textsuperscript2</sup>, L.M. Pant, P. Shukla, A. Topkar

**Tata Institute of Fundamental Research, Mumbai, India**

T. Aziz, S. Banerjee, S. Bhowmik<sup>textsuperscript19</sup>, R.M. Chatterjee, R.K. Dewanjee, S. Dugad, S. Ganguly, S. Ghosh, M. Guchait, A. Gurtu<sup>textsuperscript20</sup>, G. Kole, S. Kumar, M. Maity<sup>textsuperscript19</sup>, G. Majumder, K. Mazumdar, G.B. Mohanty, B. Parida, K. Sudhakar, N. Wickramaget<sup>textsuperscript21</sup>

**Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**

H. Bakhshiansohi, H. Behnamian, S.M. Etesamit<sup>textsuperscript22</sup>, A. Fahim<sup>textsuperscript23</sup>, R. Goldouzian, M. Khakzad, M. Mohammadi Najafabadi, M. Naseri, S. Paktinat Mehdiabadi, F. Rezaei Hosseinabadi, B. Safarzadeht<sup>textsuperscript24</sup>, M. Zeinali

**University College Dublin, Dublin, Ireland**

M. Felcini, M. Grunewald

**INFN Sezione di Bari <sup>a</sup>, Università di Bari <sup>b</sup>, Politecnico di Bari <sup>c</sup>, Bari, Italy**

M. Abbrescia<sup>a,b</sup>, C. Calabria<sup>a,b</sup>, S.S. Chhibra<sup>a,b</sup>, A. Colaleo<sup>a</sup>, D. Creanza<sup>a,c</sup>, N. De Filippis<sup>a,c</sup>, M. De Palma<sup>a,b</sup>, L. Fiore<sup>a</sup>, G. Iaselli<sup>a,c</sup>, G. Maggi<sup>a,c</sup>, M. Maggi<sup>a</sup>, S. My<sup>a,c</sup>, S. Nuzzo<sup>a,b</sup>, A. Pompili<sup>a,b</sup>, G. Pugliese<sup>a,c</sup>, R. Radogna<sup>a,b, textsuperscript2</sup>, G. Selvaggi<sup>a,b</sup>, A. Sharma, L. Silvestris<sup>a, textsuperscript2</sup>, R. Venditti<sup>a,b</sup>

**INFN Sezione di Bologna <sup>a</sup>, Università di Bologna <sup>b</sup>, Bologna, Italy**

G. Abbiendi<sup>a</sup>, A.C. Benvenuti<sup>a</sup>, D. Bonacorsi<sup>a,b</sup>, S. Braibant-Giacomelli<sup>a,b</sup>, L. Brigliadori<sup>a,b</sup>, R. Campanini<sup>a,b</sup>, P. Capiluppi<sup>a,b</sup>, A. Castro<sup>a,b</sup>, F.R. Cavallo<sup>a</sup>, G. Codispoti<sup>a,b</sup>, M. Cuffiani<sup>a,b</sup>,

G.M. Dallavalle<sup>a</sup>, F. Fabbri<sup>a</sup>, A. Fanfani<sup>a,b</sup>, D. Fasanella<sup>a,b</sup>, P. Giacomelli<sup>a</sup>, C. Grandi<sup>a</sup>, L. Guiducci<sup>a,b</sup>, S. Marcellini<sup>a</sup>, G. Masetti<sup>a</sup>, A. Montanari<sup>a</sup>, F.L. Navarria<sup>a,b</sup>, A. Perrotta<sup>a</sup>, F. Primavera<sup>a,b</sup>, A.M. Rossi<sup>a,b</sup>, T. Rovelli<sup>a,b</sup>, G.P. Siroli<sup>a,b</sup>, N. Tosi<sup>a,b</sup>, R. Travaglini<sup>a,b</sup>

**INFN Sezione di Catania <sup>a</sup>, Università di Catania <sup>b</sup>, CSFNSM <sup>c</sup>, Catania, Italy**

S. Albergo<sup>a,b</sup>, G. Cappello<sup>a</sup>, M. Chiorboli<sup>a,b</sup>, S. Costa<sup>a,b</sup>, F. Giordano<sup>a,c, textsuperscript2</sup>, R. Potenza<sup>a,b</sup>, A. Tricomi<sup>a,b</sup>, C. Tuve<sup>a,b</sup>

**INFN Sezione di Firenze <sup>a</sup>, Università di Firenze <sup>b</sup>, Firenze, Italy**

G. Barbagli<sup>a</sup>, V. Ciulli<sup>a,b</sup>, C. Civinini<sup>a</sup>, R. D'Alessandro<sup>a,b</sup>, E. Focardi<sup>a,b</sup>, E. Gallo<sup>a</sup>, S. Gozzi<sup>a,b</sup>, V. Gori<sup>a,b, textsuperscript2</sup>, P. Lenzi<sup>a,b</sup>, M. Meschini<sup>a</sup>, S. Paoletti<sup>a</sup>, G. Sguazzoni<sup>a</sup>, A. Tropiano<sup>a,b</sup>

**INFN Laboratori Nazionali di Frascati, Frascati, Italy**

L. Benussi, S. Bianco, F. Fabbri, D. Piccolo

**INFN Sezione di Genova <sup>a</sup>, Università di Genova <sup>b</sup>, Genova, Italy**

R. Ferretti<sup>a,b</sup>, F. Ferro<sup>a</sup>, M. Lo Vetere<sup>a,b</sup>, E. Robutti<sup>a</sup>, S. Tosi<sup>a,b</sup>

**INFN Sezione di Milano-Bicocca <sup>a</sup>, Università di Milano-Bicocca <sup>b</sup>, Milano, Italy**

M.E. Dinardo<sup>a,b</sup>, S. Fiorendi<sup>a,b</sup>, S. Gennai<sup>a, textsuperscript2</sup>, R. Gerosa<sup>a,b, textsuperscript2</sup>, A. Ghezzi<sup>a,b</sup>, P. Govoni<sup>a,b</sup>, M.T. Lucchini<sup>a,b, textsuperscript2</sup>, S. Malvezzi<sup>a</sup>, R.A. Manzoni<sup>a,b</sup>, A. Martelli<sup>a,b</sup>, B. Marzocchi<sup>a,b</sup>, D. Menasce<sup>a</sup>, L. Moroni<sup>a</sup>, M. Paganoni<sup>a,b</sup>, D. Pedrini<sup>a</sup>, S. Ragazzi<sup>a,b</sup>, N. Redaelli<sup>a</sup>, T. Tabarelli de Fatis<sup>a,b</sup>

**INFN Sezione di Napoli <sup>a</sup>, Università di Napoli 'Federico II' <sup>b</sup>, Università della Basilicata (Potenza) <sup>c</sup>, Università G. Marconi (Roma) <sup>d</sup>, Napoli, Italy**

S. Buontempo<sup>a</sup>, N. Cavallo<sup>a,c</sup>, S. Di Guida<sup>a,d, textsuperscript2</sup>, F. Fabozzi<sup>a,c</sup>, A.O.M. Iorio<sup>a,b</sup>, L. Lista<sup>a</sup>, S. Meola<sup>a,d, textsuperscript2</sup>, M. Merola<sup>a</sup>, P. Paolucci<sup>a, textsuperscript2</sup>

**INFN Sezione di Padova <sup>a</sup>, Università di Padova <sup>b</sup>, Università di Trento (Trento) <sup>c</sup>, Padova, Italy**

P. Azzi<sup>a</sup>, N. Bacchetta<sup>a</sup>, D. Bisello<sup>a,b</sup>, A. Branca<sup>a,b</sup>, R. Carlin<sup>a,b</sup>, P. Checchia<sup>a</sup>, M. Dall'Osso<sup>a,b</sup>, T. Dorigo<sup>a</sup>, M. Galanti<sup>a,b</sup>, U. Gasparini<sup>a,b</sup>, P. Giubilato<sup>a,b</sup>, A. Gozzelino<sup>a</sup>, K. Kanishchev<sup>a,c</sup>, S. Lacaprara<sup>a</sup>, M. Margoni<sup>a,b</sup>, A.T. Meneguzzo<sup>a,b</sup>, J. Pazzini<sup>a,b</sup>, N. Pozzobon<sup>a,b</sup>, P. Ronchese<sup>a,b</sup>, F. Simonetto<sup>a,b</sup>,

E. Torassa<sup>a</sup>, M. Tosi<sup>a,b</sup>, S. Vanini<sup>a,b</sup>, S. Ventura<sup>a</sup>,  
P. Zotto<sup>a,b</sup>, A. Zucchetta<sup>a,b</sup>, G. Zumerle<sup>a,b</sup>

**INFN Sezione di Pavia<sup>a</sup>, Università di Pavia<sup>b</sup>,  
Pavia, Italy**

M. Gabusi<sup>a,b</sup>, S.P. Ratti<sup>a,b</sup>, V. Re<sup>a</sup>, C. Riccardi<sup>a,b</sup>,  
P. Salvini<sup>a</sup>, P. Vitulo<sup>a,b</sup>

**INFN Sezione di Perugia<sup>a</sup>, Università di Perugia<sup>b</sup>,  
Perugia, Italy**

M. Biasini<sup>a,b</sup>, G.M. Bilei<sup>a</sup>, D. Ciangottini<sup>a,b</sup>,  
L. Fanò<sup>a,b</sup>, P. Lariccia<sup>a,b</sup>, G. Mantovani<sup>a,b</sup>,  
M. Menichelli<sup>a</sup>, A. Saha<sup>a</sup>, A. Santocchia<sup>a,b</sup>,  
A. Spiezia<sup>a,b</sup>, <sup>textsuperscript2</sup>

**INFN Sezione di Pisa<sup>a</sup>, Università di Pisa<sup>b</sup>,  
Scuola Normale Superiore di Pisa<sup>c</sup>, Pisa, Italy**

K. Androsov<sup>a</sup>, <sup>textsuperscript25</sup>, P. Azzurri<sup>a</sup>,  
G. Bagliesi<sup>a</sup>, J. Bernardini<sup>a</sup>, T. Boccali<sup>a</sup>, G. Broccolo<sup>a,c</sup>,  
R. Castaldi<sup>a</sup>, M.A. Ciocci<sup>a</sup>, <sup>textsuperscript25</sup>,  
R. Dell'Orso<sup>a</sup>, S. Donato<sup>a,c</sup>, F. Fiori<sup>a,c</sup>, L. Foà<sup>a,c</sup>,  
A. Giassi<sup>a</sup>, M.T. Grippo<sup>a</sup>, <sup>textsuperscript25</sup>,  
F. Ligabue<sup>a,c</sup>, T. Lomtadze<sup>a</sup>, L. Martini<sup>a,b</sup>,  
A. Messineo<sup>a,b</sup>, C.S. Moon<sup>a</sup>, <sup>textsuperscript26</sup>,  
F. Palla<sup>a</sup>, <sup>textsuperscript2</sup>, A. Rizzi<sup>a,b</sup>, A. Savoy-  
Navarro<sup>a</sup>, <sup>textsuperscript27</sup>, A.T. Serban<sup>a</sup>, P. Spagnolo<sup>a</sup>,  
P. Squillacioti<sup>a</sup>, <sup>textsuperscript25</sup>, R. Tenchini<sup>a</sup>,  
G. Tonelli<sup>a,b</sup>, A. Venturi<sup>a</sup>, P.G. Verdini<sup>a</sup>,  
C. Vernieri<sup>a,c</sup>, <sup>textsuperscript2</sup>

**INFN Sezione di Roma<sup>a</sup>, Università di Roma<sup>b</sup>,  
Roma, Italy**

L. Barone<sup>a,b</sup>, F. Cavallari<sup>a</sup>, G. D'imperio<sup>a,b</sup>,  
D. Del Re<sup>a,b</sup>, M. Diemoz<sup>a</sup>, C. Jorda<sup>a</sup>,  
E. Longo<sup>a,b</sup>, F. Margaroli<sup>a,b</sup>, P. Meridiani<sup>a</sup>,  
F. Micheli<sup>a,b</sup>, <sup>textsuperscript2</sup>, S. Nourbakhsh<sup>a,b</sup>,  
G. Organtini<sup>a,b</sup>, R. Paramatti<sup>a</sup>, S. Rahatlou<sup>a,b</sup>,  
C. Rovelli<sup>a</sup>, F. Santanastasio<sup>a,b</sup>,  
L. Soffi<sup>a,b</sup>, <sup>textsuperscript2</sup>, P. Traczyk<sup>a,b</sup>

**INFN Sezione di Torino<sup>a</sup>, Università di Torino<sup>b</sup>,  
Università del Piemonte Orientale (Novara)<sup>c</sup>,  
Torino, Italy**

N. Amapane<sup>a,b</sup>, R. Arcidiacono<sup>a,c</sup>, S. Argiro<sup>a,b</sup>,  
M. Arneodo<sup>a,c</sup>, R. Bellan<sup>a,b</sup>, C. Biino<sup>a</sup>, N. Cartiglia<sup>a</sup>,  
S. Casasso<sup>a,b</sup>, <sup>textsuperscript2</sup>, M. Costa<sup>a,b</sup>,  
A. Degano<sup>a,b</sup>, N. Demaria<sup>a</sup>, L. Finco<sup>a,b</sup>, C. Mariotti<sup>a</sup>,  
S. Maselli<sup>a</sup>, E. Migliore<sup>a,b</sup>, V. Monaco<sup>a,b</sup>, M. Musich<sup>a</sup>,  
M.M. Obertino<sup>a,c</sup>, <sup>textsuperscript2</sup>, G. Ortona<sup>a,b</sup>,  
L. Pacher<sup>a,b</sup>, N. Pastrone<sup>a</sup>, M. Pelliccioni<sup>a</sup>, G.L. Pinna  
Angioni<sup>a,b</sup>, A. Potenza<sup>a,b</sup>, A. Romero<sup>a,b</sup>, M. Ruspa<sup>a,c</sup>,  
R. Sacchi<sup>a,b</sup>, A. Solano<sup>a,b</sup>, A. Staiano<sup>a</sup>, U. Tamponi<sup>a</sup>

**INFN Sezione di Trieste<sup>a</sup>, Università di Trieste<sup>b</sup>,  
Trieste, Italy**

S. Belforte<sup>a</sup>, V. Candelise<sup>a,b</sup>, M. Casarsa<sup>a</sup>, F. Cossutti<sup>a</sup>,  
G. Della Ricca<sup>a,b</sup>, B. Gobbo<sup>a</sup>, C. La Licata<sup>a,b</sup>,  
M. Marone<sup>a,b</sup>, A. Schizzi<sup>a,b</sup>, T. Umer<sup>a,b</sup>, A. Zanetti<sup>a</sup>

**Kangwon National University, Chunchon, Korea**  
S. Chang, A. Kropivnitskaya, S.K. Nam

**Kyungpook National University, Daegu, Korea**  
D.H. Kim, G.N. Kim, M.S. Kim, D.J. Kong, S. Lee,  
Y.D. Oh, H. Park, A. Sakharov, D.C. Son

**Chonbuk National University, Jeonju, Korea**  
T.J. Kim

**Chonnam National University, Institute for Uni-  
verse and Elementary Particles, Kwangju, Korea**  
J.Y. Kim, S. Song

**Korea University, Seoul, Korea**

S. Choi, D. Gyun, B. Hong, M. Jo, H. Kim, Y. Kim,  
B. Lee, K.S. Lee, S.K. Park, Y. Roh

**University of Seoul, Seoul, Korea**

M. Choi, J.H. Kim, I.C. Park, G. Ryu, M.S. Ryu

**Sungkyunkwan University, Suwon, Korea**

Y. Choi, Y.K. Choi, J. Goh, D. Kim, E. Kwon, J. Lee,  
H. Seo, I. Yu

**Vilnius University, Vilnius, Lithuania**

A. Juodagalvis

**National Centre for Particle Physics, Universiti  
Malaya, Kuala Lumpur, Malaysia**

J.R. Komaragiri, M.A.B. Md Ali

**Centro de Investigacion y de Estudios Avanzados  
del IPN, Mexico City, Mexico**

E. Casimiro Linares, H. Castilla-Valdez, E. De La  
Cruz-Burelo, I. Heredia-de La Cruz<sup>textsuperscript28</sup>,  
A. Hernandez-Almada, R. Lopez-Fernandez,  
A. Sanchez-Hernandez

**Universidad Iberoamericana, Mexico City, Mex-  
ico**

S. Carrillo Moreno, F. Vazquez Valencia

**Benemerita Universidad Autonoma de Puebla,  
Puebla, Mexico**

I. Pedraza, H.A. Salazar Ibarguen

**Universidad Autónoma de San Luis Potosí, San  
Luis Potosí, Mexico**

A. Morelos Pineda

**University of Auckland, Auckland, New Zealand**

D. Krofcheck

**University of Canterbury, Christchurch, New  
Zealand**

P.H. Butler, S. Reucroft

**National Centre for Physics, Quaid-I-Azam Uni-  
versity, Islamabad, Pakistan**

A. Ahmad, M. Ahmad, Q. Hassan, H.R. Hoorani,  
W.A. Khan, T. Khurshid, M. Shoaib

**National Centre for Nuclear Research, Swierk,  
Poland**

H. Bialkowska, M. Bluj, B. Boimska, T. Frueboes,  
M. Górski, M. Kazana, K. Nawrocki, K. Romanowska-  
Rybinska, M. Szleper, P. Zalewski

**Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**

G. Brona, K. Bunkowski, M. Cwiok, W. Dominik, K. Doroba, A. Kalinowski, M. Konecki, J. Krolikowski, M. Misiura, M. Olszewski, W. Wolszczak

**Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**

P. Bargassa, C. Beirão Da Cruz E Silva, P. Faccioli, P.G. Ferreira Parracho, M. Gallinaro, L. Lloret Iglesias, F. Nguyen, J. Rodrigues Antunes, J. Seixas, J. Varela, P. Vischia

**Joint Institute for Nuclear Research, Dubna, Russia**

S. Afanasiev, P. Bunin, I. Golutvin, V. Karjavin, V. Konoplyanikov, G. Kozlov, A. Lanev, A. Malakhov, V. Matveev<sup>29</sup>, P. Moisenz, V. Palichik, V. Perelygin, M. Savina, S. Shmatov, S. Shulha, N. Skatchkov, V. Smirnov, A. Zarubin

**Petersburg Nuclear Physics Institute, Gatchina (St. Petersburg), Russia**

V. Golovtsov, Y. Ivanov, V. Kim<sup>30</sup>, P. Levchenko, V. Murzin, V. Oreshkin, I. Smirnov, V. Sulimov, L. Uvarov, S. Vavilov, A. Vorobyev, An. Vorobyev

**Institute for Nuclear Research, Moscow, Russia**

Yu. Andreev, A. Dermenev, S. Gninenko, N. Golubev, M. Kirsanov, N. Krasnikov, A. Pashenkov, D. Tlisov, A. Toropin

**Institute for Theoretical and Experimental Physics, Moscow, Russia**

V. Epshteyn, V. Gavrilov, N. Lychkovskaya, V. Popov, I. Pozdnyakov, G. Safronov, S. Semenov, A. Spiridonov, V. Stolin, E. Vlasov, A. Zhokin

**P.N. Lebedev Physical Institute, Moscow, Russia**

V. Andreev, M. Azarkin, I. Dremin, M. Kirakosyan, A. Leonidov, G. Mesyats, S.V. Rusakov, A. Vinogradov

**Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia**

A. Belyaev, E. Boos, V. Bunichev, M. Dubinin<sup>31</sup>, L. Dudko, A. Ershov, A. Gribushin, V. Klyukhin, O. Kodolova, I. Lokhtin, S. Obraztsov, V. Savrin, A. Snigirev

**State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, Russia**

I. Azhgirey, I. Bayshev, S. Bitioukov, V. Kachanov, A. Kalinin, D. Konstantinov, V. Krychkin, V. Petrov, R. Ryutin, A. Sobol, L. Tourtchanovitch, S. Troshin, N. Tyurin, A. Uzunian, A. Volkov

**University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia**

P. Adzic<sup>32</sup>, M. Ekmedzic, J. Milosevic, V. Rekovic

**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**

J. Alcaraz Maestre, C. Battilana, E. Calvo, M. Cerrada, M. Chamizo Llatas, N. Colino, B. De La Cruz, A. Delgado Peris, D. Domínguez Vázquez, A. Escalante Del Valle, C. Fernandez Bedoya, J.P. Fernández Ramos, J. Flix, M.C. Fouz, P. Garcia-Abia, O. Gonzalez Lopez, S. Goy Lopez, J.M. Hernandez, M.I. Josa, E. Navarro De Martino, A. Pérez-Calero Yzquierdo, J. Puerta Pelayo, A. Quintario Olmeda, I. Redondo, L. Romero, M.S. Soares

**Universidad Autónoma de Madrid, Madrid, Spain**

C. Albajar, J.F. de Trocóniz, M. Missiroli, D. Moran

**Universidad de Oviedo, Oviedo, Spain**

H. Brun, J. Cuevas, J. Fernandez Menendez, S. Folgueras, I. Gonzalez Caballero

**Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain**

J.A. Brochero Cifuentes, I.J. Cabrillo, A. Calderon, J. Duarte Campderros, M. Fernandez, G. Gomez, A. Graziano, A. Lopez Virto, J. Marco, R. Marco, C. Martinez Rivero, F. Matorras, F.J. Munoz Sanchez, J. Piedra Gomez, T. Rodrigo, A.Y. Rodríguez-Marrero, A. Ruiz-Jimeno, L. Scodellaro, I. Vila, R. Vilar Cortabitarte

**CERN, European Organization for Nuclear Research, Geneva, Switzerland**

D. Abbaneo, E. Auffray, G. Auzinger, M. Bachtis, P. Baillon, A.H. Ball, D. Barney, A. Benaglia, J. Bendavid, L. Benhabib, J.F. Benitez, C. Bernett<sup>7</sup>, P. Bloch, A. Bocci, A. Bonato, O. Bondu, C. Botta, H. Breuker, T. Camporesi, G. Cerminara, S. Colafranceschi<sup>33</sup>, M. D'Alfonso, D. d'Enterria, A. Dabrowski, A. David, F. De Guio, A. De Roeck, S. De Visscher, E. Di Marco, M. Dobson, M. Dordevic, N. Dupont-Sagorin, A. Elliott-Peisert, J. Eugster, G. Franzoni, W. Funk, D. Gigi, K. Gill, D. Giordano, M. Girone, F. Glege, R. Guida, S. Gundacker, M. Guthoff, J. Hammer, M. Hansen, P. Harris, J. Hegeman, V. Innocente, P. Janot, K. Kousouris, K. Krajczar, P. Lecoq, C. Lourenço, N. Magini, L. Malgeri, M. Mannelli, J. Marrouche, L. Masetti, F. Meijers, S. Mersi, E. Meschi, F. Moortgat, S. Morovic, M. Mulders, P. Musella, L. Orsini, L. Pape, E. Perez, L. Perrozzi, A. Petrilli, G. Petrucciani, A. Pfeiffer, M. Pierini, M. Pimiä, D. Piparo, M. Plagge, A. Racz,

G. Rolanditextsuperscript34, M. Rovere, H. Sakulin, C. Schäfer, C. Schwick, A. Sharma, P. Siegrist, P. Silva, M. Simon, P. Sphicastextsuperscript35, D. Spiga, J. Steggemann, B. Stieger, M. Stoye, Y. Takahashi, D. Treille, A. Tsirou, G.I. Verestextsuperscript17, N. Wardle, H.K. Wöhri, H. Wollny, W.D. Zeuner

**Paul Scherrer Institut, Villigen, Switzerland**

W. Bertl, K. Deiters, W. Erdmann, R. Horisberger, Q. Ingram, H.C. Kaestli, D. Kotlinski, U. Langenegger, D. Renker, T. Rohe

**Institute for Particle Physics, ETH Zurich, Zurich, Switzerland**

F. Bachmair, L. Bäni, L. Bianchini, M.A. Buchmann, B. Casal, N. Chanon, G. Dissertori, M. Dittmar, M. Donegà, M. Dünser, P. Eller, C. Grab, D. Hits, J. Hoss, W. Lustermann, B. Mangano, A.C. Marini, P. Martinez Ruiz del Arbol, M. Masciovecchio, D. Meister, N. Mohr, C. Nägelitextsuperscript36, F. Nessi-Tedaldi, F. Pandolfi, F. Pauss, M. Peruzzi, M. Quittnat, L. Rebane, M. Rossini, A. Starodumovtextsuperscript37, M. Takahashi, K. Theofilatos, R. Wallny, H.A. Weber

**Universität Zürich, Zurich, Switzerland**

C. Amslertextsuperscript38, M.F. Canelli, V. Chiochia, A. De Cosa, A. Hinzmann, T. Hreus, B. Kilminster, C. Lange, B. Millan Mejias, J. Ngadiuba, P. Robmann, F.J. Ronga, S. Taroni, M. Verzetti, Y. Yang

**National Central University, Chung-Li, Taiwan**

M. Cardaci, K.H. Chen, C. Ferro, C.M. Kuo, W. Lin, Y.J. Lu, R. Volpe, S.S. Yu

**National Taiwan University (NTU), Taipei, Taiwan**

P. Chang, Y.H. Chang, Y.W. Chang, Y. Chao, K.F. Chen, P.H. Chen, C. Dietz, U. Grundler, W.-S. Hou, K.Y. Kao, Y.F. Liu, R.-S. Lu, D. Majumder, E. Petrakou, Y.M. Tzeng, R. Wilken

**Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand**

B. Asavapibhop, G. Singh, N. Srimanobhas, N. Suwonjandee

**Cukurova University, Adana, Turkey**

A. Adiguzel, M.N. Bakircitextsuperscript39, S. Cercitextsuperscript40, C. Dozen, I. Dumanoglu, E. Eskut, S. Girgis, G. Gokbulut, E. Gurpinar, I. Hos, E.E. Kangal, A. Kayis Topaksu, G. Onengutttextsuperscript41, K. Ozdemir, S. Ozturktextsuperscript39, A. Polatoz, D. Sunar Cercitextsuperscript40, B. Talitextsuperscript40, H. Topaklitextsuperscript39, M. Vergili

**Middle East Technical University, Physics Department, Ankara, Turkey**

I.V. Akin, B. Bilin, S. Bilmis,

H. Gamsizkantextsuperscript42, B. Isildaktextsuperscript43, G. Karapinantextsuperscript44, K. Ocalantextsuperscript45, S. Sekmen, U.E. Surat, M. Yalvac, M. Zeyrek

**Bogazici University, Istanbul, Turkey**

E.A. Albayrakttextsuperscript46, E. Gülmez, M. Kayatextsuperscript47, O. Kayatextsuperscript48, T. Yetkintextsuperscript49

**Istanbul Technical University, Istanbul, Turkey**

K. Cankocak, F.I. Vardarli

**National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine**

L. Levchuk, P. Sorokin

**University of Bristol, Bristol, United Kingdom**

J.J. Brooke, E. Clement, D. Cussans, H. Flacher, J. Goldstein, M. Grimes, G.P. Heath, H.F. Heath, J. Jacob, L. Kreczko, C. Lucas, Z. Meng, D.M. Newboldtextsuperscript50, S. Paramesvaran, A. Poll, T. Sakuma, S. Senkin, V.J. Smith, T. Williams

**Rutherford Appleton Laboratory, Didcot, United Kingdom**

K.W. Bell, A. Belyaevtextsuperscript51, C. Brew, R.M. Brown, D.J.A. Cockerill, J.A. Coughlan, K. Harder, S. Harper, E. Olaiya, D. Petyt, C.H. Shepherd-Themistocleous, A. Thea, I.R. Tomalin, W.J. Womersley, S.D. Worm

**Imperial College, London, United Kingdom**

M. Baber, R. Bainbridge, O. Buchmuller, D. Burton, D. Colling, N. Cripps, M. Cutajar, P. Dauncey, G. Davies, M. Della Negra, P. Dunne, W. Ferguson, J. Fulcher, D. Fuytan, G. Hall, G. Iles, M. Jarvis, G. Karapostoli, M. Kenzie, R. Lane, R. Lucastextsuperscript50, L. Lyons, A.-M. Magnan, S. Malik, B. Mathias, J. Nash, A. Nikitenkotextsuperscript37, J. Pela, M. Pesaresi, K. Petridis, D.M. Raymond, S. Rogerson, A. Rose, C. Seez, P. Sharp<sup>†</sup>, A. Tapper, M. Vazquez Acosta, T. Virdee, S.C. Zenz

**Brunel University, Uxbridge, United Kingdom**

J.E. Cole, P.R. Hobson, A. Khan, P. Kyberd, D. Leggat, D. Leslie, W. Martin, I.D. Reid, P. Symonds, L. Teodorescu, M. Turner

**Baylor University, Waco, USA**

J. Dittmann, K. Hatakeyama, A. Kasmi, H. Liu, T. Scarborough

**The University of Alabama, Tuscaloosa, USA**

O. Charaf, S.I. Cooper, C. Henderson, P. Rumerio

**Boston University, Boston, USA**

A. Avetisyan, T. Bose, C. Fantasia, P. Lawson, C. Richardson, J. Rohlf, J. St. John, L. Sulak

**Brown University, Providence, USA**

J. Alimena, E. Berry, S. Bhattacharya, G. Christopher, D. Cutts, Z. Demiragli, N. Dhingra, A. Ferapontov, A. Garabedian, U. Heintz, G. Kukartsev, E. Laird, G. Landsberg, M. Luk, M. Narain, M. Segala, T. Sinthuprasith, T. Speer, J. Swanson

**University of California, Davis, USA**

R. Breedon, G. Breto, M. Calderon De La Barca Sanchez, S. Chauhan, M. Chertok, J. Conway, R. Conway, P.T. Cox, R. Erbacher, M. Gardner, W. Ko, R. Lander, T. Miceli, M. Mulhearn, D. Pellett, J. Pilot, F. Ricci-Tam, M. Searle, S. Shalhout, J. Smith, M. Squires, D. Stolp, M. Tripathi, S. Wilbur, R. Yohay

**University of California, Los Angeles, USA**

R. Cousins, P. Everaerts, C. Farrell, J. Hauser, M. Ignatenko, G. Rakness, E. Takasugi, V. Valuev, M. Weber

**University of California, Riverside, Riverside, USA**

K. Burt, R. Clare, J. Ellison, J.W. Gary, G. Hanson, J. Heilman, M. Ivova Rikova, P. Jandir, E. Kennedy, F. Lacroix, O.R. Long, A. Luthra, M. Malberti, M. Olmedo Negrete, A. Shrinivas, S. Sumowidagdo, S. Wimpenny

**University of California, San Diego, La Jolla, USA**

J.G. Branson, G.B. Cerati, S. Cittolin, R.T. D'Agnolo, A. Holzner, R. Kelley, D. Klein, J. Letts, I. Macneill, D. Olivito, S. Padhi, C. Palmer, M. Pieri, M. Sani, V. Sharma, S. Simon, E. Sudano, M. Tadel, Y. Tu, A. Vartak, C. Welke, F. Würthwein, A. Yagil

**University of California, Santa Barbara, Santa Barbara, USA**

D. Barge, J. Bradmiller-Feld, C. Campagnari, T. Danielson, A. Dishaw, V. Dutta, K. Flowers, M. Franco Sevilla, P. Geffert, C. George, F. Golf, L. Gouskos, J. Incandela, C. Justus, N. Mccoll, J. Richman, D. Stuart, W. To, C. West, J. Yoo

**California Institute of Technology, Pasadena, USA**

A. Apresyan, A. Bornheim, J. Bunn, Y. Chen, J. Duarte, A. Mott, H.B. Newman, C. Pena, C. Rogan, M. Spiropulu, V. Timciuc, J.R. Vlimant, R. Wilkinson, S. Xie, R.Y. Zhu

**Carnegie Mellon University, Pittsburgh, USA**

V. Azzolini, A. Calamba, B. Carlson, T. Ferguson, Y. Iiyama, M. Paulini, J. Russ, H. Vogel, I. Vorobiev

**University of Colorado at Boulder, Boulder, USA**

J.P. Cumalat, W.T. Ford, A. Gaz, M. Krohn, E. Luiggi Lopez, U. Nauenberg, J.G. Smith, K. Stenson, K.A. Ulmer, S.R. Wagner

**Cornell University, Ithaca, USA**

J. Alexander, A. Chatterjee, J. Chaves, J. Chu, S. Dittmer, N. Eggert, N. Mirman, G. Nicolas Kaufman, J.R. Patterson, A. Ryd, E. Salvati, L. Skinnari, W. Sun,

W.D. Teo, J. Thom, J. Thompson, J. Tucker, Y. Weng, L. Winstrom, P. Wittich

**Fairfield University, Fairfield, USA**

D. Winn

**Fermi National Accelerator Laboratory, Batavia, USA**

S. Abdullin, M. Albrow, J. Anderson, G. Apollinari, L.A.T. Bauerdick, A. Beretvas, J. Berryhill, P.C. Bhat, G. Bolla, K. Burkett, J.N. Butler, H.W.K. Cheung, F. Chlebana, S. Cihangir, V.D. Elvira, I. Fisk, J. Freeman, Y. Gao, E. Gottschalk, L. Gray, D. Green, S. Grünendahl, O. Gutsche, J. Hanlon, D. Hare, R.M. Harris, J. Hirschauer, B. Hooberman, S. Jindariani, M. Johnson, U. Joshi, K. Kaadze, B. Klima, B. Kreis, S. Kwan, J. Linacre, D. Lincoln, R. Lipton, T. Liu, J. Lykken, K. Maeshima, J.M. Marraffino, V.I. Martinez Outschoorn, S. Maruyama, D. Mason, P. McBride, P. Merkel, K. Mishra, S. Mrenna, Y. Musienkotextsuperscript29, S. Nahn, C. Newman-Holmes, V. O'Dell, O. Prokofyev, E. Sexton-Kennedy, S. Sharma, A. Soha, W.J. Spalding, L. Spiegel, L. Taylor, S. Tkaczyk, N.V. Tran, L. Uplegger, E.W. Vaandering, R. Vidal, A. Whitbeck, J. Whitmore, F. Yang

**University of Florida, Gainesville, USA**

D. Acosta, P. Avery, P. Bortignon, D. Bourilkov, M. Carver, D. Curry, S. Das, M. De Gruttola, G.P. Di Giovanni, R.D. Field, M. Fisher, I.K. Furic, J. Hugon, J. Konigsberg, A. Korytov, T. Kypreos, J.F. Low, K. Matchev, P. Milenovictextsuperscript52, G. Mitselmakher, L. Muniz, A. Rinkevicius, L. Shchutska, M. Snowball, D. Sperka, J. Yelton, M. Zakaria

**Florida International University, Miami, USA**

S. Hewamanage, S. Linn, P. Markowitz, G. Martinez, J.L. Rodriguez

**Florida State University, Tallahassee, USA**

T. Adams, A. Askew, J. Bochenek, B. Diamond, J. Haas, S. Hagopian, V. Hagopian, K.F. Johnson, H. Prosper, V. Veeraraghavan, M. Weinberg

**Florida Institute of Technology, Melbourne, USA**

M.M. Baarmand, M. Hohmann, H. Kalakhety, F. Yumiceva

**University of Illinois at Chicago (UIC), Chicago, USA**

M.R. Adams, L. Apanasevich, V.E. Bazterra, D. Berry, R.R. Betts, I. Bucinskaite, R. Cavanaugh, O. Evdokimov, L. Gauthier, C.E. Gerber, D.J. Hofman, S. Khalatyan, P. Kurt, D.H. Moon, C. O'Brien, C. Silkworth, P. Turner, N. Varelas

**The University of Iowa, Iowa City, USA**

B. Bilkitextsuperscript53, W. Clarida, K. Dilisiz, F. Duru, M. Haytmyradov, J.-P. Merlo, H. Mermerkayatextsuperscript54, A. Mestvirishvili,

A. Moeller, J. Nachtman, H. Ogul, Y. Onel, F. Ozoktextsuperscript46, A. Penzo, R. Rahmat, S. Sen, P. Tan, E. Tiras, J. Wetzel, K. Yi

**Johns Hopkins University, Baltimore, USA**

B.A. Barnett, B. Blumenfeld, S. Bolognesi, D. Fehling, A.V. Gritsan, P. Maksimovic, C. Martin, M. Swartz

**The University of Kansas, Lawrence, USA**

P. Baringer, A. Bean, G. Benelli, C. Bruner, R.P. Kenny III, M. Malek, M. Murray, D. Noonan, S. Sanders, J. Sekaric, R. Stringer, Q. Wang, J.S. Wood

**Kansas State University, Manhattan, USA**

I. Chakaberia, A. Ivanov, S. Khalil, M. Makouski, Y. Maravin, L.K. Saini, S. Shrestha, N. Skhirtladze, I. Svintrazde

**Lawrence Livermore National Laboratory, Livermore, USA**

J. Gronberg, D. Lange, F. Rebassoo, D. Wright

**University of Maryland, College Park, USA**

A. Baden, A. Belloni, B. Calvert, S.C. Eno, J.A. Gomez, N.J. Hadley, R.G. Kellogg, T. Kolberg, Y. Lu, M. Marionneau, A.C. Mignerey, K. Pedro, A. Skuja, M.B. Tonjes, S.C. Tonwar

**Massachusetts Institute of Technology, Cambridge, USA**

A. Apyan, R. Barbieri, G. Bauer, W. Busza, I.A. Cali, M. Chan, L. Di Matteo, G. Gomez Ceballos, M. Goncharov, D. Gulhan, M. Klute, Y.S. Lai, Y.-J. Lee, A. Levin, P.D. Luckey, T. Ma, C. Paus, D. Ralph, C. Roland, G. Roland, G.S.F. Stephans, F. Stöckli, K. Sumorok, D. Velicanu, J. Veverka, B. Wyslouch, M. Yang, M. Zanetti, V. Zhukova

**University of Minnesota, Minneapolis, USA**

B. Dahmes, A. Gude, S.C. Kao, K. Klapoetke, Y. Kubota, J. Mans, N. Pastika, R. Rusack, A. Singovsky, N. Tambe, J. Turkewitz

**University of Mississippi, Oxford, USA**

J.G. Acosta, S. Oliveros

**University of Nebraska-Lincoln, Lincoln, USA**

E. Avdeeva, K. Bloom, S. Bose, D.R. Claes, A. Dominguez, R. Gonzalez Suarez, J. Keller, D. Knowlton, I. Kravchenko, J. Lazo-Flores, S. Malik, F. Meier, F. Ratnikov, G.R. Snow, M. Zvada

**State University of New York at Buffalo, Buffalo, USA**

J. Dolen, A. Godshalk, I. Iashvili, A. Kharchilava, A. Kumar, S. Rappocci

**Northeastern University, Boston, USA**

G. Alverson, E. Barberis, D. Baumgartel, M. Chasco, J. Haley, A. Massironi, D.M. Morse, D. Nash, T. Orioto, D. Trocino, R.-J. Wang, D. Wood, J. Zhang

**Northwestern University, Evanston, USA**

K.A. Hahn, A. Kubik, N. Mucia, N. Odell, B. Pollack, A. Pozdnyakov, M. Schmitt, S. Stoynev, K. Sung, M. Velasco, S. Won

**University of Notre Dame, Notre Dame, USA**

A. Brinkerhoff, K.M. Chan, A. Drozdetskiy, M. Hildreth, C. Jessop, D.J. Karmgard, N. Kellams, K. Lannon, W. Luo, S. Lynch, N. Marinelli, T. Pearson, M. Planer, R. Ruchti, N. Valls, M. Wayne, M. Wolf, A. Woodard

**The Ohio State University, Columbus, USA**

L. Antonelli, J. Brinson, B. Bylsma, L.S. Durkin, S. Flowers, A. Hart, C. Hill, R. Hughes, K. Kotov, T.Y. Ling, D. Puigh, M. Rodenburg, G. Smith, B.L. Winer, H. Wolfe, H.W. Wulsin

**Princeton University, Princeton, USA**

O. Driga, P. Elmer, J. Hardenbrook, P. Hebda, A. Hunt, S.A. Koay, P. Lujan, D. Marlow, T. Medvedeva, M. Mooney, J. Olsen, P. Piroué, X. Quan, H. Saka, D. Sticklandtextsuperscript2, C. Tully, J.S. Werner, A. Zuranski

**University of Puerto Rico, Mayaguez, USA**

E. Brownson, H. Mendez, J.E. Ramirez Vargas

**Purdue University, West Lafayette, USA**

V.E. Barnes, D. Benedetti, D. Bortoletto, M. De Mattia, L. Gutay, Z. Hu, M.K. Jha, M. Jones, K. Jung, M. Kress, N. Leonardo, D. Lopes Pegna, V. Maroussov, D.H. Miller, N. Neumeister, B.C. Radburn-Smith, X. Shi, I. Shipsey, D. Silvers, A. Svyatkovskiy, F. Wang, W. Xie, L. Xu, H.D. Yoo, J. Zablocki, Y. Zheng

**Purdue University Calumet, Hammond, USA**

N. Parashar, J. Stupak

**Rice University, Houston, USA**

A. Adair, B. Akgun, K.M. Ecklund, F.J.M. Geurts, W. Li, B. Michlin, B.P. Padley, R. Redjimi, J. Roberts, J. Zabel

**University of Rochester, Rochester, USA**

B. Betchart, A. Bodek, R. Covarelli, P. de Barbaro, R. Demina, Y. Eshaq, T. Ferbel, A. Garcia-Bellido, P. Goldenzweig, J. Han, A. Harel, A. Khukhunaishvili, S. Korjenevski, G. Petrillo, D. Vishnevskiy

**The Rockefeller University, New York, USA**

R. Ciesielski, L. Demortier, K. Goulios, G. Lungu, C. Mesropian

**Rutgers, The State University of New Jersey, Piscataway, USA**

S. Arora, A. Barker, J.P. Chou, C. Contreras-Campana, E. Contreras-Campana, D. Duggan, D. Ferencek, Y. Gershtein, R. Gray, E. Halkiadakis, D. Hidas, S. Kaplan, A. Lath, S. Panwalkar, M. Park, R. Patel, S. Salur, S. Schnetzer, S. Somalwar, R. Stone, S. Thomas, P. Thomassen, M. Walker

**University of Tennessee, Knoxville, USA**

K. Rose, S. Spanier, A. York

**Texas A&M University, College Station, USA**

O. Bouhalitextsuperscript55, A. Castaneda Hernandez, R. Eusebi, W. Flanagan, J. Gilmore, T. Kamontextsuperscript56, V. Khotilovich, V. Krute-lyov, R. Montalvo, I. Osipenkov, Y. Pakhotin, A. Perloff, J. Roe, A. Rose, A. Safonov, I. Suarez, A. Tatarinov

**Texas Tech University, Lubbock, USA**

N. Akchurin, C. Cowden, J. Damgov, C. Dragoiu, P.R. Duderu, J. Faulkner, K. Kovitanggoon, S. Kunori, S.W. Lee, T. Libeiro, I. Volobouev

**Vanderbilt University, Nashville, USA**

E. Appelt, A.G. Delannoy, S. Greene, A. Gurrola, W. Johns, C. Maguire, Y. Mao, A. Melo, M. Sharma, P. Sheldon, B. Snook, S. Tuo, J. Velkovska

**University of Virginia, Charlottesville, USA**

M.W. Arenton, S. Boutle, B. Cox, B. Francis, J. Goodell, R. Hirosky, A. Ledovskoy, H. Li, C. Lin, C. Neu, J. Wood

**Wayne State University, Detroit, USA**

C. Clarke, R. Harr, P.E. Karchin, C. Kottachchi Kankanamge Don, P. Lamichhane, J. Sturdy

**University of Wisconsin, Madison, USA**

D.A. Belknap, D. Carlsmith, M. Cepeda, S. Dasu, L. Dodd, S. Duric, E. Friis, R. Hall-Wilton, M. Herndon, A. Hervé, P. Klabbers, A. Lanaro, C. Lazaridis, A. Levine, R. Loveless, A. Mohapatra, I. Ojalvo, T. Perry, G.A. Pierro, G. Polese, I. Ross, T. Sarangi, A. Savin, W.H. Smith, D. Taylor, P. Verwilligen, C. Vuosalo, N. Woods

†: Deceased

1: Also at Vienna University of Technology, Vienna, Austria

2: Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

3: Also at Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg, Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France

4: Also at National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

5: Also at Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia

6: Also at Universidade Estadual de Campinas, Campinas, Brazil

7: Also at Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France

8: Also at Joint Institute for Nuclear Research, Dubna, Russia

9: Also at Suez University, Suez, Egypt

10: Also at British University in Egypt, Cairo, Egypt

11: Also at Fayoum University, El-Fayoum, Egypt

12: Also at Ain Shams University, Cairo, Egypt

13: Now at Sultan Qaboos University, Muscat, Oman

14: Also at Université de Haute Alsace, Mulhouse, France

15: Also at Brandenburg University of Technology, Cottbus, Germany

16: Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary

17: Also at Eötvös Loránd University, Budapest, Hungary

18: Also at University of Debrecen, Debrecen, Hungary

19: Also at University of Visva-Bharati, Santiniketan, India

20: Now at King Abdulaziz University, Jeddah, Saudi Arabia

21: Also at University of Ruhuna, Matara, Sri Lanka

22: Also at Isfahan University of Technology, Isfahan, Iran

23: Also at University of Tehran, Department of Engineering Science, Tehran, Iran

24: Also at Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran

25: Also at Università degli Studi di Siena, Siena, Italy

26: Also at Centre National de la Recherche Scientifique (CNRS) - IN2P3, Paris, France

27: Also at Purdue University, West Lafayette, USA

28: Also at Universidad Michoacana de San Nicolas de Hidalgo, Morelia, Mexico

29: Also at Institute for Nuclear Research, Moscow, Russia

30: Also at St. Petersburg State Polytechnical University, St. Petersburg, Russia

31: Also at California Institute of Technology, Pasadena, USA

32: Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia

33: Also at Facoltà Ingegneria, Università di Roma, Roma, Italy

34: Also at Scuola Normale e Sezione dell'INFN, Pisa, Italy

35: Also at University of Athens, Athens, Greece

36: Also at Paul Scherrer Institut, Villigen, Switzerland

37: Also at Institute for Theoretical and Experimental Physics, Moscow, Russia

38: Also at Albert Einstein Center for Fundamental Physics, Bern, Switzerland

39: Also at Gaziosmanpasa University, Tokat, Turkey

40: Also at Adiyaman University, Adiyaman, Turkey

41: Also at Cag University, Mersin, Turkey

42: Also at Anadolu University, Eskisehir, Turkey

43: Also at Ozyegin University, Istanbul, Turkey

- 44: Also at Izmir Institute of Technology, Izmir, Turkey  
45: Also at Necmettin Erbakan University, Konya, Turkey  
46: Also at Mimar Sinan University, Istanbul, Istanbul, Turkey  
47: Also at Marmara University, Istanbul, Turkey  
48: Also at Kafkas University, Kars, Turkey  
49: Also at Yildiz Technical University, Istanbul, Turkey  
50: Also at Rutherford Appleton Laboratory, Didcot, United Kingdom  
51: Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom  
52: Also at University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia  
53: Also at Argonne National Laboratory, Argonne, USA  
54: Also at Erzincan University, Erzincan, Turkey  
55: Also at Texas A&M University at Qatar, Doha, Qatar  
56: Also at Kyungpook National University, Daegu, Korea