



HAL
open science

Spectroscopic characterization of the inclusion complexes between the antifungal drugs Naftifine and Terbinafine and Cyclodextrins

Maite Uzqueda, Gustavo González-Gaitano, Denis Wouessidjewe, Arantza Zornoza, Miguel Sánchez, Carmen Martín, Itziar Velaz

► To cite this version:

Maite Uzqueda, Gustavo González-Gaitano, Denis Wouessidjewe, Arantza Zornoza, Miguel Sánchez, et al.. Spectroscopic characterization of the inclusion complexes between the antifungal drugs Naftifine and Terbinafine and Cyclodextrins. *Supramolecular Chemistry, Taylor & Francis: STM, Behavioural Science and Public Health Titles*, 2009, 21 (08), pp.759-769. 10.1080/10610270902980606. hal-00528957


HAL Id: hal-00528957

<https://hal.archives-ouvertes.fr/hal-00528957>

Submitted on 23 Oct 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




**Spectroscopic characterization of the inclusion complexes
between the antifungal drugs Naftifine and Terbinafine and
Cyclodextrins**

Journal:	<i>Supramolecular Chemistry</i>
Manuscript ID:	GSCH-2008-0177.R1
Manuscript Type:	Full Paper
Date Submitted by the Author:	13-Mar-2009
Complete List of Authors:	Uzqueda, Maite; University of Navarra González-Gaitano, Gustavo; University of Navarra Wouessidjewe, Denis; Université Joseph Fourier, Pharmacotechnie Zornoza, Arantza; University of Navarra Sánchez, Miguel; University of Navarra Martín, Carmen; University of Navarra Velaz, Itziar; University of Navarra, Chemistry and Soil Sciences
Keywords:	Terbinafine, Naftifine, Cyclodextrins, inclusion complexes characterization

Graphical abstract

Complexes of naftifine and terbinafine with cyclodextrins have been characterized by UV, ^1H NMR spectroscopy, ROESY techniques and ESI-MS. Stoichiometry, stability constants and topology of the complexes have been investigated.



M. Uzqueda et al.
Supramolecular Chemistry

RESEARCH ARTICLE

Spectroscopic characterization of the inclusion complexes between the antifungal drugs Naftifine and Terbinafine and Cyclodextrins

M. Uzqueda^a, G. González-Gaitano^a, D. Wouessidjewe^b, A. Zornoza^a, M. Sánchez^a, C.

Martín^a, I. Vélaz^{a,*}

^aDepartment of Chemistry and Soil Sciences (Physical Chemistry). University of Navarra. C/Irunlarrea s/n 31080, Pamplona, Navarra, SPAIN.

^bPharmacotechnie, DPM-UMR 5063 UJF/CNRS, Université Joseph Fourier – Grenoble I, 5 av. de Verdun, 38243 Meylan cedex, FRANCE.

Abstract

The complexation of naftifine (NF) and terbinafine (TB) with cyclodextrins (CDs) has been investigated by UV/visible and ¹H NMR spectroscopy, ROESY techniques and also ESI-MS. Both drugs form 1:1 inclusion complexes with all the cyclodextrins tested except with α-CD, as deduced from the Benesi-Hildebrand plots and confirmed by ESI-MS and by NMR (Job plot method). The K₁₁ values for NF decrease in the order β-CD > Mβ-CD > HPβ-CD > γ-CD. The determination of the enthalpy and entropy provides information about the main driving forces in the process. The stability constants of the complexes NF-βCD, TB-βCD and TB-γCD determined by ¹H NMR are in agreement with the values obtained by UV. For TB-βCD the value is higher, due to the fact that the length of the TB aliphatic chain allows a deeper inclusion of the naphthalene group inside the corresponding β-CD molecule, according to the two dimensional ROESY experiments.

Keywords: Terbinafine; Naftifine; Cyclodextrins; inclusion complexes characterization

*Corresponding author. Email: itzvelaz@unav.es

1. Introduction

Cyclodextrins (CDs) are torus shaped cyclic oligosaccharides made up of α -D-glucopyranose residues linked by glycosidic bonds, the most common are those formed by six, seven or eight glucose units (α -, β - and γ -CD, respectively) (Figure 1). This class of carbohydrates exhibits a doughnut shape, in which the cavity has a hydrophobic character compared to water, whereas the rims, bearing the OH groups, are hydrophilic. This structure makes CDs capable of forming inclusion compounds with a variety of guest molecules both in solution and in the solid state (1). The first condition required for a molecule to form an inclusion complex with CD is to fit in the cavity, either completely or partially. In addition, a favourable energetic balance is required, which depends on the nature of the guest, the inner diameter of the CD, and its substitution degree (2). The formation of these inclusion compounds has been widely used to improve the aqueous solubility of poorly soluble drugs, together with their bioavailability, dissolution rate, permeability and stability (3). Natural CDs can be modified by substitution of the hydroxyl groups by methyl or hydroxypropyl groups in order to improve properties such as the solubility or to avoid undesired effects. 2-Hydroxypropyl- β -cyclodextrin (HP β -CD) is a hydroxyalkyl β -cyclodextrin derivative which has been studied most thoroughly. It is highly hydrophilic and generally forms complexes with many poorly water-soluble substances. In addition, both β -CD and HP β -CD can be safely used in oral and parenteral formulations. The preparation of the methylated CD (M β -CD) is relatively simple and inexpensive. Both CD derivatives (M β - and HP β -CD) are found to be permeable into the skin, particularly when they are applied under the occlusive-dressing conditions and/or by using vehicles containing absorption-promoting agents (4,5).

In this study, naftifine (NF) and terbinafine (TB) have been selected as guests to characterize their respective complexes with CDs. Naftifine, (E)-N-methyl-N-(1-naphthylmethyl)-3-phenyl-2-propen-1-amine ($pK_{aNF} = 8.0 \pm 0.2$) and terbinafine, (E)-N-(6,6-dimethyl-2-hepten-4-ynyl)-N-methyl-1-naphthalene methanamine ($pK_{aTB} = 6.7 \pm 0.3$) hydrochlorides, are two antifungal agents of the chemical group of allylamines (Figure 2). They present a high selectivity for a single fungal enzyme and an apparent lack of interferences with other enzymes of fungal and mammalian origin (6). Their mechanism of action is based on the blocking of the biosynthesis of ergosterol, which is an essential component of fungal cell membranes (7). TB is indicated for both oral and topical treatment of mycoses and NF only for topical treatment. These drugs present gastrointestinal adverse effects; which are expected to be reduced or avoided by microencapsulation in CDs.

1
2
3 The study of complexation with cyclodextrins is usually focused on two main objectives: the
4 accurate determination of the stability constants and the structural elucidation of the complex.
5
6 There is a wide variety of techniques that can be used to determine the stability constants (1).
7
8 The accuracy of the constant is improved by the use of different methods in its calculation,
9
10 although disagreements between techniques are not uncommon.

11
12 In relation to the study of the mode of inclusion or topology of the complex, $^1\text{H-NMR}$ has
13 become the most important method for structural elucidation of inclusion complexes with
14 CDs, particularly in solution. For structural studies, the Nuclear Overhauser Effect (NOE) is
15 one of the most important tools for the study of the topology of the complexes (8), because the
16 dipolar interactions that are detected as NOE enhancements reflect the spatial proximity
17 between protons. For medium-sized systems, such as those studied here, the appropriate
18 technique is rotating frame Overhauser effect spectroscopy (ROESY) (9).

19
20 The aim of this paper is to study the inclusion process of NF and TB with natural and
21 derivative CDs by different techniques. This characterization includes the determination of
22 the stability constants, stoichiometries and the corresponding thermodynamic parameters
23 together with the structural elucidation of the complexes.
24
25
26
27
28
29
30
31
32

33 2. Experimental

34 2.1. Materials

35
36 NF hydrochloride (molecular weight 323.5) was kindly supplied by Schering (Milan, Italy)
37 and TB hydrochloride (molecular weight 327.9) by Novartis (Basel, Switzerland). α - and β -
38 CD were from Wacker Chemie GmbH (Munich, Germany), having a water content of 9.8 %
39 and 11.2 %, respectively, as determined by thermal analysis. γ -CD was from Roquette
40 (Lestrem, France, with 11.0% of water. M β -CD and HP β -CD were from Sigma (Missouri,
41 USA). M β -CD and HP β -CD had average substitution degrees DS \approx 12 and 4 and a water
42 content of 5.2 and 6.5 %, respectively. D $_2$ O (deuterium content 99.9 %) was obtained from
43 Aldrich Chemical (Barcelona, Spain).
44
45
46
47
48
49
50
51
52
53
54
55

56 2.2. Spectrophotometric method

57 The interactions of NF with α -, β -, M β -, HP β - and γ -CD in aqueous solutions (pH 4) were
58 studied at different temperatures: 290, 293, 298, 303 and 310 K. The temperature was
59 controlled by recirculating water from a thermostat bath Lauda Ecoline RE 104. The spectra
60

of NF and NF+CD were recorded with a diode array spectrophotometer HP 8452. The NF concentration was kept constant at 5.0×10^{-5} M while that of CD varied between 4.8×10^{-4} M and 3.8×10^{-3} M by addition of a concentrated solution of NF+CD to the cell. This procedure permits to minimise the errors due to the manipulation of the cuvette or to slight changes in its position in relation to the source. The measurements were made at 254 nm. The Benesi-Hildebrand equation was used to calculate the stability constant (K_{11}) (10).

$$\frac{1}{\Delta A} = \frac{1}{S_t K_{11} \Delta \varepsilon [L]} + \frac{1}{S_t \Delta \varepsilon}$$

where ΔA is the difference between the absorbance of the drug in the presence and absence of CD, $\Delta \varepsilon$ is the difference between the molar absorption coefficient of NF and the complex, S_t and $[L]$ are the initial concentrations of NF and CD, respectively. K_{11} can be calculated from the slope of the straight line obtained by plotting $1/\Delta A$ against $1/[L]$.

The thermodynamic parameters of the inclusion process were determined from the temperature dependence of the association constant using the van't Hoff equation.

2.3. $^1\text{H-NMR}$ measurements

All the NMR measurements were performed at 25°C in D_2O as the solvent. For collecting the 1D spectra we used a Bruker Avance 400 Ultrashield spectrometer, by averaging 600 scans. The solvent signal (HDO, at 4.792 ppm) was used as reference. For NF, the concentration was kept constant at 2.7 mM and that of β -CD varied from 0.5 mM to 10 mM by adding NF to vials with different amounts of β -CD. In the case of TB, the Job plot method was used to ascertain the stoichiometry of the complexes with both β - and γ -CD (10), by mixing a 4 mM solution of the drug with either 4 mM β - or γ -CD in different amounts.

For the 2D ROESY experiments a Bruker Avance AV-500 spectrometer (11.7 T) was used by applying the pulse sequence defined in the literature (11). Different spin-lock mixing times were tested until getting the best signal-to-noise ratio with 600 ms. The FT of the FIDs (1024×1024 matrixes, obtained by averaging at least 64 scans), 2D phase tuning and signal integration were performed with MestRe-C software (version 4.7, Mestrelab Research, 2004). For determining the stability constants of the complexes with TB and β -CD or γ -CD we used a multivariable non-linear analysis with selected protons of the host and guest. The main advantage of the multivariable analysis arises from the increase in the number of fitted data. The protons chosen for the analysis were those that underwent more pronounced shifts, assuming that the stability constant is the same for each proton. The equation used for the

determination of the stability constants is based on the fast exchange between the host and guest in the NMR time scale. If such is the case, the measured chemical shifts are the average of the chemical shifts of the species present (10,12). $\Delta\delta$ (the difference between the chemical shift of the host or guest alone and the measured at each concentration) can be related to the concentration of TB and CD at each point:

$$\Delta\delta = \Delta\delta_0 \frac{1}{2} \left\{ \left(1 + \frac{w}{x} + \frac{1}{K_{11}x} \right) - \sqrt{\left(1 + \frac{w}{x} + \frac{1}{K_{11}x} \right)^2 - 4 \frac{w}{x}} \right\}$$

where $\Delta\delta_0$ is the difference in the limit when all the drug is complexed, w is the TB or CD concentration and x is that of CD or TB, depending if the stability constant is obtained from the host or the guest. All the calculations were performed with ORIGIN software (version 5.0, Microcal. Software, Inc. 1997).

2.4. Electrospray Ionization Mass spectrometry

The ESI-MS spectra of NF, TB and their complexes with cyclodextrins (α -, β -, $M\beta$ -, $HP\beta$ - and γ -CD) were obtained using a Waters micromass ZQ mass spectrometer equipped with an electrospray interface. 1:1 mixtures CD:TB and CD:NF (0.1 mg/mL) were prepared in methanol/water 50/50 (v/v) and were introduced into the ion source of the mass spectrometer with a flow-rate of 20 μ L/min using a syringe pump. The ionization voltage was 3.51 and 3.33 kV for NF and TB, respectively. The temperature of the inlet capillary was 200°C and the detection was performed in the positive ion mode.

3. Results and discussion

3.1. Spectrophotometric characterization of NF-CDs complexes

The addition of increasing amounts of every CD, except for α -CD, caused a decrease in the absorbance intensity at 254 nm and also the appearance of an isosbestic point; both facts indicate complex formation (10) (Figure 3).

These changes in the spectral properties are probably due to the influence of the high electron density inside the CD cavity on the electrons of the guest (13). Plots of $1/\Delta A$ as a function of $1/[CD]$ for NF confirm the formation of a 1:1 inclusion complex. As an example, Figure 4 shows the Benesi–Hildebrand plot of NF in the presence of β -CD at 25°C. The stability constants determined for the complexes at the different temperatures are compiled in Table 1.

1
2
3 The K_{11} values decrease in the order β -CD > M β -CD > HP β -CD > γ -CD. In the case of α -
4 CD, the absence of changes in the UV spectra suggests its cavity is too narrow to include NF.
5 With respect to β -CD and its derivatives, the complexation between NF and the natural
6 cyclodextrin is stronger, probably due to interactions between the amino group and the
7 hydroxyls of the β -CD rims, which are not present in the corresponding derivatives of β -CD.
8 In addition, taking into account that the drug is charged in aqueous solution ($pK_a = 8.0 \pm 0.2$),
9 the substitution of some of the hydroxyls of the CD by the hydroxypropyl and methyl groups
10 probably reduces the affinity of the charged drug for the CD. Although HP β -CD bears more
11 OH groups than M β -CD, the lower constant obtained for HP β -CD in relation to M β -CD could
12 be explained by steric hindrance of the hydroxypropyl group, which could hamper the guest
13 molecule entering the CD cavity (14).
14

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The corresponding enthalpy (ΔH) and entropy (ΔS) values were calculated from the van't Hoff relationship from the temperature dependence of the equilibrium constants. The results are shown in Table 1. The enthalpies are always negative and close to each other, meaning that the attractive forces involved in the complexation of NF with the different CDs, such as Van der Waals and hydrogen bonding between the drug and the CD cavity, are similar (15). In relation to the entropy, it is positive for the complex with β -CD whereas it is negative with the other macrocycles; this can be interpreted in terms of the specific interactions between the CD and the guest. The OH groups of β -CD are initially solvated by structured water, but the formation of hydrogen bonds between the amino group of NF and the hydroxyl groups of the CD would break this structure and release the water molecules in the bulk. However, for the β -CD derivatives, the negative values of entropy are related to an increase of order in the system upon complexation. This could be associated with the solvation of the amino group by water molecules in the complex and also to a pronounced restriction of the rotational and vibrational degrees of freedom in the complex due to the presence of the hydroxyl and methyl groups.

The interactions of neutral NF with CDs have been studied by the solubility method at pH 12 and 298 K in a previous work (16), resulting $K_{11} = 512, 829, 1123$ and 184 M^{-1} for the complexation with β -CD, HP β -CD, M β -CD and γ -CD, respectively. The stability constants resulted higher at pH 12, except for β -CD, showing that the interaction is stronger for the neutral form of NF, according with the explanation of Szejtli (13). Since the drug is uncharged at pH 12, it can penetrate deeper in the CD cavity.

1
2
3 It was intended to study the interactions between TB and CDs but no changes in the TB
4 spectra were observed.
5
6
7

8 9 3.2. ¹H-NMR Spectroscopy

10 As a preliminary step, we have investigated the possible concentration dependence of the
11 chemical shifts of both NF and TB (from 0.5 to 3.5 mM). This is important if aggregation
12 processes in solution are expected, but this has not been the case. The signal assignment of
13 NF and TB is shown in Figures 5 and 6, respectively. These antifungal drugs have similar
14 chemical structures and in both cases the signal of proton d overlaps that of HDO in the ¹H-
15 NMR spectra.
16
17
18
19
20
21

22 2.2.1. NF-βCD interactions

23 The ¹H NMR spectrum for NF is quite similar to that of TB (Figures 5 and 6). The difference
24 is the presence of the benzene ring in the case of NF (signals between 7.4 and 7.3 ppm)
25 instead of the tert-butyl group of TB. Upon addition of a solution of NF to variable amounts
26 of β-CD, the protons that shift are a, g, h, i of NF, the inner protons of β-CD (H3 and H5) and
27 H6. All of them move upfield, with the maximum changes corresponding to those of NF g, h
28 and i, which shift -0.592, -0.963 and -0.515 ppm, respectively, while in the case of CD H5
29 moves -0.065 ppm. Figure 7 displays the spectra for all the measured molar ratios in the
30 regions where the changes are more remarkable.
31
32
33
34
35
36
37

38 The changes in the chemical shifts with respect to those of the pure substances in water, Δδ,
39 have been plotted versus the molar ratio in Figure 8. These plots indicate a 1:1 stoichiometry,
40 as deduced from the extrapolations from high and low molar ratios at $R = 1$.
41
42

43 Provided a fast exchange in the scale of NMR, and a 1:1 stoichiometry, the observed chemical
44 shift δ is the sum of the chemical shift due to the complex ($\delta_{NF:CD}$) and that due to the
45 host/guest (δ_i , being $i = CD$ or NF), each one averaged with its molar fraction (X_i , $X_{NF:CD}$),
46 that is (17):
47
48
49
50

$$51 \delta = \delta_i X_i + \delta_{NF:CD} X_{NF:CD}$$

52 The binding constant for each proton can be estimated by a nonlinear fit of the increments of
53 the chemical shifts with respect to that of the pure host or guest pure compound, Δδ, versus
54 the molar ratio (Figure 8). The values deduced for the stability constants are compiled in
55 Table 2. The average K_{11} obtained with the protons of NF and β-CD are $(8.3 \pm 0.8) \times 10^2 \text{ M}^{-1}$
56 and $(6.5 \pm 1.3) \times 10^2 \text{ M}^{-1}$, respectively. These values are within the experimental uncertainty
57
58
59
60

and agree with the stability constant calculated by UV spectrophotometry at 25°C, despite the different range of concentrations used.

2.2.2. TB- β -CD and TB- γ -CD interactions

The presence of β - or γ -CD induces important shifts in the signals of TB, as well as changes in those of the CDs, especially in the inner H3 and H5. These protons move upfield in the presence of TB (Figure 9). Those that shift to a larger extent are H3 and H5 (-0.070 and -0.092 ppm, respectively). It can also be observed that H1, H2 and H4 change -0.018, +0.043 and -0.002 ppm, respectively. The displacement of H2 may suggest a possible contact of TB with the external surface of the cyclodextrin.¹⁶ The shifts of the TB protons are, in general, greater than those of the CD (Figure 9). The most remarkable changes occur in g, h (vinylic) and e, in an extent of -0.413, -0.263 and +0.153 ppm, respectively. The methyl protons (i) and those of the naphthalene group (a) shift less (-0.016 ppm and -0.099 ppm, respectively).

In the case of γ -CD, the protons of the macrocycle that exhibit the most remarkable changes are H3, -0.065 ppm, and H5, -0.069 ppm. Unlike β -CD, H6 moves less (-0.019 ppm) and the external protons do not shift. The protons of TB that undergo larger shifts are those of the vinyl group and of the tert-butyl group, which move upfield (g, -0.49 ppm, h, -0.353, i, -0.086). On the other hand, e proton moves downfield (+0.158 ppm) as it occurred with β -CD. The interaction with γ -CD is weaker than that with β -CD, as deduced from the lower shifts with the former.

With both cyclodextrins it seems that the interaction takes place between the internal protons of the CD and the aliphatic part of TB, although the inclusion of the naphthalene nucleus can not be discarded. Its size fits both CDs (19), and the changes in some of their protons are significant enough to conclude that inclusion has taken place. This could suggest as most probable a 2:1 stoichiometry. In order to elucidate if this is the case, an NMR titration according to the Job plot method (10) has been performed. The results for TB- β CD are shown in Figure 10, where r_{TB} and r_{CD} are the mole fractions of host and guest, respectively, and $\Delta\delta$ is the change in the chemical shift of the CD and TB protons. In both cases, the maxima appear at a mole fraction of ca. 0.5, what unquestionably indicates a 1:1 stoichiometry for the complex.

The stability constants have been also deduced for both complexes. For TB- β CD, protons g, h, H3, H5 and H6 yield a binding constant of $(1.4 \pm 0.3) \times 10^3 \text{ L mol}^{-1}$. In the case of TB-

1
2
3
4 γ CD, the protons chosen were i, H3 and H5 (Figure 11). The multivariable analysis produces
5 a $K_{11} = (2.2 \pm 1.1) \times 10^2 \text{ L mol}^{-1}$. This constant is lower than that of the complex with β -CD
6 and reflects the fact that the cavity size of γ -CD is wider for TB whereas that of β -CD
7 provides a tighter fitting.
8
9

10 11 12 **3.3. 2D Structure of the complexes**

13
14
15
16 2D ROESY experiments provide valuable information about the actual topology of the
17 inclusion complexes. Figure 12 shows an expanded region of the ROESY spectrum of the
18 system NF- β CD. Although the interpretation is complicated due to signal overlapping, the
19 spectrum reveals intense correlations between signals of the aromatic protons of the guest and
20 those of the inner part of the β -CD (H3 and H5), what confirms the intracavity binding. The
21 most intense are the NOEs between i,j and k protons (phenyl group) and H3 and H5, although
22 those of the naphthalene nucleus correlate also strongly with H3 and scarcely with H5,
23 indicating that the naphthalene enters the cavity by its wider side. In view of the 1:1
24 stoichiometry evidenced by the ^1H NMR data, the only explanation is that there must be a
25 mixture of 1:1 complexes, in which either the benzene part or the naphthalene moiety of NF is
26 included. There would be no possibility to share the molecule between two CDs, at least
27 within the interval of concentration used.
28
29

30
31
32
33
34
35
36
37 In the case of TB and β -CD the spectrum shows (Figure 13) a clear correlation between
38 naphthalene group and the inner protons H3 and H5 of the β -CD. Also, tert-butyl signal of the
39 TB (i protons) correlate strongly with H5 (figure not shown), whereas H2 and H4 give weaker
40 crosspeaks with both naphthalene and tert-butyl group. The fact that the most intense NOE
41 signals are paired as naphthalene-H3 and H5 and i-H5 suggests, as in the case of NF, that the
42 naphthalene group enters the cavity by its wider rim and also the tert-butyl group (this one
43 more deeply buried than the naphthalene). The 1:1 stoichiometry, confirmed by the Job plot,
44 must imply that both topologies must be present in solution.
45
46
47
48
49

50
51
52
53
54
55
56
57
58
59
60
In the case of TB with the wider γ -CD, the qualitative interpretation is the same than that of
the precedent cases: intense cross-peaks appear between the inner protons of γ -CD and i ones
(relative NOE intensities=100), being weaker those between the protons of naphthalene group
and H3 and H5 (NOE volume = 14). This suggests that the affinity of TB for this macrocycle
is mainly due to the inclusion of the aliphatic part of the guest.

3.4. Electrospray Ionization Mass Spectrometry

Full-scan positive-ion mass spectra of TB and NF showed the protonated molecular species $[M+H]^+$ at m/z 292.2 and 288.2, respectively. The most abundant ion in the product ion spectra was at m/z 140.7 for TB and 116.9 for NF, as the result of the dissociation of the drugs.

Ions due to the inclusion of Na^+ by α -, β -, $M\beta$ -, $HP\beta$ - and γ -CD are observed at m/z 995, 1157, 1353, 1563 and 1319, respectively. All the CD-NF and CD-TB complexes, except those with $HP\beta$ -CD, can be detected by ESI-MS in the positive ion mode. As an example, the spectrum obtained for NF- β CD is shown in Figure 14. The complexes of α -, β -, $M\beta$ -, $HP\beta$ - and γ -CD with NF give ions at m/z 1260, 1422, 1618 and 1584 corresponding to the respective drug:CD complexes, which demonstrate the presence of 1:1 stoichiometry in all the cases. Complexes of the same cyclodextrins with TB give ions at m/z 1264, 1426, 1622 and 1588, respectively. Therefore, ESI-MS can provide a direct proof of the formation of 1:1 non covalent complexes in the gas phase. Thus, the stoichiometries determined by ESI are in agreement with those obtained by 1H -NMR and UV spectroscopy.

The signals for TB- α CD and NF- α CD are weaker than those of α -CD alone, probably due to a weak host-guest interaction. As expected, the interactions of the antifungal drugs with β -CD are stronger, in agreement with the fact that the cavity of α -CD is too narrow for lodging aromatic molecules, while β -CD fits adequately (19). The interaction of NF with γ -CD is weak but it seems to be stronger for TB, in accordance with a better fit in this wider macrocycle.

Acknowledgments

Authors are grateful to the Gobierno de Navarra for M. Uzqueda's grant and to Andrés Guerrero-Martínez (Universidad Complutense de Madrid) for the NMR and ROESY measurements.

References

- (1) Martín del Valle, E. M. Cyclodextrins and their uses: a review. *Process Biochem.* **2004**, *39*, 1033-1046.
- (2) Connors, K. A. The stability of cyclodextrin complexes in solution. *Chem. Rev.* **1997**, *97*, 1325-1358.
- (3) Hirayama, F.; Uekama, K. Cyclodextrin-based controlled drug release system. *Adv. Drug Deliver. Rev.* **1999**, *36*, 125-141.
- (4) Jicsinszky, L.; Fenyvesi, E.; Hashimoto, H.; Ueno, A. In *Comprehensive Supramolecular Chemistry. Cyclodextrins*; Szejtli, J.; Osa, T., Eds.; Elsevier: UK, 1996; Vol. 3; Chapter 4.
- (5) Uekama, K.; Irie, T. In *Comprehensive Supramolecular Chemistry. Cyclodextrins*; Szejtli, J.; Osa, T., Eds.; Elsevier: UK, 1996; Vol. 3; Chapter 15
- (6) Schuster, I.; Ryder, N. S. Mode and selectivity of action compared to azole antifungals and biological fate in mammalian organisms. *J. Dermatol. Treat.* **1990**, *1*, 7-9.
- (7) Balfour, J. A.; Faulds, D. A review of its Pharmacodynamic and Pharmacokinetic properties, and Therapeutic potential in Superficial Mycoses. *Drugs* **1992**, *43*, 259-284.
- (8) Schneider, H-J; Hacket, F.; Rüdiger, V.; Ikeda, H. NMR Studies of Cyclodextrin and Cyclodextrin Complexes. *Chem. Rev.* **1998**, *98*, 1755-1785.
- (9) González-Gaitano, G.; Rodríguez Sainz-Rozas, P.; Isasi, J. R.; Guerrero-Martínez, A.; Tardajos, G. Site-Specific Interaction between 2-Dibenzofuran Carboxilate and β - and γ -Cyclodextrins Determined by Intermolecular NOE and Molecular Modelling. *J. Phys. Chem. B.* **2004**, *108*, 14154-14162.
- (10) Connors, K.A. *Binding constants. The Measurement of Molecular Complex Stability*. Wiley: USA, 1987.
- (11) Bax, A.; Davis, D. G. Practical aspects of two-dimensional transverse NOE spectroscopy. *J. Magn. Reson.* **1985**, *63*, 207-213.
- (12) González-Gaitano, G.; Tardajos, G. Chemical Equilibrium in Supramolecular Systems as Studied by NMR Spectrometry. *J. Chem. Edu.* **2004**, *81*, 270-274.
- (13) Szejtli, J. In *Comprehensive Supramolecular Chemistry. Cyclodextrins*; Szejtli, J.; Osa, T., Eds.; Elsevier: UK, 1996; Vol. 3; Chapter 5.
- (14) Liu, L.; Zhu, S. Preparation and characterization of inclusion complexes of prazosin hydrochloride with β -cyclodextrin and hydroxypropyl- β -cyclodextrin. *J. Pharm. Biomed. Anal.* **2005**, *40*, 122-127.
- (15) Liu, L.; Guo, Q. X. The driving forces in the inclusion complexation of cyclodextrins. *J. Incl. Phenom.* **2002**, *42*, 1-14.

- 1
2
3 (16) Uzqueda, M.; Martín, C.; Zornoza, A.; Sánchez, M.; Martínez-Ohárriz, M. C.; Vélaz, I.
4 Characterization of complexes between naftifine and cyclodextrins in solution and in the
5 solid state. *Pharm. Res.* **2006**, *23*, 980-988.
6
7
8 (17) Loukas, Y. L.; Vraka, V.; Gregoriadis, G. Novel non-acidic formulations of haloperidol
9 complexed with beta-cyclodextrin derivatives. *J. Pharm. Biomed. Anal.* **1997**, *16*, 263-
10 268.
11
12
13 (18) Bratu, I.; Gavira-Vallejo, J. M.; Hernanz, A. ¹H-NMR Study of the Inclusion Processes
14 for α - and γ -cyclodextrin with Fenbufen. *Biopolymers* **2005**, *77*, 361-367.
15
16
17 (19) Szente, L. In *Comprehensive Supramolecular Chemistry. Cyclodextrins*; Szejtli, J.; Osa,
18 T., Eds.; Elsevier: UK, 1996; Vol. 3; Chapter 7.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Tables

Table 1. Stability constants^a ($K_{11} \cdot 10^{-2} / M^{-1}$) and thermodynamic parameters of NF complexes with CDs in water.

	290 K	293 K	298 K	303 K	310 K	ΔH° (kJ mol ⁻¹)	ΔS° (JK ⁻¹ mol ⁻¹)
β -CD	10.8 ± 0.4	10.2 ± 0.3	9.0 ± 0.3	8.2 ± 0.3	6.9 ± 0.6	-17.0 ± 0.8	0.6 ± 3
M β -CD	4.1 ± 0.1	3.8 ± 0.1	3.3 ± 0.4	2.9 ± 0.4	2.5 ± 0.4	-19.2 ± 0.5	-16.3 ± 1.7
HP β -CD	3.0 ± 0.2	2.8 ± 0.1	2.5 ± 0.4	2.2 ± 0.1	1.9 ± 0.1	-18.7 ± 0.5	-16.9 ± 0.2
γ -CD	2.3 ± 0.1	----	2.1 ± 0.2	1.7 ± 0.2	----	-14.0 ± 0.1	-3.0 ± 0.4

^astandard errors are indicated

Table 2. Binding constants^a ($K_{11} \times 10^{-2} / M^{-1}$) corresponding to the NF- β CD system, calculated with NF and β -CD protons.

NF proton	K_{11}	β -CD proton	K_{11}
a	5.1 ± 1.1	H3	7.0 ± 1.0
g	8.9 ± 0.6	H5	5.5 ± 1.2
h	9.6 ± 0.8	H6	7.1 ± 1.6
i	9.7 ± 0.7		

^astandard errors are indicated

Figure captions

Figure 1. β -cyclodextrin molecule

Figure 2. Chemical structures of NF (a) and TB (b) hydrochlorides

Figure 3. Absorption spectra of 5.0×10^{-5} M NF as a function of $[\beta\text{-CD}]$ at 25°C.

Figure 4. Benesi–Hildebrand plot of $1/\Delta A$ vs. $1/[\beta\text{-CD}]$ for NF at 25°C.

Figure 5. ^1H -NMR spectrum of 2.7 mM NF in D_2O .

Figure 6. ^1H -NMR spectrum of 0.5 mM TB in D_2O .

Figure 7. Selected regions of ^1H NMR spectra in D_2O at different $[\text{NF}]/[\beta\text{-CD}]$ ratios.

Figure 8. Increments in the chemical shifts versus the molar ratio for selected protons of (a) NF and (b) $\beta\text{-CD}$. Solid lines are obtained by nonlinear fit assuming a 1:1 stoichiometry.

Figure 9. Selected regions of the ^1H NMR spectra of (a) TB 4 mM; (b) $\beta\text{-CD}$ 4 mM; (c) and (d) $[\text{TB}]/[\beta\text{-CD}] = 0.5$; (e) and (f) $[\text{TB}]/[\beta\text{-CD}] = 0.3$.



Figure 10. Job plots for selected protons of (a) TB and (b) $\beta\text{-CD}$.

Figure 11. Job plots for selected protons of (a) TB and (b) $\gamma\text{-CD}$.



Figure 12. Zoomed view of the ROESY spectrum of the system NF- βCD .

Figure 13. Partial view of the ROESY spectrum for the TB- βCD system (aromatic region).


Figure 14. ESI-MS spectrum of $\beta\text{-CD}$ with NF.




Peer Review Only




For Peer Review Only




For Peer Review Only



For Peer Review Only





1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60