

2023

The impact of smoking on response to tumor necrosis factor- α inhibitor treatment in patients with ankylosing spondylitis

HANDAN YARKAN TUĞSAL

GÖKÇE KENAR ARTIN

GERÇEK CAN

SEDAT ÇAPAR

BERRİN ZENGİN

See next page for additional authors

Follow this and additional works at: <https://journals.tubitak.gov.tr/medical>



Part of the [Medical Sciences Commons](#)

Recommended Citation

TUĞSAL, HANDAN YARKAN; ARTIN, GÖKÇE KENAR; CAN, GERÇEK; ÇAPAR, SEDAT; ZENGİN, BERRİN; AKAR, SERVET; DALKILIÇ, HÜSEYİN EDİZ; ŞENEL, ABDURRAHMAN SONER; KOCA, SÜLEYMAN SERDAR; GÖKER, BERNA; YAZICI, AYTEN; İNANÇ, GÜZİDE NEVSUN; ELLİDOKUZ, HÜLYA; AKKOÇ, NURULLAH; and ÖNEN, FATOŞ (2023) "The impact of smoking on response to tumor necrosis factor- α inhibitor treatment in patients with ankylosing spondylitis," *Turkish Journal of Medical Sciences*: Vol. 53: No. 4, Article 15. <https://doi.org/10.55730/1300-0144.5661>

Available at: <https://journals.tubitak.gov.tr/medical/vol53/iss4/15>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Medical Sciences by an authorized editor of TÜBİTAK Academic Journals. For more information, please contact academic.publications@tubitak.gov.tr.

The impact of smoking on response to tumor necrosis factor- α inhibitor treatment in patients with ankylosing spondylitis

Authors

HANDAN YARKAN TUĞSAL, GÖKÇE KENAR ARTIN, GERÇEK CAN, SEDAT ÇAPAR, BERRİN ZENGİN, SERVET AKAR, HÜSEYİN EDİZ DALKILIÇ, ABDURRAHMAN SONER ŞENEL, SÜLEYMAN SERDAR KOCA, BERNA GÖKER, AYTEN YAZICI, GÜZİDE NEVSUN İNANÇ, HÜLYA ELLİDOKUZ, NURULLAH AKKOÇ, and FATOŞ ÖNEN

The impact of smoking on response to tumor necrosis factor- α inhibitor treatment in patients with ankylosing spondylitis

Handan YARKAN TUĞSAL^{1,*}, Gökçe KENAR¹, Gerçek CAN¹, Sedat ÇAPAR², Berrin ZENGİN¹, Servet AKAR³, Ediz DALKILIÇ⁴, Soner ŞENEL⁵, Süleyman Serdar KOCA⁶, Berna GÖKER⁷, Ayten YAZICI⁸, Nevsun İNANÇ⁹, Hülya ELLİDOKUZ¹⁰, Nurullah AKKOÇ¹¹, Fatoş ÖNEN¹

¹Department of Internal Medicine, Division of Rheumatology, Faculty of Medicine, Dokuz Eylül University, İzmir, Türkiye

²Department of Statistics, Faculty of Medicine, Dokuz Eylül University, İzmir, Türkiye

³Division of Rheumatology, Department of Internal Medicine, Faculty of Medicine, Katip Çelebi University, İzmir, Türkiye

⁴Division of Rheumatology, Department of Internal Medicine, Faculty of Medicine, Uludağ University, Bursa, Türkiye

⁵Division of Rheumatology, Department of Internal Medicine, Faculty of Medicine, Erciyes University, Kayseri, Türkiye

⁶Division of Rheumatology, Department of Internal Medicine, Faculty of Medicine, Fırat University, Elazığ, Türkiye

⁷Division of Rheumatology, Department of Internal Medicine, Faculty of Medicine, Gazi University, Ankara, Türkiye

⁸Division of Rheumatology, Department of Internal Medicine, Faculty of Medicine, Kocaeli University, Kocaeli, Türkiye

⁹Division of Rheumatology, Department of Internal Medicine, Faculty of Medicine, Marmara University, İstanbul, Türkiye

¹⁰Department of Biostatistics and Medical Informatics, Faculty of Medicine, Dokuz Eylül University, İzmir, Türkiye

¹¹Division of Rheumatology, Department of Internal Medicine, Faculty of Medicine, Celal Bayar University, Manisa, Türkiye

Received: 12.09.2022 • Accepted/Published Online: 01.02.2023 • Final Version: 18.08.2023

Background/aim: To investigate the impact of smoking on disease activity, treatment retention, and response in patients with ankylosing spondylitis (AS) treated with their first tumor necrosis factor- α inhibitor (TNFi).

Materials and methods: AS patients who started their first TNFi treatment for the active axial disease (BASDAI \geq 4) from TURKBIO Registry were included. Treatment response of smoker (current and ex-smokers) and nonsmoker (never smoker) patients were primarily evaluated as achievement of BASDAI50 or improvement in BASDAI at least 20 mm at 3 months and 6 months compared to baseline.

Results: There were 322 patients with AS (60% male, 59% smoker, mean age: 38.3 years). The median follow-up time was 2.8 years (Q1–Q3: 1.3–3.8), and disease duration was 3.5 years (Q1–Q3: 0.7–8.2). Smokers had male predominance ($p < 0.001$), lower ESR ($p = 0.03$), higher BASDAI ($p = 0.02$), BASFI ($p = 0.05$), HAQ-AS ($p = 0.007$), and ASDAS-CRP ($p = 0.04$) compared with nonsmokers at baseline. In the multivariate analysis, male gender [OR 2.7 (95%CI 1.4–5), $p = 0.002$], and concomitant conventional synthetic disease-modifying antirheumatic drug use [OR 2.4 (95%CI 1.1–5.2), $p = 0.03$] were associated with better treatment response. There was an association of male gender [HR 2.4 (95%CI 1.6–3.7), $p < 0.001$], older age (≥ 30 years) [HR 1.8 (95%CI 1.1–2.8), $p = 0.01$], and response to treatment [HR 1.8 (95%CI 1.2–2.9), $p = 0.008$] with better treatment retention. No impact of smoking status was found on treatment retention and response in univariate and multivariate analyses.

Conclusion: This study suggested that smoking was associated with poorer patient-reported outcomes in biologic naïve AS patients initiating their first TNFi treatment, but it had no impact on the TNFi treatment response and retention rate.

Key words: Ankylosing spondylitis, smoking, tumor necrosis factor-alpha inhibitor, treatment response, registry

1. Introduction

Cigarette smoking, one of the most severe public health problems, is a modifiable risk factor for several diseases. It is a major cardiovascular risk factor and estimated that 8 million deaths per year worldwide¹. In the literature, a relationship between rheumatoid arthritis (RA) and smoking was first shown among inflammatory diseases,

and smoking was reported to be the strongest known environmental risk factor for the development of RA [1]. Then, interest has been directed towards other rheumatic diseases to see whether smoking affects development, progression and treatment response in other rheumatic diseases. In patients with systemic lupus erythematosus, negative effects of smoking on the course of the disease and

¹World Health Organization (2019). Tobacco fact sheet [online]. Website <https://www.who.int/news-room/fact-sheets/detail/tobacco> [accessed 24 May 2022].

* Correspondence: drhandanyarkan@gmail.com

its treatment have been reported [2]. (The role of smoking in rheumatic diseases other than autoimmune spectrum (less female predominance and autoantibodies) is less clear. In terms of spondyloarthritis, a large population-based cohort study has shown that current smoking is significantly associated with incident ankylosing spondylitis (AS) [3]. Data from previous cross-sectional and longitudinal studies showed that smoking was an independent risk factor for structural damage in patients with axial spondyloarthritis (axSpA) [4–7]. However, its impact on global disease activity, functional status, and health-related quality of life in axSpA patients is unclear based on the limited data with contradictory results [8].

The tumor necrosis factor- α inhibitor (TNFi) drugs (adalimumab, certolizumab pegol, etanercept, golimumab, and infliximab) have revolutionized the treatment of AS over the past 20 years. Nevertheless, there are still a considerable number of patients who are resistant to treatments with TNFi. The response rates of the *Assessment of SpondyloArthritis International Society* (ASAS)20, ASAS40, ASAS partial remission were 58%–64%, 40%–47%, and 20%–23%, respectively in AS patients treated with the five TNFi drugs in the phase III trials [9–10]. Furthermore, real-life data showed that as many as 45% of patients with AS discontinued TNFi therapy within the first two years [11]. Older age, negative HLA-B27, and higher baseline disease activity were defined as the independent predictors of the nonresponse to first TNFi [12–14]. Furthermore, conflicting results had been reported in previous studies on the effect of smoking on the TNFi response [15].

The primary aim of this study was to investigate the impact of smoking on baseline disease activity, treatment response (BASDAI50/20) [9] and, treatment retention in Turkish patients with AS treated with their first TNFi therapy. We also aimed to determine the factors that could be associated with treatment efficacy and treatment retention, and the effect of smoking on treatment response evaluated by ASDAS [16].

2. Material and method

2.1 Patients

In this observational cohort study, we included adult AS patients who started their first TNFi treatment for active axial disease (BASDAI \geq 4) and who had at least two visits after the initiation of therapy by the time of 1 January 2018 in the TURKBIO registry.

We excluded patients with baseline BASDAI < 4 or who initiated TNFi therapy for an indication other than axial disease, patients with <2 follow-up visits, patients who discontinued follow-up with no recorded reason, and patients with unknown smoking status.

TURKBIO registry is a nationwide biological database contributed by 14 different centers across Turkey. It includes

adult (\geq 18 years) patients with RA, AS, nonradiographic axial spondyloarthritis (nr-AxSpA), and psoriatic arthritis. Demographic and clinical features, follow-up parameters related to disease and treatment, current and previous treatments, adverse events and discontinuation rates, and reasons were registered electronically using open-source software at each of the 3–6 months visits. The Drug Regulatory Authority of the Health Ministry of Turkey and the Dokuz Eylül University Ethics Committee has approved the Registry Project as a phase IV observational study. The whole study was performed according to the Declaration of Helsinki and the ethical approval for the secondary use of the data included in the TURKBIO has been obtained previously (Dokuz Eylül University Ethics Committee; 13.04.2017; protocol no: 304-SBKAEK). Patients signed a written informed consent form before their inclusion in the study. We followed the STROBE checklist of items for reports of cohort studies.

2.2 Smoking status

Smoking was defined as using >1 cigarette/day without reference to the quantity (e.g., pack-years). Ex-smoker was defined as one who had smoked more than 100 cigarettes in the lifetime but has not smoked in the last 28 days. Patients' smoking status at the start of TNFi therapy was obtained from the data of the closest yearly visit in the TURKBIO registry. They were stratified into two groups: 1. Smokers (including current- and ex-smokers) and 2. Nonsmokers (including never-smokers).

2.3 Disease activity, function, mobility and health-related quality of life

Disease activity was evaluated by the Bath ankylosing spondylitis disease activity index (BASDAI) [17] and, the ankylosing spondylitis disease activity score calculated using C-reactive protein (ASDAS-CRP) [18]. Functional status and mobility were assessed using the Bath ankylosing spondylitis functional index (BASFI) [19] and the Bath ankylosing spondylitis metrology index (BASMI) [20], respectively. Patients completed the health assessment questionnaire for ankylosing spondylitis (HAQ-AS) for health-related quality of life assessment [21].

2.4 Treatment response

Clinical response to TNFi was primarily evaluated by BASDAI50/20 [improvement of at least 50% in the BASDAI score or an absolute change of 20 units (on a 0 to 100 scale) after three months of therapy] as recommended by ASAS [9]. Patients who achieved clinical response at both third- and sixth-months' visits compared to baseline were defined as responders. The patients with a clinical response at either the third- or sixth-months' visits were also classified as 'responders' when there was a reported reason for missing visits. If there were no visits within the given month, the closest visit was chosen not exceeding

eight weeks. For baseline visit, the period between one month before and one week after the TNFi therapy were allowed.

Furthermore, ASDAS clinical responses [16] as ASDAS-LDA (Low Disease Activity, ASDAS 1.3–2.1), ASDAS CII (Clinically Important Improvement, $\Delta \geq 1.1$), and ASDAS MI (Major Improvement, $\Delta \geq 2$) were evaluated at third and sixth months of therapy.

2.5 Treatment retention

Treatment retention was defined as the number of years from the initiation to the end of therapy. Treatment cessation for any reason (e.g., local reactions, infections, surgery) less than two months were allowed.

The reasons for treatment discontinuation were recorded in TURKBIO as lack of effect, adverse events, cancer, follow-up at other centers, pregnancy, infections, discontinuation of follow-up, death, surgery, disease remission, patient preference, and other reasons. For this study, the reasons for discontinuation were collected into three groups; adverse events (including infection, death, and cancer), lack of effect, and others. The patients without a record of the reason for treatment cessation were excluded from the study.

2.6 Statistics

Demographic and descriptive data are presented as median, first and third quartile (Q1–Q3). Baseline characteristics in categorical variables were compared between smokers (ever or current) and nonsmokers using nonparametric tests (χ^2 , Kruskal Wallis and Mann Whitney tests). In all tests, a 5% type-1 error level was used to infer statistical significance.

The univariate analyses to identify variables associated with treatment response were investigated using Pearson's chi-square, Fisher exact, Student's *t*, and Mann-Whitney *U* tests, where appropriate. Age (<30 years old vs. ≥ 30 years old) and disease duration (<4 years vs. ≥ 4 years) were included as categorical variables to allow for possible nonlinear effects. For the multivariate analysis, the possible factors [age, concomitant conventional synthetic disease-modifying antirheumatic drug (csDMARD) use, and gender] identified with univariate analyses were further entered into the logistic regression analysis to determine independent predictors of treatment response. The odds ratios (OR) and 95% confidence intervals (95% CI) were also calculated. In the regression model, both age and gender were retained as covariates to control their influence on other covariates. Since baseline BASDAI were higher in smokers, parameter estimates from the multivariable model were adjusted for baseline difference. Hosmer-Lemeshow goodness of fit statistics was used to assess model fit.

The effect of smoking and other factors on treatment retention was investigated using the log-rank test. The Kaplan-Meier survival estimates were calculated. The

possible factors identified with univariate analyses (age, gender, disease duration, and response to treatment) were further entered into Cox regression analysis, with backward selection, to determine independent predictors of adherence and calculated hazard ratios (HRs). The effects of potential covariates (age and gender) on treatment retention were adjusted by retaining them in the multivariable model. However, adjustment for baseline BASDAI was performed since smokers have higher baseline BASDAI.

Stratified analyses were performed according to gender.

3. Results

3.1 Baseline characteristics

There were 322 patients (60% male, mean age: 38.3 years) who fulfilled the modified New York classification criteria for AS [22] and followed up since the first TNFi therapy in TURKBIO by 2018. Among the 322 patients, 191 (59%) were smokers (125 current- and 66 ex-smokers) and 73% of smokers were male.

At baseline, smoker patients had higher BASDAI, BASFI, HAQ-AS, ASDAS-CRP, and lower erythrocyte sedimentation rate (ESR) than nonsmokers. Median BASMI was slightly higher in smokers than nonsmokers but statistically nonsignificant ($p = 0.34$). There were no differences concerning age, disease duration, HLA-B27 status, serum CRP levels, BMI measurements, medications including csDMARDs, and the reasons for stopping TNFi treatment between smoker and nonsmoker patient groups (Table 1).

3.2 Treatment response

Among 322 AS patients, 267 (83%) responded to the first TNFi therapy based on BASDAI50/20. Smoker and nonsmoker patients had similar treatment responses (BASDAI50/20: 85% and 79%, respectively). There was no statistically significant difference between current vs. never smokers ($p = 0.19$), current vs. ex-smokers ($p = 0.94$), and current vs. never+ ex-smokers ($p = 0.31$) regarding TNFi response. When men and women were analyzed separately, TNFi response was not different between smoker vs. nonsmoker ($p = 0.98$, $p = 0.71$), current vs. never-smoker ($p = 0.90$, $p = 0.47$), current vs. ex-smoker ($p = 0.73$, $p = 0.38$), or current vs. never+ ex-smoker ($p = 0.81$, $p = 0.40$) groups in both men and women respectively.

There was no statistically significant difference between smoker and nonsmoker patients regarding ASDAS-CII ($p = 0.32$) and ASDAS-MI response ($p = 0.65$) and achievement of ASDAS-LDA ($p = 0.31$) at both 3 and 6 months. ASDAS-CII ($p = 0.75$, $p = 0.83$) and ASDAS-MI response ($p = 0.25$, $p = 0.80$), and achievement of ASDAS-LDA ($p = 0.37$, $p = 0.62$) were also not different between smoker and nonsmoker patients in both men and women respectively.

Table 1. Demographics and patient characteristics.

	Smoking status		
	Smoker n = 191	Nonsmoker n = 131	p
Age, median (Q1–Q3), years	38 (30–44)	39 (30–46)	0.397
Disease duration, median (Q1–Q3), years	3.3 (0.6–8.9)	3.7 (0.9–7.3)	0.795
Male, n (%)	139 (72)	56 (42)	<0.001
HLA positive ^a , n (%)	95 (69)	58 (62)	0.228
Body Mass Index ^b , kg/m ² , median (Q1–Q3)	25.1 (22.6–29.4)	27.1 (23.9–29.5)	0.142
CRP, mg/L, median (Q1–Q3)	11.5 (5–25.2)	13 (5–29)	0.772
ESR, mm/h, median (Q1–Q3)	26 (12–42.2)	32 (19–49)	0.029
Baseline disease characteristics, median (Q1–Q3)			
BASDAI	58 (49–66)	54 (46–62)	0.019
BASFI	38 (25–52)	34 (21.25–45.5)	0.052
BASMI	20 (4–50)	15 (4–30)	0.338
HAQ-AS	0.75 (0.5–1)	0.625 (0–0.875)	0.007
ASDAS-CRP	3.7 (2.7–4.2)	3.35 (0–3.975)	0.042
TNFi type, n (%)			0.315
Adalimumab	39 (20)	33 (25)	
Etanercept	57 (30)	31 (24)	
Infliximab	58 (30)	32 (24)	
Golimumab	26 (14)	25 (19)	
Certolizumab	11 (6)	10 (8)	
Current DMARD use	29 (15)	17 (13)	
Methotrexate, n (%)	16 (8)	6 (5)	0.492
Sulphasalazine, n (%)	13 (7)	11 (8)	
Previous DMARD use, n (%)	18 (10)	15 (11)	0.556
Methotrexate	7 (4)	6 (4.5)	
Sulphasalazine	9 (5)	6 (4.5)	
Reason of treatment discontinuation ^c n (%)	67 (35)	37 (28)	0.588
Advers events	8 (4)	6 (5)	
Lack of efficacy	25 (13)	16 (12)	
Other	34 (18)	15 (11)	
Total follow-up time, median (Q1–Q3), years	2.9 (1.5–4.3)	2 (1.06–3.5)	0.003
Treatment response ^d , n (%)	163 (85)	104 (79)	0.163
Changes at 3 rd months ^e , median (Q1–Q3)			
BASDAI	4.4 (3.5–5.6)	4.2 (3–5.3)	0.278
BASFI	2.2 (0.4–3.7)	2.1 (0.7–3.5)	0.809
BASMI	1 (–0.4–4.8)	0.4 (–0.4–3)	0.287
ASDAS-CRP	2.2 (1.4–3.1)	2.1 (0.7–3.3)	0.282
HAQ-AS	0.37 (–0.1–0.7)	0.62 (0.25–0.87)	0.006
Changes at 6 th months ^e , median (Q1–Q3)			
BASDAI	4.4 (3.7–5.5)	4.4 (3.6–5.5)	0.817
BASFI	2.4 (2.7–4)	2.4 (1–3.8)	0.683

Table 1. (Continued)

BASMI	3 (0.4–5.2)	1 (0.4–3)	0.108
ASDAS-CRP	2.5 (0.8–3.3)	2.2 (–0.05–3.3)	0.437
HAQ-AS	0.5 (–0.05–0.75)	0.6 (0.20–0.99)	0.055

Abbreviations: ASDAS-CRP: Ankylosing Spondylitis Disease Activity Score calculated using C-Reactive Protein; BASDAI: Bath Ankylosing Spondylitis Disease Activity Index; BASFI: Bath Ankylosing Spondylitis Functional Index; BASMI: Bath Ankylosing Spondylitis Metrology Index; DMARD: Disease-Modifying AntiRheumatic Drug; HAQ-S: Health Assessment Questionnaire for Ankylosing Spondylitis; TNFi: Tumor necrosis factor- α inhibitor; Q1–Q3: First and third quartile

^aNot all patients had HLA B27 status, HLA B27 was available for 231 patients. Percentages were reported as positivity among known HLA status. ^bNot all patients had BMI, analyses were done with 193 patients. ^cPercentages of patients who have stopped TNFi treatment according to smoking status. ^dTreatment response was defined as BASDAI50/20 mm response both at 3th and 6th months' visits compared to baseline. ^eChanges at 3rd and 6th month mean decreases from baseline

3.3 Change in the follow-up outcomes

Third and 6th-month improvement in HAQ-AS was significantly better in nonsmokers than smokers. However, improvement in the other follow-up parameters (Δ BASDAI, Δ BASFI, Δ BASMI, and Δ ASDAS-CRP) at 3 and 6 months was not different between the groups. When males and females were analyzed separately, nonsmoker males had better improvement in third month BASFI and HAQ-AS and 6th-month HAQ-AS than smoker males. However, the improvement of HAQ-AS at third and 6th-month visits was not different between smoker and nonsmoker females.

The possible factors affecting treatment response, including gender, concomitant csDMARD use, and age (grouped as <30 and \geq 30) in the univariate analysis, were further entered into the logistic regression analysis. Multivariate regression analysis showed that males [OR: 2.7; (95% CI 1.4–5), $p = 0.002$] and concomitant csDMARD users [OR: 2.4; (95% CI 1.1–5.2) $p = 0.03$] had better response rates of BASDAI50/20 (Table 2).

3.4 Treatment retention

The total follow-up time was 908.4 patient years. The median follow-up time was 2.8 years (Q1–Q3: 1.3–3.8), and ever-smokers had followed longer [2.9 years (Q1–Q3: 1.5–4.3)] than never smokers [2 years (Q1–Q3: 1.1–3.5)].

Smokers had similar drug survival [2.36 (1.00–3.72) years] to nonsmokers [1.75 (0.77–3.22) years; Kaplan Meier log rank = 0.06] (Figure). Both female smokers [1.6 (0.5–2.9) years] and male smokers [2.5 (1.3–4.1) years] had similar drug survival to female nonsmokers [1.4 (0.5–2.9) years] and male nonsmokers [2.6 (1–3.6) years] (both Kaplan Meier log rank = 0.7) respectively.

The possible factors affecting drug survival in univariate analysis were gender, age (\geq 30 years old), response to treatment, and disease duration (\geq 4 years). In female patients, age (\geq 30 years old), disease duration (\geq 4 years), and response to TNFi therapy, in males age (\geq 30 years old) were the possible factors affecting drug survival

determined by univariate analysis.

Multivariate analysis showed that treatment retention was better in men [HR: 2.4, (95% CI 1.6–3.7), $p < 0.001$], patients who were older than 30 years of age [HR: 1.8, (95% CI 1.1–2.8), $p = 0.01$] and treatment responders [HR: 1.8, (95% CI 1.2–2.9) $p = 0.008$] (Table 3).

4. Discussion

This study showed that smoking was associated with higher baseline disease activity, decreased function, and impaired health-related quality of life in biologic naïve AS patients initiating their first TNFi in the TURKBIO registry. However, it had no impact on their first TNFi treatment response and drug survival.

Several studies reported higher baseline disease activity, functional disability, and impaired quality of life in smoker AS patients than nonsmokers, similar to our study [7,8,23–28]. Matthey et al. showed the negative and dose-dependent impact of smoking on measures of disease severity in AS patients and was independent of age, gender, deprivation level, and disease duration [25]. Another report from the Scotland Registry for AS suggested that smoking cessation was associated with lower disease activity and better physical function and quality of life in patients with AS [27].

Evidence suggests that smoking is associated with axial skeletal radiographic severity and disease progression over time in ax-SpA patients [5, 7, 29–31]. A study in the OASIS cohort, Ramiro et al., showed that the impact of smoking on radiographic progression was through impairment in disease activity in patients with AS [30]. Another study suggested a strong association between the amount of smoking and spinal radiographic progression evaluated by mSASSS in AS patients [32].

There is limited data with conflicting results regarding the impact of smoking on TNFi response in AS patients [33]. Two studies with adjusted analysis reported its negative impact on the TNFi treatment efficacy. The first

Table 2. Results of multivariate logistic regression analysis for factors affecting treatment response.

Variable	OR (95% CI)	P-value
Gender (male vs. female)	2.7 (1.4–5)	0.002
csDMARD use (yes vs. no)	2.4 (1.1–5.2)	0.033
Age (≥ 30 vs. < 30 years old)	1.4 (0.6–3.1)	0.372

Abbreviations: csDMARD: conventional synthetic Disease-Modifying AntiRheumatic Drug CI: Confidence Interval OR: Odds ratio
Parameter estimates from the multivariable model were adjusted for baseline BASDAI

Figure. Treatment retention according to smoking status results from Kaplan-Meier analysis [median (95% CI)]




Figure. Treatment retention according to smoking status results from Kaplan-Meier analysis [median (95% CI)].

Table 3. Results of multivariate cox regression analysis for factors affecting treatment retention.

Variable	HR (95% CI)	P-value
Gender (male vs. female)	2.4 (1.6–3.7)	< 0.001
Age (≥ 30 vs. < 30 years old)	1.8 (1.1–2.8)	0.012
Disease duration (≥ 4 vs. < 4 years)	1.5 (0.9–2.2)	0.072
Treatment response (yes vs. no)	1.8 (1.2–2.9)	0.008

Abbreviations: CI: Confidence Interval HR: Hazard ratio
Parameter estimates from the multivariable model were adjusted for baseline BASDAI

study revealed that current and ex-smoker AS patients were less likely to achieve BASDAI50/20 response than nonsmokers at 3 and 6 months in the DANBİO cohort, including 1425 patients [34]. Current smokers had 46% lower odds of achieving BASDAI50/20. Ciurea et al. also showed that current smokers had poorer BASDAI50 and ASDAS responses rates than nonsmokers among patients with elevated baseline CRP. However, there was no lower response rate in smokers with normal serum CRP levels [35]. Furthermore, the difference in BASDAI change over time (Δ BASDAI) was not clinically or statistically significant, despite current smokers having 54% lower odds of BASDAI50 at one year.

No statistically significant differences were found in studies investigating change in continuous outcomes of follow-up with TNFi between smoker and nonsmoker ax-SpA patients [33, 36, 37]. Kydd et al. showed that smoking did not affect TNFi response in terms of health-related quality of life among Australian AS patients [37]. Recently Zhao et al. also reported that response to TNFi drugs did not differ according to smoking status in the British AxSpA cohort [33]. In our study, TNFi response investigated by several follow-up parameters, including both continuous or binary outcomes and drug survival were not different between smoker and nonsmoker AS patients. Only HAQ-AS score improvement at 3 and 6 months was found to be better in nonsmokers than smokers.

Contradictory results between the studies may be related to using continuous or binary response outcomes. Since smokers had higher baseline disease activity than nonsmokers, they would be less likely to reach a binary response, although they had similar absolute improvement in disease activity or other follow-up parameters over time [33]. On the other hand, better response rates could be expected in smoker AS patients with higher disease activity since higher baseline disease activity was associated with better clinical response in patients with AS initiating TNFi.

In our study, males, and active csDMARD users had better response rates. Male gender was reported as a baseline predictor of TNFi response in previous studies, consistent with our study [38].

To date, the impact of smoking on TNFi response was investigated primarily in observational database studies with different methodological features and significant limitations due to their retrospective designs. The criteria used for either inclusion or exclusion of patients also may lead to essential differences in the results. In this study,

we excluded patients with baseline BASDAI < 4, patients who initiated TNFi for nonaxial involvements, and those with follow-up visits less than two (without known cause) to provide a homogenous group and evaluate clinical response more accurately. All these are the strengths of our study, although they led to a decrease in the number of patients as a limitation.

The definition of treatment success is another issue. Some studies define response by continuous variables (Etc. BASDAI, ASDAS), others by dichotomized variables as responder or nonresponder. Our study used binary BASDAI50/20 response as the primary outcome and evaluated other outcomes, including ASDAS-LDA, ASDAS-CII, and ASDAS-MI. Furthermore, changes at 3 and 6 months from baseline in all these parameters were investigated to see absolute improvements. We consider that the main strength of this study was the exact definition of clinical response to TNFi treatment.

One of the main limitations of this study was not to include smoking duration and intensity to evaluate whether there was a dose-dependent effect. In addition, potential confounders such as the socioeconomic and exercise status of the patients were not included in our database.

In conclusion, our findings suggested that although the baseline disease activity was higher in smoker AS patients than nonsmokers, smoking had no impact on their first TNFi responses. Based on contradictory results in the literature, there is a need to investigate the association between smoking and response to TNFi in prospective studies that evaluated the effects of the amount and duration of smoking and its cessation on treatment. However, irrespective of the results of the studies, smoking cessation should be strongly recommended in all AS patients because of its negative effects on disease activity and radiographic progression and well-known risk for several comorbidities, including cardiovascular and chronic lung diseases.

Author contributions

All the authorship contributions were declared in line with the International Committee of Medical Journal Editors 4 criteria: FO, HYT: conceptualization, study design; SA, ED, SŞ, SSK, AT, AY, Nİ, NA, GK, HYT, BZ, GC, NA, FO: study advertisement, data acquisition; HYT, SÇ, HE, SA, FO: data analysis and interpretation. All authors approved the final submitted version of the manuscript.

References

1. Sugiyama D, Nishimura K, Tamaki K, Tsuji G, Nakazawa T et al. Impact of smoking as a risk factor for developing rheumatoid arthritis: a meta-analysis of observational studies. *Annals of the Rheumatic Diseases* 2010; 69 (1): 70-81. <https://doi.org/10.1136/ard.2008.096487>
2. Arnaud L, Tektonidou MG. Long-term outcomes in systemic lupus erythematosus: trends over time and major contributors. *Rheumatology (Oxford)* 2020; 59 (5): 29-38. <https://doi.org/10.1093/rheumatology/keaa382>.

3. Videm V, Cortes A, Thomas R, Brown MA. Current Smoking is Associated with Incident Ankylosing Spondylitis — The HUNT Population-based Norwegian Health Study. *The Journal of Rheumatology* 2014; 41 (10) 2041-2048. <https://doi.org/10.3899/jrheum.140353>
4. Poddubnyy D, Haibel H, Listing J, Marker-Hermann E, Zeidler H et al. Cigarette smoking has a dose-dependent impact on progression of structural damage in the spine in patients with axial spondyloarthritis: results from the GERMAN Spondyloarthritis Inception Cohort (GESPIC). *Annals of the Rheumatic Diseases* 2013; 72 (8): 1430-2. <https://doi.org/10.1136/annrheumdis-2012-203148>
5. Poddubnyy D, Haibel H, Listing J, Marker-Hermann E, Zeidler H et al. Baseline radiographic damage, elevated acute-phase reactant levels, and cigarette smoking status predict spinal radiographic progression in early axial spondylarthritis. *Arthritis & Rheumatism* 2012; 64 (5): 1388-98. <https://doi.org/10.1002/art.33465>
6. Maas F, Arends S, Wink FR, Bos R, Bootsma H et al. Ankylosing spondylitis patients at risk of poor radiographic outcome show diminishing spinal radiographic progression during long-term treatment with TNF-alpha inhibitors. *PLoS One* 2017; 12 (6): e0177231. <https://doi.org/10.1371/journal.pone.0177231>
7. Chung HY, Machado P, van der Heijde D, D'Agostino MA, Dougados M. Smokers in early axial spondyloarthritis have earlier disease onset, more disease activity, inflammation and damage, and poorer function and health-related quality of life: results from the DESIR cohort. *Annals of the Rheumatic Diseases* 2012; 71 (6): 809-16. <https://doi.org/10.1136/annrheumdis-2011-200180>
8. Villaverde-Garcia V, Cobo-Ibanez T, Candelas-Rodriguez G, Seoane-Mato D, Campo-Fontecha PDD et al. The effect of smoking on clinical and structural damage in patients with axial spondyloarthritis: A systematic literature review. *Seminars in Arthritis and Rheumatism* 2017; 46 (5): 569-83. <https://doi.org/10.1016/j.semarthrit.2016.11.004>
9. Braun J, Davis J, Dougados M, Sieper J, van der Linden S et al. First update of the international ASAS consensus statement for the use of anti-TNF agents in patients with ankylosing spondylitis. *Annals of the Rheumatic Diseases* 2006; 65 (3): 316-20. <https://doi.org/10.1136/ard.2005.040758>
10. Poddubnyy D, Sieper J. What is the best treatment target in axial spondyloarthritis: tumour necrosis factor alpha, interleukin 17, or both? *Rheumatology (Oxford)* 2017; 57 (7) 1145-1150. <https://doi.org/10.1093/rheumatology/kex361>
11. Navarro-Compán V, Plasencia-Rodríguez C, de Miguel E, Diaz Del Campo P, Balsa A et al. Switching biological disease-modifying antirheumatic drugs in patients with axial spondyloarthritis: results from a systematic literature review. *Rheumatic & Musculoskeletal Diseases Open* 2017; 3 (2): e000524. <https://doi.org/10.1136/rmdopen-2017-000524>
12. Lee JW, Kang JH, Yim YR, Kim JE, Wen L et al. Predictors of switching anti-tumor necrosis factor therapy in patients with ankylosing spondylitis. *PLoS One* 2015;10(7): e0131864. <https://doi.org/10.1371/journal.pone.0131864>
13. Gulyas K, Bodnar N, Nagy Z, Szamosi S, Horvath A et al. Real-life experience with switching TNF-alpha inhibitors in ankylosing spondylitis. *The European Journal of Health Economics* 2014; 15 (1): 93-100. <https://doi.org/10.1007/s10198-014-0598-0>
14. Alazmi M, Sari I, Krishnan B, Inman RD, Haroon N. Profiling Response to TNF-Inhibitor Treatment in Axial Spondyloarthritis. *Arthritis Care & Research (Hoboken)* 2017; 70 (9): 1393-1399. <https://doi.org/10.1002/acr.23465>
15. Zurita Prada PA, Urrego Laurín CL, Guillén Astete CA, Kanaffo Caltelblanco S, Navarro-Compán V. Influence of smoking and obesity on treatment response in patients with axial spondyloarthritis: a systematic literature review. *Clinical Rheumatology* 2021; 40 (5): 1673-86. <https://doi.org/10.1007/s10067-020-05319-6>
16. Machado P, Landewé R, Lie E, Kvien TK, Braun J et al. Ankylosing Spondylitis Disease Activity Score (ASDAS): defining cut-off values for disease activity states and improvement scores. *Annals of the Rheumatic Diseases* 2011 ;70 (1): 47-53. <https://doi.org/10.1136/ard.2010.138594>
17. Garrett S, Jenkinson T, Kennedy LG, Whitelock H, Gaisford P et al. A new approach to defining disease status in ankylosing spondylitis: the Bath Ankylosing Spondylitis Disease Activity Index. *The Journal of Rheumatology* 1994; 21: 2286-91.
18. Lukas C, Landewé R, Sieper J, Dougados M, Davis J et al. Development of an ASAS-endorsed disease activity score (ASDAS) in patients with ankylosing spondylitis. *Annals of the Rheumatic Diseases* 2009; 68 (1): 18-24. <https://doi.org/10.1136/ard.2008.094870>
19. Calin A, Garrett S, Whitelock H, Kennedy LG, O'Hea J et al. A new approach to defining functional ability in ankylosing spondylitis: the development of the Bath Ankylosing Spondylitis Functional Index. *The Journal of Rheumatology* 1994; 21 (12): 2281-5.
20. Jones SD, Porter J, Garrett SL, Kennedy LG, Whitelock H et al. A new scoring system for the Bath Ankylosing Spondylitis Metrology Index (BASMI). *The Journal of Rheumatology* 1995; 22 (8): 1609
21. Guillemin F, Briançon S, Poulrel J, Gaucher A. Long-term disability and prolonged sick leaves as outcome measurements in ankylosing spondylitis. Possible predictive factors. *Arthritis & Rheumatism* 1990; 33 (7): 1001-6. <https://doi.org/10.1002/art.1780330712>
22. van der Linden S, Valkenburg HA, Cats A. Evaluation of diagnostic criteria for ankylosing spondylitis. A proposal for modification of the New York criteria. *Arthritis & Rheumatism* 1984; 27 (4): 361-8. <https://doi.org/10.1002/art.1780270401>
23. Ward MM. Predictors of the progression of functional disability in patients with ankylosing spondylitis. *The Journal of Rheumatology* 2002; 29 (7): 1420-5.
24. Reed MD, Dharmage S, Boers A, Martin BJ, Buchanan RR et al. Ankylosing spondylitis: an Australian experience. *Internal Medicine Journal* 2008; 38 (5): 321-7. <https://doi.org/10.1111/j.1445-5994.2007.01471.x>

25. Matthey DL, Dawson SR, Healey EL, Packham JC. Relationship between smoking and patient-reported measures of disease outcome in ankylosing spondylitis. *The Journal of Rheumatology* 2011; 38 (12): 2608-15. <https://doi.org/10.3899/jrheum.110641>
26. Kaan U, Ferda O. Evaluation of clinical activity and functional impairment in smokers with ankylosing spondylitis. *Rheumatology International* 2005; 25 (5): 357-60. <https://doi.org/10.1007/s00296-004-0451-3>
27. Jones GT, Ratz T, Dean LE, Macfarlane GJ, Atzeni F. Disease Severity in Never Smokers, Ex-Smokers, and Current Smokers With Axial Spondyloarthritis: Results From the Scotland Registry for Ankylosing Spondylitis. *Arthritis Care & Research (Hoboken)* 2017; 69 (9): 1407-13. <https://doi.org/10.1002/acr.23157>
28. Chen CH, Chen HA, Lu CL, Liao HT, Liu CH et al. Association of cigarette smoking with Chinese ankylosing spondylitis patients in Taiwan: a poor disease outcome in systemic inflammation, functional ability, and physical mobility. *Clinical Rheumatology* 2013; 32 (5): 659-63. <https://doi.org/10.1007/s10067-013-2165-y>
29. Ward MM, Hendrey MR, Malley JD, Learch TJ, Davis JC et al. Clinical and immunogenetic prognostic factors for radiographic severity in ankylosing spondylitis. *Arthritis Care & Research* 2009; 61 (7): 859-66. <https://doi.org/10.1002/art.24585>
30. Ramiro S, Landewe R, van Tubergen A, Boonen A, Stolwijk C et al. Lifestyle factors may modify the effect of disease activity on radiographic progression in patients with ankylosing spondylitis: a longitudinal analysis. *Rheumatic & Musculoskeletal Diseases Open* 2015; 1 (1): e000153. <https://doi.org/10.1136/rmdopen-2015-000153>
31. Fallahi S JAR, Gharibdoost F, Mahmoudi M, Ahmadzadeh N, Nicknam M H. The Correlation between Pack-Years of Smoking and Disease Activity, Quality of Life, Spinal Mobility, and Sacroiliitis Grading in Patients with Ankylosing Spondylitis. *Turkish Journal of Rheumatology* 2013; 28 (3): 181-8. <https://doi.org/10.5606/tjr.2013.3269>
32. Haroon N, Inman RD, Learch TJ, Weisman MH, Lee M et al. The impact of tumor necrosis factor alpha inhibitors on radiographic progression in ankylosing spondylitis. *Arthritis & Rheumatism* 2013; 65 (10): 2645-54. <https://doi.org/10.1002/art.38070>
33. Zhao S, Yoshida K, Jones GT, Hughes DM, Tedeschi SK et al. The impact of smoking on response to TNF inhibitors in axial spondyloarthritis: methodological considerations for longitudinal observational studies. *Arthritis Care & Research (Hoboken)* 2019; 72 (4): 591-599. <https://doi.org/10.1002/acr.23851>
34. Grintborg B, Hojgaard P, Lund Hetland M, Steen Krogh N, Kollerup G et al. Impact of tobacco smoking on response to tumour necrosis factor-alpha inhibitor treatment in patients with ankylosing spondylitis: results from the Danish nationwide DANBIO registry. *Rheumatology (Oxford)* 2016; 55 (4): 659-68. <https://doi.org/10.1093/rheumatology/kev392>
35. Ciurea A, Scherer A, Weber U, Exer P, Bernhard J et al. Impaired response to treatment with tumour necrosis factor alpha inhibitors in smokers with axial spondyloarthritis. *Annals of the Rheumatic Diseases* 2016; 75 (3): 532-9. <https://doi.org/10.1136/annrheumdis-2013-205133>
36. Lord PA, Farragher TM, Lunt M, Watson KD, Symmons DP et al. Predictors of response to anti-TNF therapy in ankylosing spondylitis: results from the British Society for Rheumatology Biologics Register. *Rheumatology (Oxford)* 2010; 49 (3): 563-70. <https://doi.org/10.1093/rheumatology/kep422>
37. Kydd AS, Chen JS, Makovey J, Chand V, Henderson L et al. Smoking did not modify the effects of anti-TNF treatment on health-related quality of life among Australian ankylosing spondylitis patients. *Rheumatology (Oxford)* 2015; 54 (2): 310-7. <https://doi.org/10.1093/rheumatology/keu314>
38. Menegatti S, Bianchi E, Rogge L. Anti-TNF Therapy in Spondyloarthritis and Related Diseases, Impact on the Immune System and Prediction of Treatment Responses. *Frontiers in Immunology* 2019; 10: 382. <https://doi.org/10.3389/fimmu.2019.00382>