



Access this article online

Quick Response Code:



Website:

<https://turkjemergmed.com/>

DOI:

10.4103/tjem.tjem\_73\_24

Submitted: 12-04-2024

Revised: 05-06-2024

Accepted: 05-06-2024

Published: 01-07-2024

ORCID:

AŞ: 0000-0002-0583-2936

MP: 0000-0002-3917-0192

MM: 0000-0001-6928-2307

EA: 0000-0002-3249-2420

SK: 0000-0002-8692-7266

GA: 0000-0002-5580-3201

NÖD: 0000-0002-5209-8076

BÖ: 0000-0002-6645-7287

MH: 0000-0001-6368-6853

ÇT: 0009-0001-5534-5781

SEE: 0000-0002-3183-3713

YY: 0000-0003-3917-9434

NA: 0000-0001-8059-4862

NGG: 0000-0001-5052-9242

NNU: 0000-0003-3509-3254

HA: 0000-0002-8396-4854

ASG: 0000-0003-2669-0532

SK: 0000-0001-8265-9615

SA: 0000-0002-2197-4715

AÖ: 0000-0002-2707-4099

GPG: 0000-0001-8531-4591

MBS: 0000-0001-9335-9001

ÖS: 0000-0002-5557-6627

SBHB: 0000-0002-7695-

9969

**Address for  
correspondence:**

Dr. Alp Şener,  
Department of Emergency  
Medicine, Faculty of  
Medicine, Ankara Bilkent  
City Hospital, Ankara  
Yıldırım Beyazıt University,  
Bilkent, Ankara, Türkiye.  
E-mail: alpsener@  
gmail.com

# Prospective, multicenter, Turkish out-of-hospital cardiac arrest study: TROHCA

Alp Şener<sup>1,2\*</sup>, Murat Pekdemir<sup>3</sup>, Mehmet Muzaffer İslam<sup>4</sup>, Ersin Aksay<sup>5</sup>, Sevilay Karahan<sup>6</sup>, Gokhan Aksel<sup>4</sup>, Nurettin Özgür Doğan<sup>3</sup>, Berkant Öztürk<sup>7</sup>, Muhammet Hacimustafaoğlu<sup>7</sup>, Çağrı Türkücü<sup>7</sup>, Serkan Emre Eroğlu<sup>4</sup>, Yusuf Yürümez<sup>8</sup>, Nuray Aslan<sup>8</sup>, Necip Gokhan Güner<sup>8</sup>, Neşe Nur User<sup>9</sup>, Hüseyin Aldemir<sup>9</sup>, Abdullah Sadık Girişgin<sup>10</sup>, Sedat Koçak<sup>10</sup>, Sami Ataman<sup>10</sup>, Ayhan Özhasenekler<sup>1,2</sup>, Gul Pamukçu Günaydın<sup>1,2</sup>, Mustafa Burak Sayhan<sup>11</sup>, Ömer Salt<sup>11</sup>, Satuk Bugra Han Bozatlı<sup>11</sup>, Engin Deniz Arslan<sup>12</sup>, Fevzi Yılmaz<sup>12</sup>, Ramazan Sivil<sup>12</sup>, Özlem Köksal<sup>13</sup>, Vahide Aslıhan Durak<sup>13</sup>, Fatma Özdemir<sup>13</sup>, Mahmut Taş<sup>14</sup>, Yenal Karakoç<sup>14</sup>, Öner Avınca<sup>14</sup>, Yunus Emre Arık<sup>15</sup>, Adem Melekoğlu<sup>15</sup>, Özgür Çevrim<sup>15</sup>, Özlem Yiğit<sup>16</sup>, Cem Oktay<sup>16</sup>, Süleyman İbze<sup>16</sup>, Salim Satar<sup>17</sup>, Muge Gülen<sup>17</sup>, Selen Acehan<sup>17</sup>, Erhan Altunbaş<sup>18</sup>, Melis Efeoğlu Saçak<sup>18</sup>, Emir Ünal<sup>18</sup>, Erdem Çevik<sup>19</sup>, Dilay Satılmış<sup>19</sup>, Hande Asan<sup>19</sup>, Yunus Karaca<sup>20</sup>, Melih İmamoğlu<sup>20</sup>, Vildan Özer<sup>20</sup>, Ahmet Demircan<sup>21</sup>, Ayfer Keleş<sup>21</sup>, Gültekin Kadı<sup>21</sup>, Orhan Delice<sup>22</sup>, Sibel Güçlü Utlu<sup>22</sup>, Senol Arslan<sup>22</sup>, Neslihan Yücel<sup>23</sup>, Şükrü Gürbüz<sup>23</sup>, Hüseyin Burak Ayhan<sup>23</sup>, Abdullah Şen<sup>24</sup>, Mahmut Yaman<sup>24</sup>, Müge Günalp<sup>25</sup>, Sinan Genç<sup>25</sup>, Ahmet Baydın<sup>26</sup>, Fatih Çalışkan<sup>26</sup>, Şeyma Arzu Temür<sup>26</sup>, Murat Ersel<sup>27</sup>, Sercan Yalçınlı<sup>27</sup>, Enver Özçete<sup>27</sup>, Bulent Erbil<sup>28</sup>, Elif Ozturk Ince<sup>28</sup>, Mehmet Ali Karaca<sup>28</sup>, Murat Çetin<sup>29</sup>, Mehmet Demirbağ<sup>30</sup>, Mustafa Sabak<sup>31</sup>, Mustafa Bozkurt<sup>31</sup>

<sup>1</sup>Department of Emergency Medicine, Ankara Bilkent City Hospital, Ankara, Türkiye, <sup>2</sup>Department of Emergency Medicine, Ankara Yıldırım Beyazıt University, Faculty of Medicine, Ankara, Türkiye, <sup>3</sup>Department of Emergency Medicine, Kocaeli University, Faculty of Medicine, Kocaeli, Türkiye, <sup>4</sup>University of Health Sciences, Umraniye Training and Research Hospital, İstanbul, Türkiye, <sup>5</sup>Department of Emergency Medicine, İzmir Ekonomi University, Medical Point Hospital, İzmir, Türkiye, <sup>6</sup>Department of Biostatistics, Hacettepe University, Faculty of Medicine, Ankara, Türkiye, <sup>7</sup>Department of Emergency Medicine, Aksaray University, Aksaray Training and Research Hospital, Aksaray, Türkiye, <sup>8</sup>Department of Emergency Medicine, Sakarya University, Faculty of Medicine, Sakarya, Türkiye, <sup>9</sup>Department of Emergency Medicine, Afyonkarahisar Health Sciences University, Afyonkarahisar, Türkiye, <sup>10</sup>Department of Emergency Medicine, Necmettin Erbakan University, Faculty of Medicine, Konya, Türkiye, <sup>11</sup>Department of Emergency Medicine, Trakya University, Faculty of Medicine, Edirne, Türkiye, <sup>12</sup>Department of Emergency Medicine, University of Health Sciences, Antalya Training and Research Hospital, Antalya, Türkiye, <sup>13</sup>Department of Emergency Medicine, Bursa Uludağ University, Faculty of Medicine, Bursa, Türkiye, <sup>14</sup>Department of Emergency Medicine, University of Health Sciences, Gazi Yaşargil Training and Research Hospital, Diyarbakır, Türkiye, <sup>15</sup>Department of Emergency Medicine, University of Health Sciences, İstanbul Şişli Hamidiye Etfal Training and Research Hospital, İstanbul, Türkiye, <sup>16</sup>Department of Emergency Medicine, Akdeniz University, Faculty of Medicine, Antalya, Türkiye, <sup>17</sup>Department of Emergency Medicine, University of Health Sciences, Adana City Training and Research Hospital, Adana, Türkiye, <sup>18</sup>Department of Emergency Medicine, Marmara University Pendik Training and Research Hospital, İstanbul, Türkiye, <sup>19</sup>Department of Emergency Medicine, University of Health Sciences, Sultan 2. Abdülhamid Han Training and Research Hospital, İstanbul, Türkiye, <sup>20</sup>Department of Emergency Medicine, Karadeniz Technical University, Faculty of Medicine, Trabzon, Türkiye, <sup>21</sup>Department of Emergency Medicine, Gazi University, Faculty of Medicine, Ankara, Türkiye, <sup>22</sup>Department of Emergency Medicine, Erzurum City Hospital, Erzurum, Türkiye, <sup>23</sup>Department of Emergency Medicine, İnönü University, Turgut Özal Medical Center, Malatya, Türkiye, <sup>24</sup>Department of Emergency Medicine, Dicle University, Faculty of Medicine, Diyarbakır, Türkiye, <sup>25</sup>Department of Emergency Medicine, Ankara University, Faculty of Medicine, Ankara, Türkiye, <sup>26</sup>Department of Emergency Medicine, Ondokuz Mayıs University, Faculty of Medicine, Samsun, Türkiye, <sup>27</sup>Department of Emergency

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

**How to cite this article:** Şener A, Pekdemir M, İslam MM, Aksay E, Karahan S, Aksel G, *et al.* Prospective, multicenter, Turkish out-of-hospital cardiac arrest study: TROHCA. Turk J Emerg Med 2024;24:133-44.

EDA: 0000-0002-0155-6903 *Medicine, Ege University, Faculty of Medicine, İzmir, Türkiye,* <sup>28</sup>*Department of Emergency Medicine, Hacettepe University, Faculty of Medicine, Ankara, Türkiye,* <sup>29</sup>*Dr.Behçet Uz Pediatric Diseases and Surgery Training and Research Hospital, İzmir, Türkiye,* <sup>30</sup>*Department of Emergency Medicine, Manisa Merkezefendi State Hospital, Manisa, Türkiye,* <sup>31</sup>*Department of Emergency Medicine, Gaziantep University, Şahinbey Research and Practice Hospital, Gaziantep, Türkiye*  
FY: 0000-0002-3675-7457  
RS: 0000-0002-6463-2117  
ÖK: 0000-0003-2271-5659  
VAD: 0000-0003-0836-7862  
FÖ: 0000-0002-4278-1576  
MT: 0000-0002-3894-4766  
YK: 0000-0003-3607-9286  
ÖA: 0000-0003-3109-6773  
YEA: 0000-0002-4521-9546  
AM: 0000-0002-5764-3678  
ÖÇ: 0000-0002-5515-5232  
ÖY: 0000-0002-3703-7590  
CO: 0000-0001-7993-0566  
Sİ: 0000-0002-0927-0029  
SS: 0000-0001-6080-4287  
MG: 0000-0002-5080-3501  
SA: 0000-0003-4390-6593  
EA: 0000-0002-5075-739X  
MES: 0000-0002-8446-9898  
EÜ: 0000-0003-2758-8214  
EÇ: 0000-0003-1474-3166  
DS: 0000-0003-3765-2208  
HA: 0000-0002-6742-302X  
YK: 0000-0002-0947-0565  
Mİ: 0000-0003-4197-8999  
VÖ: 0000-0002-2514-5674  
AD: 0000-0002-7993-8098  
AK: 0000-0003-4429-3100  
GK: 0000-0002-9195-6635  
OD: 0000-0003-1629-4245  
SGU: 0000-0001-8635-1562  
SA: 0000-0002-6636-5307  
NY: 0000-0001-5845-2614  
ŞG: 0000-0003-2616-0304  
HBA: 0000-0002-8876-1378  
AŞ: 0000-0001-6705-4079  
MY: 0000-0002-0783-1419  
MG: 0000-0001-8140-6720  
SG: 0000-0002-0516-1028  
AB: 0000-0003-4987-0878  
FÇ: 0000-0001-7786-3929  
ŞAT: 0009-0003-6564-2434  
ME: 0000-0003-2282-5559  
SY: 0000-0003-1780-8391  
EÖ: 0000-0002-1685-2369  
BE: 0000-0001-8555-0017  
EÖİ: 0000-0002-6396-6668  
MAK: 0000-0003-0876-2811  
MÇ: 0000-0002-0667-7966  
MD: 0000-0002-0475-872X  
MS: 0000-0003-2777-2003  
MB: 0000-0002-4400-6696

### Abstract:

**OBJECTIVES:** There is no sufficient data to provide a clear picture of out-of-hospital cardiac arrest (OHCA) across Türkiye. This study is the first to present the prognostic outcomes of OHCA cases and the factors associated with these outcomes.

**MATERIALS AND METHODS:** The study was conducted in a prospective, observational, multicenter design under the leadership of the Emergency Medicine Association of Turkey Resuscitation Study Group. OHCA cases aged 18 years and over who were admitted to 28 centers from Türkiye were included in the study. Survived event, return of spontaneous circulation (ROSC), survival to hospital discharge, and neurological outcome at discharge were investigated as primary outcomes.

**RESULTS:** One thousand and three patients were included in the final analysis. 61.1% of the patients were male, and the average age was  $67.0 \pm 15.2$ . Cardiopulmonary resuscitation (CPR) was performed on 86.5% of the patients in the prehospital period by emergency medical service, and bystander CPR was performed on only 2.9% by nonhealth-care providers. As a result, the survived event rate was found to be 6.9%. The survival rate upon hospital discharge was 4.4%, with 2.7% of patients achieving a good neurological outcome upon discharge. In addition, the overall ROSC and sustained ROSC rates were 45.2% and 33.4%, respectively. In the multiple logistic regression analysis, male gender, initial shockable rhythm, a shorter prehospital duration of CPR, and the lack of CPR requirement in the emergency department were determined to be independent predictors for the survival to hospital discharge.

**CONCLUSION:** Compared to global data, survival to hospital discharge and good neurological outcome rates appear to be lower in our study. We conclude that this result is related to low bystander CPR rates. Although not the focus of this study, inadequate postresuscitative care and intensive care support should also be discussed in this regard. It is obvious that this issue should be carefully addressed through political moves in the health and social fields.

### Keywords:

Bystander cardiopulmonary resuscitation, cardiopulmonary resuscitation, cardiopulmonary resuscitation, out-of-hospital cardiac arrest, registry, return of spontaneous circulation, survival, survived event, Turkey, Türkiye

### Box-ED section

#### What is already known about the study topic?

- Out-of-hospital cardiac arrest (OHCA) remains a global public health problem. Although there are varying rates, survival rates are far from satisfactory. Türkiye's data on this subject are limited; therefore, a multicenter study was needed.

#### How is this study structured?

- The study was planned under the supervision of the Emergency Medicine Association of Turkey Resuscitation Study Group for 1-year period with a prospective, observational, multicenter design in OHCA patients aged 18 years and over.

#### What does this study tell us?

- In this study, the rate of survival to

hospital discharge for OHCA was found to be 4.4%; there are similar results in the literature. However, it is thought provoking that bystander cardiopulmonary resuscitation (CPR) was applied only in 2.9% of the patients.

#### What is the conflict on the issue? Is it important for readers?

- Bystander-initiated CPR is one of the key points of a chain of survival. In our country, where the prehospital system and emergency medicine are well developed, the low rate of hospital discharge of out-of-cardiac arrest victims may be due to the low bystander CPR rate. Therefore, improved policies are needed in the political and health fields to enable the public to early recognize OHCA and start early bystander-initiated CPR.



arrest to the ED were included 24/7 in the study. Informed consent was obtained from all patients/relatives. Patients experiencing traumatic cardiac arrest, those transferred to study centers from other hospitals, defined as deceased prior to admission, whose primary outcome data couldn't be obtained, and whose relatives did not provide consent for their participation in the study, were excluded from the study. According to the sample size calculation, based on the ROSC (24.8%) and survival to hospital discharge (5.6%) rates in the study of Şener *et al.* and assuming a 5% error, a total of at least 837 cases were planned to be included in the study.<sup>[6]</sup> Upon reaching this sample, the study was terminated on October 15, 2023, before the planned date.

Ethical approval was received from the Kocaeli University Non-Interventional Clinical Research Ethics Committee (No: KÜ GOKAEK-2022/18.24) at the coordinator center on November 10, 2022, and the study was carried out in accordance with the principles of the World Medical Association Declaration of Helsinki.

### Data collection

Data obtained from the patients themselves or their relatives and recorded forms in Hospital Information Management Systems were checked and transferred to the online registry form (Google® Forms) by the local coordinator of the relevant center. Afterward, the relevant data were processed by the authorized researchers and directed for statistical analysis.

### Definitions and outcomes

The data collection forms were primarily designed based on the Utstein Resuscitation Registry Templates for out-of-hospital cardiac arrest (OHCA), with additional variables incorporated.<sup>[7]</sup> Obtaining any rhythm with pulse for at least 5 min during resuscitation was defined as ROSC, and persistence of this condition for at least 20 min was defined as sustained ROSC. Patients who were admitted to the ED with a pulse that persisted 20 min or who completed 20 min with a pulse in the ED were defined as a survived event. Patients who were discharged with sustained ROSC were recorded as survival to hospital discharge. Patients who were still in ROSC at day 30 were also recorded as 30-day survival. Cerebral performance category (CPC) was evaluated at discharge and on the 30<sup>th</sup> day, and according to this scale, categories 1 and 2 were recorded as good neurological outcome and 3 and above were recorded as poor neurological outcome. The hospital parameter as cardiac arrest localization refers to patients who had cardiac arrest at the hospital entrance (parking lot, garden, entrance of the outpatient clinics or triage, etc.) before being delivered to the relevant health-care professional. A cardiac arrest which is seen by other people or is monitored by EMS is defined as witnessed arrest. CPR procedure initiated by witnessed


people other than the EMS staff is defined as bystander CPR. The rhythm of the patient detected in the first monitoring was recorded as initial cardiac arrest rhythm, and pulseless ventricular tachycardia and ventricular fibrillation rhythms were recorded as shockable rhythm. The concept of airway procedure was used as the most advanced method performed. The concept of unsuccessful airway procedure refers to a patient who cannot be ventilated in any way. Survived event, ROSC, survival to hospital discharge, and neurological outcome at discharge were investigated as primary outcomes.

### Statistical analysis

Statistical analyses of the study were performed with IBM SPSS Statistics for Windows, Version 26.0 (Armonk, NY, USA: IBM Corp). In comparing continuous data between two independent groups, the Mann-Whitney *U*-test or Independent samples *t*-test was used, depending on distribution. Continuous data are expressed as mean with standard deviation or median with interquartile range in accordance with distribution. The comparison of categorical data between independent groups was performed with Pearson's Chi-square and Fisher's exact tests; these data were expressed as sample numbers and percentages. In primary analyses,  $P < 0.05$  level was used for statistical significance. Multiple logistic regression analysis was performed for sustained ROSC and discharge, and variables with  $P < 0.250$  in univariate analysis but thought to be clinically significant were included in the model. For discharge outcome, "prehospital CPR, arrest cause, prehospital airway management, arrest location, use of magnesium sulfate in the emergency department (ED), and experiencing cardiac arrest again in the ED" were unable to be included in the model due to imbalances in sample counts across groups. Similarly, for sustained ROSC, "prehospital CPR, arrest etiology, prehospital airway management, and CPR administered in the ED" couldn't be included due to group imbalances. As a result, odds ratios were expressed with 95% confidence intervals. Cutoff values of pH variable used in logistic regression analysis were determined by ROC analysis.

### Results

A total of 1060 patients were recorded in the data system from 28 centers in Türkiye. After ruling out missing data and samples that did not meet the criteria (such as trauma patients, inhospital arrests, and referrals from other hospitals), 1002 patients were included in the final analysis [Figure 2]. It was determined that 61.1% of the patients were male, the mean age was  $67.0 \pm 15.2$  years, the etiology was acute coronary syndrome in 46.3%, and the localization of cardiac arrest was "home" in 61.3%. Witnessed arrest was detected in 83.9% of patients, transport with EMS in 97.4%, and shockable first arrest rhythm in 10.8%. CPR was performed on 86.5% of the



EMS: Emergency medical services, OHCA: Out-of-hospital cardiac arrest, CPR: Cardio-pulmonary resuscitation, ROSC: Return of spontaneous circulation, ED: Emergency department

Figure 2: Flow diagram

patients in the prehospital period by EMS personnel, and bystander CPR was performed on only 2.9% by nonhealth-care providers (higher in male gender; 3.8 vs. 1.5). While the prehospital bag-valve-mask rate was 31.7%, successful endotracheal intubation (ETI) was found to be 47.6%. Peripheral intravenous access was presented in 88.4% of the patients, and 11.2% were delivered to the ED without any venous access [Table 1].

### Primary outcomes

As a result, while the prehospital ROSC rate was 13.6%, the survived event rate was found to be 6.9% [Table 1].

While the mortality rate in the ED was 72.3%, 4.4% were discharged alive from the hospital and 2.7% were discharged with good neurological outcome. In addition, the ROSC rate for the entire process was determined as 45.2% and the sustained ROSC rate was 33.4%.

### Secondary outcomes

In the EDs, mechanical chest compression device was used in 41.5% of patients, EtCO<sub>2</sub> monitoring in 30.1%, and ultrasound-guided CPR performed in 52.1% of patients [Table 2]. Primary comparative

**Table 1: Demographic characteristics and prehospital descriptive variables**

| Variables                                    | n (%) or mean±SD or median (25%–75%) |
|--|--------------------------------------|
| Time zone                                    |                                      |
| Day shift                                    | 386 (38.5)                           |
| Night shift and holidays                     | 616 (61.5)                           |
| Gender                                       |                                      |
| Female                                       | 390 (38.9)                           |
| Male   | 612 (61.1)                           |
| Age (year), mean±SD                          | 67.0±15.2                            |
| Etiology                                     |                                      |
| Acute coronary syndrome                      | 464 (46.3)                           |
| Electrolyte disorder                         | 51 (5.1)                             |
| Pulmonary embolism                           | 43 (4.3)                             |
| Asthma                                       | 16 (1.6)                             |
| Drowning                                     | 16 (1.6)                             |
| Intoxication                                 | 9 (0.9)                              |
| Stroke                                       | 9 (0.9)                              |
| Electrocution                                | 3 (0.3)                              |
| Anaphylaxis                                  | 2 (0.2)                              |
| Postprimary coronary intervention            | 2 (0.2)                              |
| Postbypass                                   | 2 (0.2)                              |
| Others                                       | 140 (14)                             |
| Unknown                                      | 245 (24.5)                           |
| Independent living                           | 721 (77.4)                           |
| Chronic neurological disease                 | 168 (17.9)                           |
| Chronic pulmonary disease                    | 183 (19.4)                           |
| Chronic cardiovascular disease               | 496 (52.3)                           |
| Active malignancy                            | 126 (13.4)                           |
| Ventricular assist device                    | 7 (0.7)                              |
| ICD (or external cardioverter defibrillator) | 24 (2.5)                             |
| Cardiac arrest location                      |                                      |
| Home   | 603 (61.3)                           |
| Workplace                                    | 15 (1.5)                             |
| Public area                                  | 120 (12.2)                           |
| Nursing home                                 | 36 (3.7)                             |
| Ambulance                                    | 148 (15.1)                           |
| Hospital                                     | 40 (4.1)                             |
| Others                                       | 21 (2.1)                             |
| Witnessed arrest                             | 776 (83.9)                           |
| Transport with EMS                           | 975 (97.4)                           |
| Instructions from EMS dispatcher             | 98 (15.5)                            |
| Prehospital CPR                              |                                      |
| None   | 105 (10.6)                           |
| Health-care provider                         | 859 (86.5)                           |
| Bystander CPR                                | 29 (2.9)                             |
| Prehospital CPR duration (min)               | 15 (8–20)                            |
| Time (min) from the arrest to the CPR        | 5 (1–10)                             |
| Initial cardiac arrest rhythm                |                                      |
| Asystole                                     | 682 (78.1)                           |
| PEA  | 97 (11.1)                            |
| VF   | 85 (9.7)                             |
| pVT  | 9 (1)                                |
| Shockable rhythm                             |                                      |
| pVT/VF                                       | 94 (10.8)                            |

Contd...

**Table 1: Contd...**

| Variables  | n (%) or mean±SD or median (25%–75%) |
|--|--------------------------------------|
| Asystole/PEA   | 779 (89.2)                           |
| Prehospital defibrillation                             | 118 (12.1)                           |
| Number of prehospital defibrillations                  | 2 (1–3)                              |
| Time (min) from the arrest to the first defibrillation | 6.5 (3–12.75)                        |
| Prehospital airway procedure                           |                                      |
| Not performed  | 117 (11.7)                           |
| BVM  | 318 (31.7)                           |
| ETI - successful                                       | 477 (47.6)                           |
| Supraglottic   | 54 (5.4)                             |
| Surgical   | 4 (0.4)                              |
| Unsuccessful   | 32 (3.2)                             |
| Prehospital chest compression                          | 888 (89.4)                           |
| Prehospital ventilation support                        | 899 (90.8)                           |
| Prehospital mechanical CPR device                      | 9 (0.9)                              |
| Prehospital venous access                              |                                      |
| None   | 112 (11.2)                           |
| Peripheral   | 886 (88.4)                           |
| Intraosseous   | 2 (0.2)                              |
| Central  | 2 (0.2)                              |
| Prehospital vascular access - exists                   | 890 (88.8)                           |
| Prehospital epinephrine                                | 741 (76.6)                           |
| Prehospital epinephrine dose (mg)                      | 3 (2–5)                              |
| Prehospital amiodarone                                 | 25 (2.5)                             |
| Prehospital lidocaine                                  | 1 (0.1)                              |
| Prehospital sodium bicarbonate                         | 5 (0.5)                              |
| Prehospital calcium                                    | 2 (0.2)                              |
| Reported by EMS before ED admission                    | 710 (75.3)                           |
| Presence of pulse at ED admission - ROSC               | 136 (13.6)                           |
| Pulse duration in ED admission, if exists (min)        |                                      |
| <20  | 100 (77.5)                           |
| ≥20  | 29 (22.5)                            |
| Survived event   |                                      |
| No   | 933 (93.1)                           |
| Yes  | 69 (6.9)                             |

SD: Standard deviation, ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardiopulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, BVM: Bag-valve mask, ETI: Endotracheal intubation, ED: Emergency department, ROSC: Return of the spontaneous circulation

analyses between the two groups for the outcomes of survived event, overall sustained ROSC, survival to hospital discharge, and discharge with good neurological outcome are provided as supplemental files [Supplementary Tables 1-8]. It has been determined that the survival to hospital discharge rate is higher in male gender and patients with shockable rhythm and is lower, especially in the group of home localization and with mechanical chest compression devices [Supplementary Table 5]. In addition, it has been determined that both survived event, survival and good neurological outcome rates were higher in cases where bystander CPR was performed [Supplementary Tables 1, 5, and 7].

**Table 2: Descriptive variables of emergency department period**

| Variables   | n (%) or median (25%–75%) |
|---|---------------------------|
| First monitored rhythm at ED                            |                           |
| Asystole  | 682 (68.3)                |
| PEA   | 124 (12.4)                |
| VF  | 75 (7.5)                  |
| pVT   | 4 (0.4)                   |
| Pulse exists (any rhythm)                               | 113 (11.3)                |
| Shockable rhythm  |                           |
| pVT/VF  | 79 (7.9)                  |
| Asystole/PEA  | 806 (80.8)                |
| Pulse exists (any rhythm)                               | 113 (11.3)                |
| Advanced airway in ED                                   |                           |
| Exists at admission                                     | 432 (43.1)                |
| BVM   | 8 (0.8)                   |
| ETI   | 547 (54.6)                |
| Supraglottic  | 7 (0.7)                   |
| Surgical  | 8 (0.8)                   |
| Venous access in ED                                     |                           |
| None  | 1 (0.1)                   |
| Peripheral  | 943 (94.1)                |
| Intraosseous  | 6 (0.6)                   |
| Central   | 52 (5.2)                  |
| Cardiac arrest again in ED                              |                           |
| No  | 232 (23.3)                |
| Yes   | 218 (21.9)                |
| No ROSC in ED   | 547 (54.9)                |
| CPR time (min) to the first ROSC in ED                  | 13 (7.5–20)               |
| CPR in ED   | 950 (94.8)                |
| Mechanical CPR device in ED                             | 416 (41.5)                |
| Defibrillation in ED                                    | 229 (22.9)                |
| Number of defibrillations before first ROSC in ED       | 2 (1–3)                   |
| Total number of defibrillations in ED                   | 3 (1–4)                   |
| Epinephrine in ED                                       | 941 (93.9)                |
| Epinephrine dose before the first ROSC in ED (mg)       | 6 (3–10)                  |
| Total epinephrine dose in ED (mg)                       | 6 (7–15)                  |
| Amiodarone in ED  | 157 (15.7)                |
| Lidocaine in ED   | 5 (0.5)                   |
| Magnesium sulfate in ED                                 | 21 (2.1)                  |
| Sodium bicarbonate in ED                                | 251 (25)                  |
| Calcium in ED   | 139 (13.9)                |
| USG use during CPR in ED                                | 522 (52.1)                |
| Cardiac activity detected by USG                        | 172 (33.1)                |
| EtCO <sub>2</sub> device use during CPR in ED           | 302 (30.1)                |
| 5 <sup>th</sup> min ETCO <sub>2</sub> during CPR (mmHg) | 14 (10–21.5)              |
| Last ETCO <sub>2</sub> during CPR (mmHg)                | 14 (8–22)                 |
| Highest ETCO <sub>2</sub> during CPR (mmHg)             | 22 (15–34)                |
| Initial pH in ED  | 6.98 (6.85–7.08)          |
| Initial lactate (mmol/L) in ED                          | 11.80 (8.04–15.40)        |
| Outcome in ED   |                           |
| Exitus  | 724 (72.3)                |
| Ward/ICU  | 212 (21.2)                |
| Transferred to other hospitals                          | 66 (6.6)                  |

Contd...

**Table 2: Contd...**

| Variables                             | n (%) or median (25%–75%) |
|---------------------------------------|---------------------------|
| Outcome in hospital                   |                           |
| Exitus                                | 958 (95.6)                |
| Discharged                            | 44 (4.4)                  |
| Neurological outcome at discharge     |                           |
| Poor                                  | 967 (97.3)                |
| Good                                  | 27 (2.7)                  |
| 30-day survival                       |                           |
| Exitus                                | 951 (94.9)                |
| Survived                              | 51 (5.1)                  |
| 30-day neurological outcome           |                           |
| Poor                                  | 966 (97.2)                |
| Good                                  | 28 (2.8)                  |
| ST-segment elevation after ROSC       | 144 (34.4)                |
| Coronary revascularization after ROSC | 100 (23.9)                |
| TTM after ROSC                        | 9 (2.2)                   |
| Vasoactive drug after ROSC            | 271 (65.1)                |
| ECLS after ROSC                       | 1 (0.2)                   |
| IABP after ROSC                       | 0                         |
| Surgery after ROSC                    | 8 (2.0)                   |
| First pH after ROSC                   | 7.05 (6.92–7.22)          |
| First lactate (mmol/L) after ROSC     | 9.85 (5.59–13.93)         |
| ROSC any (overall)                    | 453 (45.2)                |
| Sustained ROSC (overall)              | 334 (33.4)                |
| Forensic case report                  | 100 (10.0)                |

PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, BVM: Bag-valve mask, ETI: Endotracheal intubation, ROSC: Return of the spontaneous circulation, ED: Emergency department, med: median, CPR: Cardiopulmonary resuscitation, min: minute, USG: Ultrasonography, EtCO<sub>2</sub>: End-tidal carbon dioxide, ICU: Intensive care unit, TTM: Targeted temperature management, ECLS: Extracorporeal Life Support, IABP: Intra-aortic balloon pump

According to regional distribution, the highest bystander CPR rates were found in the Marmara and Aegean regions and the highest survival rates in the Mediterranean and Black Sea regions. The good neurological outcome rate is higher in the Mediterranean region [Supplementary Table 9].

### Multiple logistic regression analysis

As a result of multiple regression analyses, male gender (odds ratio [OR]: 2.691; 95% confidence interval [CI]: 1.186–6.103), prehospital initial shockable rhythm (OR: 6.480; 95% CI: 3.055–13.744), a shorter prehospital duration of CPR (OR: 0.919; 95% CI: 0.880–0.961), and the lack of CPR requirement in the ED (OR: 12.038; 95% CI: 5.000–28.984) were found to be positive independent predictors for the survival to hospital discharge [Table 3]. For sustained ROSC, presence of chronic pulmonary disease (OR: 1.677; 95% CI: 1.088–2.585), witnessed arrest (OR: 2.910; 95% CI: 1.588–5.336), arrest localization being hospital (OR: 3.703; 95% CI: 1.310–10.467; reference: home), shorter prehospital CPR duration (OR: 0.954; 95% CI: 0.933–0.975), not using mechanical chest compression device in ED (OR: 2.598; 95% CI: 1.774–3.803), the venous route used in the ED

**Table 3: Multiple regression analysis for survival to hospital discharge**

| Variables  | OR     | 95% CI       | P      |
|--|--------|--------------|--------|
| Day shift versus night shift and holidays          | 1.992  | 1.013–3.918  | 0.053  |
| Gender (male vs. female)                           | 2.691  | 1.186–6.103  | 0.018  |
| Transport with EMS (EMS vs. not)                   | 0.444  | 0.113–1.742  | 0.244  |
| Prehospital initial shockable rhythm versus others | 6.480  | 3.055–13.744 | <0.001 |
| Prehospital CPR duration (min)                     | 0.919  | 0.880–0.961  | <0.001 |
| CPR in ED (none vs. performed)                     | 12.038 | 5.000–28.984 | <0.001 |
| Sodium bicarbonate in ED (none vs. performed)      | 2.590  | 0.871–7.704  | 0.087  |

CI: Confidence interval, EMS: Emergency medical services, CPR: Cardiopulmonary resuscitation, ED: Emergency department, OR: Odds ratio

**Table 4: Multiple regression analysis for overall sustained return of the spontaneous circulation**

| Variables   | OR    | 95% CI       | P      |
|---|-------|--------------|--------|
| Chronic pulmonary disease                                 | 1.677 | 1.088–2.585  | 0.019  |
| Witnessed arrest (yes vs. no)                             | 2.910 | 1.588–5.336  | 0.001  |
| Transport with EMS (EMS vs. non-EMS)                      | 0.764 | 0.263–2.222  | 0.622  |
| Cardiac arrest location (reference: Home)                 |       |              |        |
| Workplace/public area                                     | 0.594 | 0.335–1.055  | 0.076  |
| Nursing home  | 0.853 | 0.298–2.444  | 0.767  |
| Ambulance   | 0.987 | 0.605–1.609  | 0.958  |
| Hospital  | 3.703 | 1.310–10.467 | 0.014  |
| Others  | 0.797 | 0.266–2.386  | 0.685  |
| Prehospital CPR duration (min)                            | 0.954 | 0.933–0.975  | <0.001 |
| Shockable rhythm at ED admission (reference: pVT/VF)      |       |              |        |
| Asystole/PEA  | 0.517 | 0.257–1.043  | 0.066  |
| Pulse exists  | 2.031 | 0.840–4.911  | 0.116  |
| Advanced airway in ED (ref: Exists at admission)          |       |              |        |
| BVM/supraglottic/surgical                                 | 0.151 | 0.017–1.304  | 0.086  |
| ETI   | 0.824 | 0.555–1.225  | 0.339  |
| Mechanical CPR device in ED (not performed vs. performed) | 2.598 | 1.774–3.803  | <0.001 |
| Defibrillation in ED (performed vs. not)                  | 1.583 | 0.984–2.547  | 0.058  |
| Venous access in ED (others vs. peripheral)               | 2.105 | 1.051–4.219  | 0.036  |
| Initial pH during CPR in ED (reference: pH<6.9)           |       |              |        |
| $6.9 \leq \text{pH} < 7.0$                                | 2.106 | 1.257–3.528  | 0.005  |
| $\text{pH} \geq 7.0$                                      | 3.005 | 1.941–4.652  | <0.001 |

CI: Confidence interval, EMS: Emergency medical services, CPR: Cardiopulmonary resuscitation, ED: Emergency department, pVT: Pulseless ventricular tachycardia, VF: Ventricular fibrillation, PEA: Pulseless electrical activity, BVM: Bag-valve mask, ETI: Endotracheal intubation

being other than peripheral intravenous route (OR: 2.105; 95% CI: 1.051–4.219), and higher pH levels (for “ $6.9 \leq \text{pH} < 7.0$ ” OR: 2.106; 95% CI: 1.257–3.528 and for “ $\text{pH} \geq 7.0$ ” OR: 3.005; 95% CI: 1.941–4.652) were determined as independent predictors [Table 4].

## Discussion

This descriptive prospective multicenter study investigated the epidemiology and outcomes of OHCA patients in prehospital and emergency settings across Türkiye and aims to provide a comprehensive overview of this critical patient population. Of the OHCA patients, 453 (45.2%) achieved any ROSC, 334 (33.4%) achieved sustained ROSC, and 278 (27.7%) had survived event, which were the outcomes that primarily pertain to prehospital and emergency medicine department settings. Upon reviewing our findings, it can be concluded that our results closely align with

data reported in European and Australian studies. The EuRoCa TWO study, published in 2020, documented that any ROSC was achieved in 32.7% of 25,171 patients who experienced cardiac arrest and received CPR initiated by EMS or bystanders.<sup>[1]</sup> Similar findings have also been reported (23.8-37.8% for prehospital ROSC) in Australian and New Zealand data published by Beck *et al.*<sup>[8]</sup>

In a meta-analysis that examines 141 studies and nearly 4.6 million OHCA patients, the rate of sustained ROSC was reported as 29.7%. The authors conducted subgroup analyses by region and observed the highest sustained ROSC rate in Oceania countries (38.6%), followed by Europe (36.7%), and then Asian countries (22.1%).<sup>[9]</sup> While our sustained ROSC rates may not match those observed in Oceania and Europe, it can be inferred that our results are closely approximated. In the meta-analysis by Yan *et al.*, the global incidence of survived event rates among OHCA patients was reported as 22%. Oceanic countries exhibited the highest rate of survived events at 33.5%,

while Asian countries showed the lowest at 15.6%.<sup>[9]</sup> Comparatively, British and Australian data suggest a survived event rate of 22% and 28%, respectively, while European data indicate a higher survived event rate of 35%.<sup>[1,8,10]</sup> Upon the examination of prehospital and emergency setting data, Türkiye's outcomes appear to closely align with studies conducted in developed countries.

Upon closer examination of the primary outcome measures, markedly different results stand out compared to those observed in the aforementioned studies. Of the OHCA patients in our study, 44 (4.4%) were discharged from the hospital. We can argue that our results were notably inferior compared to the findings reported in the literature conducted in the developed countries. Yan *et al.* reported a global hospital discharge rate of 8.8% in OHCA patients, which is approximately double the rate observed in our findings. In this meta-analysis, subgroup analyses by region revealed the highest survival to discharge rate in Oceania countries (16.2%) and the lowest in Asian countries (4.5%).<sup>[9]</sup> The rates of survival to discharge in Australia and New Zealand were reported to be 12.1%, while European countries reported rates of 8% and England 7.2%.<sup>[1,8,10]</sup> Consequently, our findings closely parallel the outcomes observed in Asian countries regarding survival to hospital discharge.

An intriguing detail worth noting from the EuRoCa TWO study, was a specific subgroup outcome. Among patients brought to the hospital while CPR was ongoing, the survival to hospital discharge rate was notably lower at 4%. All patients who received CPR exhibited an 8% survival to hospital discharge rate, whereas those brought directly to the hospital showed a 26% survival rate, and individuals achieving ROSC in the prehospital setting demonstrated a notably higher rate of 35%.<sup>[1]</sup> One potential explanation for the low survival-to-hospital discharge rate observed in our data could be the predominance of patients within this subgroup.

2021 data by Kotini-Shah *et al.* indicate that the rate of favorable neurological outcome of OHCA patients in the United States is 8.7%.<sup>[11]</sup> However, similar to the findings regarding survival to hospital discharge, our study revealed a relatively low rate of favorable neurological outcomes, with only 27 patients (2.7%) demonstrating positive results. We can observe similar rates in Asian countries in this aspect. In the study conducted by Okubo *et al.* in 2014, which analyzed OHCA data from a sizable cohort, the rate of favorable neurological outcomes was reported to be only 2%.<sup>[12]</sup>

It is interesting to note that while our outcomes that are related to the prehospital and emergency care are notably similar to the literature, our rates of discharge

and favorable neurological survival are considerably low. This observation prompts consideration of potential factors influencing these relatively long-term outcomes.

The factors contributing to lower survival rates in Türkiye compared to global data warrant further investigation. While factors such as ED crowding in Türkiye may play a role, the lack of intensive care unit capacity in certain regions, coupling with the need for transfer of the critical patients, may have contributed to the observed poor outcomes. A comprehensive examination of all potential causes is necessary. It is imperative to escalate quality improvement initiatives nationwide to address this issue effectively. While the effectiveness of CPR is pivotal for inpatient discharge and favorable neurological outcomes, its success hinges on the coordination of multiple disciplines and is subject to various confounding factors. In this study, while ROSC rates in EMS and EDs align with global data, the underlying factors contributing to lower rates of live discharges and favorable neurological outcomes warrant investigation. These results suggest the possibility that patients in Türkiye may lack adequate support from intensive care and other multidisciplinary aspects in subsequent stages of treatment. Future studies should conduct detailed analyses to address this issue, and we believe the data presented here can significantly contribute to enhancing CPR success nationwide.

In the EuRoCa TWO study, approximately 58.8% of OHCA patients who received CPR had bystander CPR, leading to significantly higher rates of ROSC and discharge compared to those in whom EMS initiated CPR.<sup>[1]</sup> Similarly, data from Australia and New Zealand, as reported by Beck *et al.* in 2018, indicated that bystander CPR was administered in 67% of OHCA cases.<sup>[8]</sup> Despite relatively lower rates in British data, standing at 39.5%, these rates still surpass those observed in our study.<sup>[10]</sup> The notable discrepancy in bystander CPR rates in our data may be attributed to cultural disparities and variations in CPR awareness levels.<sup>[13]</sup> Barriers to access to AEDs and the lack of awareness of AED use remain important handicaps in Türkiye. In addition, the absence of Good Samaritan Law rules in Turkey may partially explain the hesitations about performing CPR in cardiac arrest cases.

We identified several independent predictors of survival to hospital discharge in the logistic regression analysis, including male gender, recorded initial shockable rhythm, a shorter prehospital duration of CPR, and the lack of CPR requirement in the ED.

The male gender predominance, which varies between 56% and 65% in the literature, is also evident in this study.<sup>[1,11,12]</sup> In the study of Kotini-Shah *et al.*, it was

observed that better results were obtained in the male gender in terms of survival to hospital discharge and neurological outcome, but in multivariate analysis, the adjusted odds ratios were seen to be in favor of the female gender.<sup>[11]</sup> On the contrary, we observed better outcomes in male patients; we can attribute these results to the higher rates of bystander CPR in male gender.

The presence of an initial shockable rhythm as an independent predictor of survival to hospital discharge has been observed in various studies in different settings.<sup>[14-17]</sup> Additionally, in Li *et al.*'s data from Singapore, which shares similar characteristics and settings with our study and demonstrates a survival rate of 3.4%, shockable initial rhythm and shorter prehospital CPR duration were also identified as independent predictors.<sup>[14]</sup> It is evident that increasing the rate of bystander-initiated CPR can lead to improved outcome rates, highlighting the necessity for a nationwide educational campaign.

The presence of chronic pulmonary disease, witnessed arrest, cardiac arrest occurring at hospital perimeters, brief prehospital CPR duration, absence of mechanical CPR device uses in the ED, utilization of a venous access other than the peripheral intravenous route, and higher pH levels in the first blood gas obtained during CPR were identified as independent predictors of sustained ROSC.

A recent study by Balan *et al.* reported that witnessed cardiac arrest, initial shockable rhythm, and cardiac arrest occurring outside of home were found to be the independent predictors of hospital discharge in 3952 OHCA patients who achieved sustained ROSC.<sup>[16]</sup> Although our study populations and outcomes may differ, it is noteworthy that we obtained similar results.

The debate regarding whether mechanical or manual chest compressions yield different outcomes in cardiac arrest patients remains a hot topic of current research. Only a single 2015 randomized controlled study reported significant harm of mechanical compression device in OHCA patients in terms of survival with favorable neurological outcome.<sup>[18]</sup> A 2018 Cochrane review categorized existing studies as of medium-to-low quality and concluded that both methods were not superior to each other except in certain circumstances. It is important to note that this review included evaluations of both in-hospital and OHCA patients, with trauma patients excluded.<sup>[19]</sup> Therefore, our findings may not align precisely with those of the aforementioned review.

We found that the sustained ROSC rate was significantly higher in patients who received prehospital respiratory support, particularly among those who underwent successful ETI before arrival at the hospital. However,

the observational design of our study limits our ability to make definitive conclusions based solely on these findings. For instance, the recent AIRWAY-2 study investigated the impact of prehospital ETI and supraglottic airway (SGA) placement on 30-day survival in OHCA patients, finding no significant difference between the two methods.<sup>[20]</sup>

In our study, we observed a significantly higher rate of sustained ROSC among patients who received epinephrine prior to hospital arrival. This finding aligns with the PARAMEDIC2 study, a large-scale investigation conducted in 2018, which demonstrated higher rates of ROSC, survival to hospital admission, and 30-day survival in OHCA patients treated with prehospital epinephrine administration.<sup>[21]</sup> Consequently, the use of epinephrine in OHCA patients has gained stronger support, as reflected in the 2023 update of the AHA advanced cardiac life support guidelines.<sup>[22]</sup>

### Limitations

First, it should be noted that significant missing data exist, and consecutive sampling could not be properly implemented. Disruption of the operational functioning of study centers due to the earthquake on February 6, 2023, was a major contributing factor to these missing data. Additionally, two of the study centers had to be excluded from the study due to data unavailability. However, it has not been assessed whether this situation negatively impacts the regional representation of the country. The inclusion of tertiary hospitals in the study may introduce bias when extrapolating the results to the general public and rural ED.

Patients who received CPR outside and were not brought to the study centers were not included in the study; excluding these data also increases the bias factor. Most previous registry studies included only patients transported by EMS.<sup>[1,8-10,12]</sup> Although the rate of patients transported by non-EMS people is very low (2.6%), the inclusion of this patient group should be taken into consideration when discussing results with previous literature. Since reliable data could not be obtained about the time for prehospital transport, it was not included in the analysis. In most of the patients, the possible etiology of cardiac arrest was estimated by the physician based on the available history. Lack of knowledge about exact no-flow time (for lay person) may have a negative influence on results.

Another point, the hospital arrest localization group causes confusion for the results. In this group, cardiac arrest occurs at the triage or the hospital entrance. Therefore, it is understood that interventions such as prehospital chest compressions and respiratory support are not applied in a significant part of this group, and it

is understood that the prognosis is better, probably due to rapid access to health-care providers.

Given the study's nonrandomized controlled trial design, it is important to acknowledge that certain subgroup analysis findings may be influenced by confounding factors. For instance, while the lower success rate of CPR with mechanical devices might appear accurate, it is plausible that this outcome could be influenced by factors such as the preference for mechanical CPR devices in patients with a lower life expectancy and longer CPR duration.

## Conclusion

This study is the first multicenter, prospective study conducted in Türkiye revealing OHCA data. As a result, prognostic data obtained in the prehospital and ED periods are similar to those in developed countries; however, survival and good neurological outcome rates appear to be worse. Concepts such as ED crowding and intensive care quality can be discussed here. However, success in hospital discharge and neurological survival is an issue that requires multidisciplinary coordination. Although it is not the focus of this study, the correct approach would be to discuss these poor outcomes by focusing on the postcardiac arrest care step in the chain of survival.

Consistent with the literature, shockable initial rhythm and witnessed arrest were found to be associated with good outcomes. On the other hand, the bystander CPR rate, which is very low compared to developed countries, should be particularly scrutinized and efforts should be made to improve it by developing health policies and social policies.

### Acknowledgments

The researchers would like to thank the Emergency Medicine Association of Türkiye Resuscitation Study Group Members for their valuable contribution to the study.

### Author contributions

All authors conducted the study concept and design and acquisition of the data. AŞ and SK conducted the statistical analysis and interpretation of the data. AŞ and MMİ conducted the drafting of the manuscript. All authors conducted critical revision of the manuscript for important intellectual content.

### Conflicts of interest

None declared.

### Ethical approval

This multicenter study was carried out in 28 health centers in Türkiye within the scope of the Emergency Medicine Association of Türkiye (EMAT) Resuscitation Study Group, and approval was obtained from the Kocaeli University Non-Interventional Clinical Research Ethics Committee (KÜ GOKAEK-2022/18.24; November 10, 2022).

### Funding

None.

## References

- Gräsner JT, Wnent J, Herlitz J, Perkins GD, Lefering R, Tjelmeland I, et al. Survival after out-of-hospital cardiac arrest in Europe – Results of the EuReCa TWO study. *Resuscitation* 2020;148:218-26.
- Panchal AR, Bartos JA, Cabañas JG, Donnino MW, Drennan IR, Hirsch KG, et al. Part 3: Adult basic and advanced life support: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2020;142:S366-468.
- Merchant RM, Topjian AA, Panchal AR, Cheng A, Aziz K, Berg KM, et al. Part 1: Executive summary: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2020;142:S337-57.
- Tekin FC, Köylü R, Köylü Ö, Kunt M. Factors related to resuscitation success and prognosis of cardiopulmonary arrest cases. *Indian J Crit Care Med* 2023;27:26-31.
- Halhalli HC, Şancı E, Uslu T. The comparison of manual and mechanical chest compression on survival and long-term neurological outcome of nontraumatic out-of-hospital cardiac arrest patients. *J Emerg Med* 2020;59:680-6.
- Şener A, Pamukçu Günaydın G, Tanrıverdi F. Mechanical versus manual chest compression: A retrospective-cohort in out-of-hospital cardiac arrest. *Acta Med* 2021;52:325-31.
- Perkins GD, Jacobs IG, Nadkarni VM, Berg RA, Bhanji F, Biarent D, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: Update of the Utstein resuscitation registry templates for out-of-hospital cardiac arrest: A statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian and New Zealand Council on Resuscitation, Heart and Stroke Foundation of Canada, Inter American Heart Foundation, Resuscitation Council of Southern Africa, Resuscitation Council of Asia); and the American Heart Association Emergency Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. *Circulation* 2015;132:1286-300.
- Beck B, Bray J, Cameron P, Smith K, Walker T, Grantham H, et al. Regional variation in the characteristics, incidence and outcomes of out-of-hospital cardiac arrest in Australia and New Zealand: Results from the Aus-ROC epistry. *Resuscitation* 2018;126:49-57.
- Yan S, Gan Y, Jiang N, Wang R, Chen Y, Luo Z, et al. The global survival rate among adult out-of-hospital cardiac arrest patients who received cardiopulmonary resuscitation: A systematic review and meta-analysis. *Crit Care* 2020;24:61.
- Hawkes C, Booth S, Ji C, Brace-McDonnell SJ, Whittington A, Mapstone J, et al. Epidemiology and outcomes from out-of-hospital cardiac arrests in England. *Resuscitation* 2017;110:133-40.
- Kotini-Shah P, Del Rios M, Khosla S, Pugach O, Vellano K, McNally B, et al. Sex differences in outcomes for out-of-hospital cardiac arrest in the United States. *Resuscitation* 2021;163:6-13.
- Okubo M, Kiyohara K, Iwami T, Callaway CW, Kitamura T. Nationwide and regional trends in survival from out-of-hospital cardiac arrest in Japan: A 10-year cohort study from 2005 to 2014. *Resuscitation* 2017;115:120-8.
- Mentzelopoulos SD, Bossaert L, Raffay V, Askitopoulou H, Perkins GD, Greif R, et al. A survey of key opinion leaders on ethical resuscitation practices in 31 European countries. *Resuscitation* 2016;100:11-7.
- Li T, Yap J, Chng WQ, Tay JC, Shahidah N, Yeo C, et al. Clinicopathological correlates of out-of-hospital cardiac arrests. *J Arrhythm* 2022;38:416-24.
- Whittaker A, Lehal M, Calver AL, Corbett S, Deakin CD, Gray H, et al. Predictors of inhospital mortality following out-of-hospital cardiac arrest: Insights from a single-centre consecutive case

- series. *Postgrad Med J* 2016;92:250-4.
16. Balan P, Hsi B, Thangam M, Zhao Y, Monlezun D, Arain S, et al. The cardiac arrest survival score: A predictive algorithm for in-hospital mortality after out-of-hospital cardiac arrest. *Resuscitation* 2019;144:46-53.
  17. Baldi E, Caputo ML, Savastano S, Burkart R, Klersy C, Benvenuti C, et al. An utstein-based model score to predict survival to hospital admission: The UB-ROSC score. *Int J Cardiol* 2020;308:84-9.
  18. Perkins GD, Lall R, Quinn T, Deakin CD, Cooke MW, Horton J, et al. Mechanical versus manual chest compression for out-of-hospital cardiac arrest (PARAMEDIC): A pragmatic, cluster randomised controlled trial. *Lancet* 2015;385:947-55.
  19. Wang PL, Brooks SC. Mechanical versus manual chest compressions for cardiac arrest. *Cochrane Database Syst Rev* 2018;8:CD007260.
  20. Bengier JR, Kirby K, Black S, Brett SJ, Clout M, Lazaroo MJ, et al. Effect of a strategy of a supraglottic airway device versus tracheal intubation during out-of-hospital cardiac arrest on functional outcome: The AIRWAYS-2 randomized clinical trial. *JAMA* 2018;320:779-91.
  21. Perkins GD, Ji C, Deakin CD, Quinn T, Nolan JP, Scomparin C, et al. A randomized trial of epinephrine in out-of-hospital cardiac arrest. *N Engl J Med* 2018;379:711-21.
  22. Perman SM, Elmer J, Maciel CB, Uzendu A, May T, Mumma BE, et al. 2023 American Heart Association Focused Update on Adult Advanced Cardiovascular Life Support: An update to the American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2024;149:e254-73.

**Supplementary Table 1: Factors associated with survived event**

| Variables                                    | Survived event, n (%) | P     |
|--|-----------------------|-------|
| Time zone                                    |                       |       |
| Day shift                                    | 24 (6.2)              | 0.594 |
| Night shift and holidays                     | 45 (7.3)              |       |
| Gender                                       |                       |       |
| Female                                       | 34 (8.7)              | 0.068 |
| Male   | 35 (5.7)              |       |
| Independent living                           |                       |       |
| No   | 13 (6.2)              | 0.709 |
| Yes  | 52 (7.2)              |       |
| Chronic neurological disease                 |                       |       |
| No   | 56 (7.3)              | 0.666 |
| Yes  | 10 (6.0)              |       |
| Chronic pulmonary disease                    |                       |       |
| No   | 57 (7.5)              | 0.425 |
| Yes  | 10 (5.5)              |       |
| Chronic cardiovascular disease               |                       |       |
| No   | 26 (5.8)              | 0.242 |
| Yes  | 38 (7.7)              |       |
| Active malignancy                            |                       |       |
| No   | 61 (7.5)              | 0.212 |
| Yes  | 5 (4.0)               |       |
| Ventricular assist device                    |                       |       |
| No   | 66 (6.9)              | 0.399 |
| Yes  | 1 (14.3)              |       |
| ICD (or external cardioverter defibrillator) |                       |       |
| No   | 65 (6.9)              | 0.680 |
| Yes  | 2 (8.3)               |       |
| Witnessed arrest                             |                       |       |
| Yes  | 59 (7.6)              | 0.429 |
| No   | 8 (5.4)               |       |
| Instructions from EMS dispatcher             |                       |       |
| Yes  | 8 (8.2)               | 0.880 |
| No   | 38 (7.1)              |       |
| Prehospital CPR                              |                       |       |
| None   | 1 (1)                 | 0.016 |
| Health-care provider                         | 63 (7.3)              |       |
| Bystander CPR                                | 4 (13.8)              |       |
| Initial arrest rhythm                        |                       |       |
| Asystole                                     | 46 (6.7)              | 0.811 |
| PEA  | 5 (5.2)               |       |
| VF   | 7 (8.2)               |       |
| pVT  | 1 (11.1)              |       |
| Shockable rhythm                             |                       |       |
| pVT/VF                                       | 8 (8.5)               | 0.618 |
| Asystole/PEA                                 | 51 (6.5)              |       |
| Prehospital defibrillation                   |                       |       |
| No   | 58 (6.8)              | 0.408 |
| Yes  | 11 (9.3)              |       |
| Prehospital chest compression                |                       |       |
| None   | 1 (1)                 | 0.020 |
| Performed                                    | 67 (7.5)              |       |

**Supplementary Table 1: Contd...**

| Variables                           | Survived event, n (%) | P       |
|-------------------------------------|-----------------------|---------|
| Prehospital ventilation support     |                       |         |
| None                                | 2 (2.2)               | 0.097   |
| Performed                           | 67 (7.5)              |         |
| Prehospital mechanical CPR device   |                       |         |
| None                                | 66 (6.7)              | 0.122   |
| Used                                | 2 (22.2)              |         |
| Prehospital vascular access         |                       |         |
| None                                | 2 (1.8)               | 0.039   |
| Yes                                 | 67 (7.5)              |         |
| Prehospital epinephrine             |                       |         |
| None                                | 4 (1.8)               | 0.001   |
| Performed                           | 63 (8.5)              |         |
| Prehospital amiodarone              |                       |         |
| None                                | 65 (6.8)              | 0.685   |
| Performed                           | 2 (8)                 |         |
| Prehospital lidocaine               |                       |         |
| None                                | 67 (6.8)              | 0.069   |
| Performed                           | 1 (100)               |         |
| Prehospital sodium bicarbonate      |                       |         |
| None                                | 66 (6.7)              | 0.041   |
| Performed                           | 2 (40)                |         |
| Prehospital calcium                 |                       |         |
| None                                | 68 (6.9)              | 1.000   |
| Performed                           | -                     |         |
| Reported by EMS before ED admission |                       |         |
| Yes                                 | 47 (6.6)              | 1.000   |
| No                                  | 15 (6.4)              |         |
| Etiology                            |                       |         |
| Acute coronary syndrome             | 30 (6.5)              | 0.129   |
| Pulmonary embolism                  | 6 (14)                |         |
| Electrolyte disorder                | 3 (5.9)               |         |
| Asthma                              | 3 (18.8)              |         |
| Drowning                            | 3 (18.8)              |         |
| Others, unknown                     | 24 (5.8)              |         |
| Prehospital airway procedure        |                       |         |
| Not performed                       | 3 (2.6)               | <0.001* |
| BVM                                 | 13 (4.1)              |         |
| ETI - successful/surgical           | 51 (10.6)             |         |
| Supraglottic                        | 1 (1.9)               |         |
| Unsuccessful                        | 1 (3.1)               |         |
| Cardiac arrest location             |                       |         |
| Home                                | 43 (7.1)              | 0.239   |
| Workplace/public area               | 13 (9.6)              |         |
| Nursing home                        | 3 (8.3)               |         |
| Ambulance                           | 6 (4.1)               |         |
| Hospital                            | 3 (7.5)               |         |
| Others                              | -                     |         |

Pearson's Chi-square test, Fisher's exact test. ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, ED: Emergency department, BVM: Bag-valve mask, ETI: Endotracheal intubation

Contd...

**Supplementary Table 2: Factors associated with survived event - continues variables**

| <b>Variables</b>                                       | <b>Nonsurvived event</b> | <b>Survived event</b> | <b>P*</b> |
|--|--------------------------|-----------------------|-----------|
| Age (year)   | 67.1±15.1                | 65.9±15.7             | 0.655     |
| Prehospital CPR duration (min)                         | 15 (8–20)                | 15 (5.3–20)           | 0.897     |
| Time from the arrest to the CPR (min)                  | 5 (1–10)                 | 10 (2–10)             | 0.714     |
| Number of defibrillation (if exist)                    | 2 (1–3)                  | 2 (1–3)               | 0.712     |
| Time from the arrest to the first defibrillation (min) | 6.5 (3–11.8)             | 6.5 (2.3–15.3)        | 0.966     |

\*Age: Independent samples *t*-test, mean±SD. Variables other than age: Mann–Whitney *U*-test, median (25%–75%). CPR: Cardio-pulmonary resuscitation, SD: Standard deviation

**Supplementary Table 3: Factors associated with sustained return of the spontaneous circulation (overall)**

| Variables                                    | Sustained ROSC, n (%) | P      |
|--|-----------------------|--------|
| Time zone                                    |                       |        |
| Day shift                                    | 115 (29.9)            | 0.065  |
| Night shift and holidays                     | 219 (35.6)            |        |
| Gender                                       |                       |        |
| Female                                       | 136 (35.1)            | 0.388  |
| Male   | 198 (32.4)            |        |
| Independent living                           |                       |        |
| No   | 62 (29.5)             | 0.070  |
| Yes  | 261 (36.3)            |        |
| Chronic neurological disease                 |                       |        |
| No   | 264 (34.3)            | 0.693  |
| Yes  | 55 (32.7)             |        |
| Chronic pulmonary disease                    |                       |        |
| No   | 244 (32.1)            | 0.009  |
| Yes  | 77 (42.3)             |        |
| Chronic cardiovascular disease               |                       |        |
| No   | 154 (34.1)            | 0.708  |
| Yes  | 163 (33.0)            |        |
| Active malignancy                            |                       |        |
| No   | 278 (34.1)            | 0.685  |
| Yes  | 40 (32.3)             |        |
| Ventricular assist device                    |                       |        |
| No   | 320 (33.7)            | 0.049  |
| Yes  | 5 (71.4)              |        |
| ICD (or external cardioverter defibrillator) |                       |        |
| No   | 320 (34.0)            | 1.000  |
| Yes  | 8 (33.3)              |        |
| Witnessed arrest                             |                       |        |
| Yes  | 296 (38.2)            | <0.001 |
| No   | 26 (17.4)             |        |
| Instructions from EMS dispatcher             |                       |        |
| Yes  | 32 (32.7)             | 0.625  |
| No   | 187 (35.2)            |        |
| Prehospital CPR                              |                       |        |
| None   | 55 (52.4)             | <0.001 |
| Health-care provider                         | 264 (30.8)            |        |
| Bystander CPR                                | 11 (37.9)             |        |
| Prehospital initial cardiac arrest rhythm    |                       |        |
| Asystole                                     | 206 (30.2)            | <0.001 |
| PEA  | 38 (39.6)             |        |
| VF   | 46 (54.1)             |        |
| pVT  | 3 (33.3)              |        |
| Prehospital shockable rhythm                 |                       |        |
| pVT/VF                                       | 49 (52.1)             | <0.001 |
| Asystole/PEA                                 | 244 (31.4)            |        |
| Prehospital defibrillation                   |                       |        |
| No   | 283 (33.1)            | 0.367  |
| Yes  | 44 (37.3)             |        |
| Prehospital chest compression                |                       |        |
| None   | 55 (52.4)             | <0.001 |
| Performed                                    | 275 (31.1)            |        |
| Prehospital ventilation support              |                       |        |
| None   | 50 (54.9)             | <0.001 |
| Performed                                    | 282 (31.5)            |        |

**Supplementary Table 3: Contd...**

| Variables                           | Sustained ROSC, n (%) | P      |
|-------------------------------------|-----------------------|--------|
| Prehospital mechanical CPR device   |                       |        |
| None                                | 328 (33.4)            | 1.000  |
| Used                                | 3 (33.3)              |        |
| Prehospital vascular access         |                       |        |
| None                                | 47 (42.0)             | 0.042  |
| Yes                                 | 287 (32.4)            |        |
| Prehospital epinephrine             |                       |        |
| None                                | 104 (46.2)            | <0.001 |
| Performed                           | 226 (30.6)            |        |
| Prehospital amiodarone              |                       |        |
| None                                | 320 (33.5)            | 0.640  |
| Performed                           | 10 (40.0)             |        |
| Prehospital lidocaine               |                       |        |
| None                                | 330 (33.6)            | 0.337  |
| Performed                           | 1 (100)               |        |
| Prehospital sodium bicarbonate      |                       |        |
| None                                | 329 (33.6)            | 1.000  |
| Performed                           | 2 (40)                |        |
| Prehospital calcium                 |                       |        |
| None                                | 331 (33.7)            | 0.553  |
| Performed                           | -                     |        |
| Reported by EMS before ED admission |                       |        |
| Yes                                 | 206 (29.1)            | <0.001 |
| No                                  | 112 (48.3)            |        |
| Etiology                            |                       |        |
| Drowning                            | 11 (68.8)             | <0.001 |
| Acute coronary syndrome             | 156 (33.6)            |        |
| Pulmonary embolism                  | 19 (46.3)             |        |
| Electrolyte disorder                | 16 (31.4)             |        |
| Asthma                              | 12 (75)               |        |
| Others, unknown                     | 120 (29.2)            |        |
| Prehospital airway procedure        |                       |        |
| Not performed                       | 64 (54.7)             | <0.001 |
| BVM                                 | 82 (25.9)             |        |
| ETI - successful/surgical           | 170 (35.5)            |        |
| Supraglottic                        | 13 (24.1)             |        |
| Unsuccessful                        | 5 (15.6)              |        |
| Cardiac arrest location             |                       |        |
| Home                                | 177 (29.5)            | <0.001 |
| Workplace/public area               | 42 (31.1)             |        |
| Nursing home                        | 8 (22.2)              |        |
| Ambulance                           | 66 (44.9)             |        |
| Hospital                            | 31 (77.5)             |        |
| Others                              | 8 (38.1)              |        |
| Shockable rhythm at ED admission    |                       |        |
| pVT/VF                              | 44 (55.7)             | <0.001 |
| Asystole/PEA                        | 203 (25.3)            |        |
| Pulse exists                        | 85 (75.2)             |        |
| Advanced airway in ED               |                       |        |
| Exists at admission                 | 146 (34)              | 0.111  |
| BVM/Supraglottic/surgical           | 3 (13)                |        |
| ETI                                 | 185 (33.9)            |        |
| CPR in ED                           |                       |        |
| None                                | 52 (100)              | <0.001 |
| Performed                           | 282 (29.8)            |        |

Contd...

Contd...

**Supplementary Table 3: Contd...**

| <b>Variables</b>                              | <b>Sustained ROSC, n (%)</b> | <b>P</b> |
|---|------------------------------|----------|
| Mechanical CPR device in ED                   |                              |          |
| None  | 236 (40.4)                   | <0.001   |
| Used  | 98 (23.6)                    |          |
| Defibrillation in ED                          |                              |          |
| None  | 234 (30.4)                   | <0.001   |
| Performed                                     | 100 (43.7)                   |          |
| Venous access in ED                           |                              |          |
| Others  | 32 (54.2)                    | <0.001   |
| Peripheral                                    | 302 (32.1)                   |          |
| Epinephrine in ED                             |                              |          |
| None  | 61 (100)                     | <0.001   |
| Performed                                     | 273 (29.1)                   |          |
| Amiodarone in ED                              |                              |          |
| None  | 263 (31.2)                   | 0.001    |
| Performed                                     | 71 (45.2)                    |          |
| Lidocaine in ED                               |                              |          |
| None  | 332 (33.4)                   | 1.000    |
| Performed                                     | 2 (40)                       |          |
| Magnesium sulfate in ED                       |                              |          |
| None  | 324 (33.1)                   | 0.246    |
| Performed                                     | 10 (47.6)                    |          |
| Sodium bicarbonate in ED                      |                              |          |
| None  | 236 (31.5)                   | 0.022    |
| Performed                                     | 98 (39.4)                    |          |
| Calcium in ED                                 |                              |          |
| None  | 290 (33.7)                   | 0.678    |
| Performed                                     | 44 (31.9)                    |          |
| USG use during CPR in ED                      |                              |          |
| None  | 156 (32.6)                   | 0.578    |
| Used  | 178 (34.2)                   |          |
| If USG used, cardiac activity exists          |                              |          |
| None  | 34 (9.8)                     | <0.001   |
| Exists  | 143 (83.1)                   |          |
| EtCO <sub>2</sub> device use during CPR in ED |                              |          |
| None  | 257 (36.9)                   | <0.001   |
| Used  | 77 (25.5)                    |          |

Pearson's Chi-square test, Fisher's exact test. ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, ED: Emergency department, BVM: Bag-valve mask, ETI: Endotracheal intubation, USG: Ultrasonography, EtCO<sub>2</sub>: End-tidal carbon dioxide, ROSC: Return of the spontaneous circulation

**Supplementary Table 4: Factors associated with sustained return of the spontaneous circulation (overall) - continues variables**

| Variables  | No sustained ROSC | Sustained ROSC   | <i>P</i> * |
|--|-------------------|------------------|------------|
| Age (year)   | 67.4±15.4         | 66.2±14.9        | 0.242      |
| Prehospital CPR duration (min)                                     | 15 (10–25)        | 10 (3–20)        | <0.001     |
| Prehospital time from the arrest to the CPR (min)                  | 8 (2–10)          | 5 (1–10)         | <0.001     |
| Prehospital number of defibrillation (if exist)                    | 2 (1–3)           | 2 (1–3)          | 0.967      |
| Prehospital time from the arrest to the first defibrillation (min) | 10 (5.5–15)       | 4 (1–9)          | 0.001      |
| Number of defibrillations before first ROSC in ED                  | 2 (1–3)           | 3 (1–4)          | 0.074      |
| Total number of defibrillations in ED                              | 3 (1–4)           | 3 (1–4)          | 0.648      |
| Epinephrine dose (mg) before the first ROSC in ED                  | 10 (5–14)         | 4 (2–6)          | <0.001     |
| Total epinephrine dose (mg) in ED                                  | 13 (10–15)        | 6 (3–11)         | <0.001     |
| CPR to the first ROSC in ED (min)                                  | 12.5 (5.5–21.8)   | 14 (8–20)        | 0.843      |
| Initial pH during CPR in ED  | 6.93 (6.80–7.05)  | 7.02 (6.92–7.13) | <0.001     |
| Initial lactate during CPR in ED (mmol/L)                          | 12.5 (9–16)       | 10.24 (6.8–13.2) | <0.001     |
| 5 <sup>th</sup> min ETCO <sub>2</sub> during CPR in ED (mmHg)      | 12 (8.5–18)       | 18 (12–29.3)     | <0.001     |
| Last ETCO <sub>2</sub> during CPR in ED (mmHg)                     | 11 (6.75–16)      | 35 (24–54)       | <0.001     |
| Highest ETCO <sub>2</sub> during CPR in ED (mmHg)                  | 18 (13–25)        | 39 (30.5–55)     | <0.001     |

\*Age: Independent samples *t*-test, mean±SD. Variables other than age: Mann–Whitney *U*-test; median (25%–75%). ROSC: Return of the spontaneous circulation, CPR: Cardio-pulmonary resuscitation, ED: Emergency department, ETCO<sub>2</sub>: End-tidal carbon dioxide, SD: Standard deviation

**Supplementary Table 5: Factors associated with survival to hospital discharge**

| Variables                                    | Survival to hospital discharge, n (%) | P      |
|--|---------------------------------------|--------|
| Time zone                                    |                                       |        |
| Day shift                                    | 25 (6.5)                              | 0.017  |
| Night shift and holidays                     | 19 (3.1)                              |        |
| Gender                                       |                                       |        |
| Female                                       | 2 (2.3)                               | 0.016  |
| Male   | 35 (5.7)                              |        |
| Independent living                           |                                       |        |
| No   | 4 (1.9)                               | 0.044  |
| Yes  | 40 (5.5)                              |        |
| Chronic neurological disease                 |                                       |        |
| No   | 37 (4.8)                              | 0.629  |
| Yes  | 6 (3.6)                               |        |
| Chronic pulmonary disease                    |                                       |        |
| No   | 37 (4.9)                              | 0.469  |
| Yes  | 6 (3.3)                               |        |
| Chronic cardiovascular disease               |                                       |        |
| No   | 27 (6)                                | 0.061  |
| Yes  | 16 (3.2)                              |        |
| Active malignancy                            |                                       |        |
| No   | 42 (5.1)                              | 0.051  |
| Yes  | 1 (0.8)                               |        |
| Ventricular assist device                    |                                       |        |
| No   | 44 (4.6)                              | 1.000  |
| Yes  | -                                     |        |
| ICD (or external cardioverter defibrillator) |                                       |        |
| No   | 44 (4.7)                              | 0.622  |
| Yes  | -                                     |        |
| Witnessed arrest                             |                                       |        |
| Yes  | 37 (4.8)                              | 0.587  |
| No   | 5 (3.4)                               |        |
| Instructions from EMS dispatcher             |                                       |        |
| Yes  | 2 (2)                                 | 0.406  |
| No   | 24 (4.5)                              |        |
| Prehospital CPR                              |                                       |        |
| None   | 11 (10.5)                             | 0.001  |
| Health-care provider                         | 29 (3.4)                              |        |
| Bystander CPR                                | 4 (13.8)                              |        |
| Prehospital shockable rhythm                 |                                       |        |
| pVT/VF                                       | 17 (18.1)                             | <0.001 |
| Asystole/PEA                                 | 19 (2.4)                              |        |
| Prehospital defibrillation                   |                                       |        |
| No   | 31 (3.6)                              | 0.003  |
| Yes  | 12 (10.2)                             |        |
| Prehospital chest compression                |                                       |        |
| None   | 11 (10.5)                             | 0.004  |
| Performed                                    | 33 (3.7)                              |        |
| Prehospital ventilation support              |                                       |        |
| None   | 12 (13.2)                             | <0.001 |
| Performed                                    | 32 (3.6)                              |        |
| Prehospital mechanical CPR device            |                                       |        |
| None   | 43 (4.4)                              | 1.000  |
| Used   | -                                     |        |

**Supplementary Table 5: Contd...**

| Variables                           | Survival to hospital discharge, n (%) | P      |
|-------------------------------------|---------------------------------------|--------|
| Prehospital vascular access         |                                       |        |
| None                                | 9 (8)                                 | 0.081  |
| Yes                                 | 35 (3.9)                              |        |
| Prehospital epinephrine             |                                       |        |
| None                                | 21 (9.3)                              | <0.001 |
| Performed                           | 23 (3.1)                              |        |
| Prehospital amiodarone              |                                       |        |
| None                                | 43 (4.5)                              | 1.000  |
| Performed                           | 1 (4)                                 |        |
| Prehospital lidocaine               |                                       |        |
| None                                | 44 (4.5)                              | 1.000  |
| Performed                           | -                                     |        |
| Prehospital sodium bicarbonate      |                                       |        |
| None                                | 44 (4.5)                              | 1.000  |
| Performed                           | -                                     |        |
| Prehospital calcium                 |                                       |        |
| None                                | 44 (4.5)                              | 1.000  |
| Performed                           | -                                     |        |
| Survival event                      |                                       |        |
| None                                | 32 (3.4)                              | <0.001 |
| Yes                                 | 12 (17.4)                             |        |
| Reported by EMS before ED admission |                                       |        |
| Yes                                 | 25 (3.5)                              | 0.006  |
| No                                  | 19 (8.2)                              |        |
| Etiology                            |                                       |        |
| Drowning                            | 2 (12.5)                              | 0.085  |
| Acute coronary syndrome             | 28 (6)                                |        |
| Pulmonary embolism                  | 1 (2.3)                               |        |
| Electrolyte disorder                | 1 (2)                                 |        |
| Asthma                              | -                                     |        |
| Others, unknown                     | 12 (2.9)                              |        |
| Prehospital airway procedure        |                                       |        |
| Not performed                       | 17 (14.5)                             | <0.001 |
| BVM                                 | 10 (3.1)                              |        |
| ETI - successful/surgical           | 15 (3.1)                              |        |
| Supraglottic                        | 1 (1.9)                               |        |
| Unsuccessful                        | 1 (3.1)                               |        |
| Cardiac arrest location             |                                       |        |
| Home                                | 16 (2.7)                              | 0.002  |
| Workplace/public area               | 12 (8.9)                              |        |
| Nursing home                        | -                                     |        |
| Ambulance                           | 9 (6.1)                               |        |
| Hospital                            | 5 (12.5)                              |        |
| Others                              | 1 (4.8)                               |        |
| Shockable rhythm at ED admission    |                                       |        |
| pVT/VF                              | 13 (6.5)                              | <0.001 |
| Asystole/PEA                        | 17 (2.1)                              |        |
| Pulse exists                        | 14 (12.4)                             |        |
| Advanced airway in ED               |                                       |        |
| Exists at admission                 | 14 (3.2)                              | 0.295  |
| BVM/Supraglottic/Surgical           | 1 (4.3)                               |        |
| ETI                                 | 29 (5.3)                              |        |

Contd...

Contd...

**Supplementary Table 5: Contd...**

| Variables                             | Survival to hospital discharge, n (%) | P      |
|---------------------------------------|---------------------------------------|--------|
| CPR in ED                             |                                       |        |
| None                                  | 12 (23.1)                             | <0.001 |
| Performed                             | 32 (3.4)                              |        |
| Mechanical CPR device in ED           |                                       |        |
| None                                  | 33 (5.6)                              | 0.034  |
| Used                                  | 11 (2.6)                              |        |
| Defibrillation in ED                  |                                       |        |
| None                                  | 27 (3.5)                              | 0.018  |
| Performed                             | 17 (7.4)                              |        |
| Venous access in ED                   |                                       |        |
| Others                                | 3 (5.1)                               | 0.740  |
| Peripheral                            | 41 (4.3)                              |        |
| Epinephrine in ED                     |                                       |        |
| None                                  | 14 (23)                               | <0.001 |
| Performed                             | 30 (3.2)                              |        |
| Amiodarone in ED                      |                                       |        |
| None                                  | 30 (3.6)                              | 0.005  |
| Performed                             | 14 (8.9)                              |        |
| Lidocaine in ED                       |                                       |        |
| None                                  | 43 (4.3)                              | 0.201  |
| Performed                             | 1 (20)                                |        |
| Magnesium sulfate in ED               |                                       |        |
| None                                  | 40 (4.1)                              | 0.011  |
| Performed                             | 4 (19)                                |        |
| Sodium bicarbonate in ED              |                                       |        |
| None                                  | 40 (5.3)                              | 0.020  |
| Performed                             | 4 (1.6)                               |        |
| Calcium in ED                         |                                       |        |
| None                                  | 43 (5)                                | 0.040  |
| Performed                             | 1 (0.7)                               |        |
| Cardiac arrest again in ED            |                                       |        |
| None                                  | 40 (17.2)                             | <0.001 |
| Yes                                   | 4 (1.8)                               |        |
| No ROSC in ED                         | -                                     |        |
| USG use during CPR in ED              |                                       |        |
| None                                  | 21 (4.4)                              | 1.000  |
| Used                                  | 23 (4.4)                              |        |
| If USG used, cardiac activity exists  |                                       |        |
| None                                  | 2 (0.6)                               | <0.001 |
| Exists                                | 21 (12.2)                             |        |
| EtCO2 device use during CPR in ED     |                                       |        |
| None                                  | 34 (4.9)                              | 0.353  |
| Used                                  | 10 (3.3)                              |        |
| ST segment elevation after ROSC       |                                       |        |
| No                                    | 21 (7.7)                              | 0.014  |
| Yes                                   | 23 (16)                               |        |
| Coronary revascularization after ROSC |                                       |        |
| No                                    | 17 (5.3)                              | <0.001 |
| Yes                                   | 27 (27)                               |        |
| TTM after ROSC                        |                                       |        |
| No                                    | 39 (9.8)                              | 0.055  |
| Yes                                   | 3 (33.3)                              |        |

**Supplementary Table 5: Contd...**

| Variables                  | Survival to hospital discharge, n (%) | P     |
|----------------------------|---------------------------------------|-------|
| Vasoactive drug after ROSC |                                       |       |
| None                       | 22 (15.2)                             | 0.019 |
| Performed                  | 20 (7.4)                              |       |
| ECLS after ROSC            |                                       |       |
| None                       | 44 (10.7)                             | 1.000 |
| Performed                  | -                                     |       |
| Surgery after ROSC         |                                       |       |
| No                         | 43 (10.7)                             | 0.601 |
| Yes                        | 1 (12.5)                              |       |
| Forensic case report       |                                       |       |
| None                       | 39 (4.3)                              | 0.795 |
| Performed                  | 5 (5)                                 |       |

Pearson's Chi-square test, Fisher's exact test. ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, ED: Emergency department, BVM: Bag-valve mask, ETI: Endotracheal intubation, ROSC: Return of the spontaneous circulation, USG: Ultrasonography, EtCO2: End-tidal carbon dioxide, TTM: Targeted temperature management, ECLS: Extracorporeal life support

Contd...

**Supplementary Table 6: Factors associated with survival to hospital discharge - continues variables**

| Variables  | Exitus             | Discharged        | P*     |
|--|--------------------|-------------------|--------|
| Age (year)   | 67.3±15.2          | 61.0±14.7         | 0.007  |
| Prehospital CPR duration (min)                                     | 15 (10–20)         | 8.5 (0.13–12)     | <0.001 |
| Prehospital time from the arrest to the CPR (min)                  | 5 (1–10)           | 3 (1–7.3)         | 0.036  |
| Prehospital number of defibrillations (if exist)                   | 2 (1–3)            | 1.5 (1–2)         | 0.353  |
| Prehospital time from the arrest to the first defibrillation (min) | 9 (4–15)           | 2 (0.8–4.3)       | 0.001  |
| Number of defibrillations before first ROSC in ED                  | 2 (1–3)            | 3 (1–5.5)         | 0.198  |
| Total number of defibrillations in ED                              | 3 (1–4)            | 3 (1–4.5)         | 0.711  |
| Epinephrine dose (mg) before the first ROSC in ED                  | 6.5 (3–10)         | 3 (2–5)           | <0.001 |
| Total epinephrine dose (mg) in ED                                  | 11 (8–15)          | 3 (2–4.8)         | <0.001 |
| CPR to the first ROSC in ED (min)                                  | 14.5 (8–22)        | 10 (5–12)         | 0.002  |
| Initial pH during CPR in ED  | 6.97 (6.84–7.07)   | 7.11 (7–7.24)     | <0.001 |
| Initial lactate during CPR in ED (mmol/L)                          | 12 (8.33–15.46)    | 7.98 (5.63–11.93) | <0.001 |
| 5 <sup>th</sup> min ETCO <sub>2</sub> during CPR in ED (mmHg)      | 14 (10–21)         | 16 (11–43.8)      | 0.148  |
| Last ETCO <sub>2</sub> during CPR in ED (mmHg)                     | 14 (8–20)          | 47.5 (33–55.8)    | <0.001 |
| Highest ETCO <sub>2</sub> during CPR in ED (mmHg)                  | 20 (15–33.3)       | 49 (34.8–55.8)    | <0.001 |
| pH after ROSC in ED  | 7.03 (6.91–7.18)   | 7.25 (7.09–7.31)  | <0.001 |
| Lactate after ROSC in ED (mmol/L)                                  | 10.63 (6.33–14.13) | 4.59 (3.08–7.91)  | <0.001 |

\*Age: Independent Samples *t*-test; mean±SD. Variables other than age: Mann–Whitney *U*-test; median (25%–75%). CPR: Cardio-pulmonary resuscitation, ROSC: Return of the spontaneous circulation, ED: Emergency department, EtCO<sub>2</sub>: End-tidal carbon dioxide, SD: Standard deviation

**Supplementary Table 7: Factors associated with good neurological outcome at discharge**

| Variables                                    | Good neurological outcome at discharge, n (%) | P      |
|--|---|--------|
| Time zone                                    |   |        |
| Day shift                                    | 16 (4.2)                                      | 0.041  |
| Night shift and holidays                     | 11 (1.8)                                      |        |
| Gender                                       |   |        |
| Female                                       | 5 (1.3)                                       | 0.044  |
| Male   | 22 (3.6)                                      |        |
| Independent living                           |   |        |
| No   | -   | 0.009  |
| Yes  | 27 (3.8)                                      |        |
| Chronic neurological disease                 |   |        |
| No   | 25 (3.3)                                      | 0.067  |
| Yes  | 1 (0.6)                                       |        |
| Chronic pulmonary disease                    |   |        |
| No   | 25 (3.3)                                      | 0.076  |
| Yes  | 1 (0.6)                                       |        |
| Chronic cardiovascular disease               |   |        |
| No   | 17 (3.8)                                      | 0.100  |
| Yes  | 9 (1.8)                                       |        |
| Active malignancy                            |   |        |
| No   | 26 (3.2)                                      | 0.038  |
| Yes  | -   |        |
| Ventricular assist device                    |   |        |
| No   | 27 (2.9)                                      | 1.000  |
| Yes  | -   |        |
| ICD (or external cardioverter defibrillator) |   |        |
| No   | 27 (2.9)                                      | 1.000  |
| Yes  | -   |        |
| Witnessed arrest                             |   |        |
| Yes  | 24 (3.1)                                      | 0.160  |
| No   | 1 (0.7)                                       |        |
| Instructions from EMS dispatcher             |   |        |
| Yes  | 1 (1)   | 0.488  |
| No   | 15 (2.8)                                      |        |
| Prehospital CPR                              |   |        |
| None   | 9 (8.7)                                       | <0.001 |
| Health-care provider                         | 14 (1.6)                                      |        |
| Bystander CPR                                | 4 (13.8)                                      |        |
| Prehospital shockable rhythm                 |   |        |
| pVT/VF                                       | 15 (16)                                       | <0.001 |
| Asystole/NEA                                 | 5 (0.6)                                       |        |
| Prehospital defibrillation                   |   |        |
| No   | 16 (1.9)                                      | <0.001 |
| Yes  | 10 (8.5)                                      |        |
| Prehospital chest compression                |   |        |
| None   | 9 (8.7)                                       | 0.001  |
| Performed                                    | 18 (2)  |        |
| Prehospital ventilation support              |   |        |
| None   | 9 (10)  | <0.001 |
| Performed                                    | 18 (2)  |        |
| Prehospital mechanical CPR device            |   |        |
| None   | 26 (2.7)                                      | 1.000  |
| Used   | -   |        |

Contd...

**Supplementary Table 7: Contd...**

| Variables                           | Good neurological outcome at discharge, n (%) | P      |
|-------------------------------------|---|--------|
| Prehospital vascular access         |   |        |
| None                                | 6 (5.4)                                       | 0.110  |
| Yes                                 | 21 (2.4)                                      |        |
| Prehospital epinephrine             |   |        |
| None                                | 18 (8)  | <0.001 |
| Performed                           | 9 (1.2)                                       |        |
| Prehospital amiodarone              |   |        |
| None                                | 26 (2.7)                                      | 0.508  |
| Performed                           | 1 (4)   |        |
| Prehospital lidocaine               |   |        |
| None                                | 27 (2.8)                                      | 1.000  |
| Performed                           | -   |        |
| Prehospital sodium bicarbonate      |   |        |
| None                                | 27 (2.8)                                      | 1.000  |
| Performed                           | -   |        |
| Prehospital calcium                 |   |        |
| None                                | 27 (2.8)                                      | 1.000  |
| Performed                           | -   |        |
| Survival event                      |   |        |
| None                                | 20 (2.2)                                      | 0.001  |
| Yes                                 | 7 (10.4)                                      |        |
| Reported by EMS before ED admission |   |        |
| Yes                                 | 14 (2)  | 0.008  |
| No                                  | 13 (5.6)                                      |        |
| Etiology                            |   |        |
| Drowning                            | 2 (12.5)                                      | <0.001 |
| Acute coronary syndrome             | 22 (4.8)                                      |        |
| Pulmonary embolism                  | -   |        |
| Electrolyte disorder                | 1 (2)   |        |
| Asthma                              | -   |        |
| Others, unknown                     | 2 (0.5)                                       |        |
| Prehospital airway procedure        |   |        |
| Not performed                       | 14 (12.1)                                     | <0.001 |
| BVM                                 | 6 (1.9)                                       |        |
| ETI - successful/surgical           | 7 (1.5)                                       |        |
| Supraglottic                        | -   |        |
| Unsuccessful                        | -   |        |
| Cardiac arrest location             |   |        |
| Home                                | 6 (1.0)                                       | <0.001 |
| Workplace/public area               | 8 (6)   |        |
| Nursing home                        | -   |        |
| Ambulance                           | 6 (4.1)                                       |        |
| Hospital                            | 5 (12.5)                                      |        |
| Others                              | 1 (4.8)                                       |        |
| Shockable rhythm at ED admission    |   |        |
| pVT/VF                              | 12 (15.2)                                     | <0.001 |
| Asystole/PEA                        | 6 (0.8)                                       |        |
| Pulse exists                        | 9 (8.1)                                       |        |
| Advanced airway in ED               |   |        |
| Exists at admission                 | 7 (1.6)                                       | 0.188  |
| BVM/supraglottic/surgical           | 1 (4.3)                                       |        |
| ETI                                 | 19 (3.5)                                      |        |
| CPR in ED                           |   |        |
| None                                | 7 (14)  | <0.001 |
| Performed                           | 20 (2.1)                                      |        |

Contd...

Supplementary Table 7: Contd...

| Variables                                     | Good neurological outcome at discharge, n (%) | P      |
|---|---|--------|
| Mechanical CPR device in ED                   |   |        |
| None  | 20 (3.5)                                      | 0.133  |
| Used  | 7 (1.7)                                       |        |
| Defibrillation in ED                          |   |        |
| None  | 11 (1.4)                                      | <0.001 |
| Performed                                     | 16 (7)  |        |
| Venous access in ED                           |   |        |
| Others  | 1 (1.7)                                       | 1.000  |
| Peripheral                                    | 26 (2.8)                                      |        |
| Epinephrine in ED                             |   |        |
| None  | 9 (15.3)                                      | <0.001 |
| Performed                                     | 18 (1.9)                                      |        |
| Amiodarone in ED                              |   |        |
| None  | 15 (1.8)                                      | <0.001 |
| Performed                                     | 12 (7.7)                                      |        |
| Lidocaine in ED                               |   |        |
| None  | 26 (2.6)                                      | 0.129  |
| Performed                                     | 1 (20)  |        |
| Magnesium sulfate in ED                       |   |        |
| None  | 23 (2.4)                                      | 0.002  |
| Performed                                     | 4 (19)  |        |
| Sodium bicarbonate in ED                      |   |        |
| None  | 25 (3.4)                                      | 0.054  |
| Performed                                     | 2 (0.8)                                       |        |
| Calcium in ED                                 |   |        |
| None  | 26 (3)  | 0.160  |
| Performed                                     | 1 (0.7)                                       |        |
| Cardiac arrest again in ED                    |   |        |
| None  | 24 (10.7)                                     | <0.001 |
| Yes   | 3 (1.4)                                       |        |
| No ROSC in ED                                 | -   |        |
| USG use during CPR in ED                      |   |        |
| None  | 14 (3)  | 0.807  |
| Used  | 13 (2.5)                                      |        |
| If USG used, cardiac activity exists          |   |        |
| None  | -   | <0.001 |
| Exists  | 13 (7.6)                                      |        |
| EtCO <sub>2</sub> device use during CPR in ED |   |        |
| None  | 21 (3)  | 0.470  |
| Used  | 6 (2)   |        |
| ST elevation after ROSC                       |   |        |
| No  | 8 (3)   | <0.001 |
| Yes   | 19 (13.3)                                     |        |
| Coronary revascularization after ROSC         |   |        |
| No  | 5 (1.6)                                       | <0.001 |
| Yes   | 22 (22.2)                                     |        |
| TTM after ROSC                                |   |        |
| No  | 24 (6.2)                                      | 0.445  |
| Yes   | 1 (11.1)                                      |        |
| Vasoactive drug after ROSC                    |   |        |
| None  | 17 (12.1)                                     | 0.001  |
| Performed                                     | 9 (3.4)                                       |        |

Contd...

Supplementary Table 7: Contd...

| Variables            | Good neurological outcome at discharge, n (%) | P     |
|----------------------|---|-------|
| ECLS after ROSC      |   |       |
| None                 | 27 (6.7)                                      | 1.000 |
| Performed            | -   |       |
| Surgery after ROSC   |   |       |
| No                   | 26 (6.6)                                      | 0.430 |
| Yes                  | 1 (12.5)                                      |       |
| Forensic case report |   |       |
| None                 | 24 (2.7)                                      | 0.747 |
| Performed            | 3 (3)   |       |

Pearson's Chi-square test, Fisher's exact test. ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, ED: Emergency department, BVM: Bag-valve mask, ETI: Endotracheal intubation, ROSC: Return of the spontaneous circulation, USG: Ultrasonography, EtCO<sub>2</sub>: End-tidal carbon dioxide, TTM: Targeted temperature management, ECLS: Extracorporeal life support

**Supplementary Table 8: Factors associated with good neurological outcome at discharge - continuous variables**

| Variables  | Poor             | Good              | P*     |
|--|------------------|-------------------|--------|
| Age (year)   | 67.3±15.2        | 57.3±12.4         | 0.001  |
| Prehospital CPR duration (min)                                     | 15 (10–20)       | 2 (0–10)          | <0.001 |
| Prehospital time from the arrest to the CPR (min)                  | 5 (1–10)         | 2 (0.5–5.5)       | 0.047  |
| Prehospital number of defibrillations (if exist)                   | 2 (1–3)          | 1.5 (1–2.5)       | 0.457  |
| Prehospital time from the arrest to the first defibrillation (min) | 9 (4–15)         | 2 (0.5–4.5)       | 0.002  |
| Number of defibrillations before first ROSC in ED                  | 2 (1–3)          | 3 (1–5.5)         | 0.198  |
| Total number of defibrillations in ED                              | 3 (1–4)          | 3 (1–4.8)         | 0.856  |
| Epinephrine dose (mg) before the first ROSC in ED                  | 6 (3–10)         | 3 (1.5–4.5)       | <0.001 |
| Total epinephrine dose (mg) in ED                                  | 11 (8–15)        | 4 (1.8–5)         | <0.001 |
| CPR to the first ROSC in ED (min)                                  | 14 (8–21.8)      | 8 (5–14.3)        | 0.009  |
| Initial pH during CPR in ED  | 6.97 (6.84–7.07) | 7.16 (7.04–7.27)  | <0.001 |
| Initial lactate during CPR in ED (mmol/L)                          | 12 (8.33–15.42)  | 7.10 (5.30–11.80) | <0.001 |
| 5 <sup>th</sup> min ETCO <sub>2</sub> during CPR in ED (mmHg)      | 14 (10–21)       | 25.5 (12.5–47)    | 0.109  |
| Last ETCO <sub>2</sub> during CPR in ED (mmHg)                     | 14 (8–20)        | 43 (30–60.5)      | 0.001  |
| Highest ETCO <sub>2</sub> during CPR in ED (mmHg)                  | 21 (15–34)       | 44.5 (34.8–60.5)  | 0.002  |
| pH after ROSC in ED  | 7.05 (6.91–7.19) | 7.28 (7.09–7.33)  | <0.001 |
| Lactate after ROSC in ED (mmol/L)                                  | 10.35 (6.25–14)  | 4.05 (2.13–10.02) | <0.001 |

\*Age: Independent samples *t*-test, mean±SD. Variables other than age: Mann–Whitney *U*-test, median (25%–75%). CPR: Cardio-pulmonary resuscitation, ROSC: Return of the spontaneous circulation, ED: Emergency department, ETCO<sub>2</sub>: End-tidal carbon dioxide, SD: Standard deviation

**Supplementary Table 9: Distribution of outcomes in terms of regions**

|   | Regions        |               |                      |                         |                  |                         |                              |
|---|----------------|---------------|----------------------|-------------------------|------------------|-------------------------|------------------------------|
|   | Marmara, n (%) | Aegean, n (%) | Mediterranean, n (%) | Central Anatolia, n (%) | Black Sea, n (%) | Eastern Anatolia, n (%) | Southeastern Anatolia, n (%) |
| EMS   | 378 (97.7)     | 326 (96.4)    | 101 (98.1)           | 62 (96.9)               | 39 (97.5)        | 40 (100.0)              | 29 (100.0)                   |
| Bystander CPR                                 | 14 (3.6)       | 12 (3.6)      | 2 (2.0)              | 0                       | 0                | 1 (2.5)                 | 0                            |
| Shockable rhythm                              | 37 (11.5)      | 3 (11.5)      | 23 (24.2)            | 24 (8)                  | 2 (5.1)          | 3 (8.1)                 | 2 (3.8)                      |
| Prehospital airway procedure                  |                |               |                      |                         |                  |                         |                              |
| Not performed                                 | 39 (10.1)      | 2 (6.9)       | 20 (19.4)            | 40 (11.8)               | 4 (10)           | 2 (5.0)                 | 10 (15.6)                    |
| BVM   | 134 (34.6)     | 3 (10.3)      | 31 (30.1)            | 93 (27.4)               | 11 (27.5)        | 25 (62.5)               | 21 (32.8)                    |
| ETI - successful                              | 185 (47.8)     | 22 (75.9)     | 48 (46.6)            | 159 (46.9)              | 22 (55)          | 11 (27.5)               | 30 (46.9)                    |
| Supraglottic                                  | 12 (3.1)       | 2 (6.9)       | 3 (2.9)              | 33 (9.7)                | 0                | 1 (2.5)                 | 3 (4.7)                      |
| Surgical                                      | 1 (0.3)        | 0             | 0                    | 1 (0.3)                 | 2 (5.0)          | 0                       | 0                            |
| Unsuccessful                                  | 16 (4.1)       | 0             | 1 (1)                | 13 (3.8)                | 1 (2.5)          | 1 (2.5)                 | 0                            |
| Prehospital venous access                     |                |               |                      |                         |                  |                         |                              |
| None  | 35 (9.0)       | 1 (3.4)       | 9 (8.7)              | 45 (13.3)               | 5 (12.5)         | 9 (22.5)                | 8 (12.5)                     |
| Peripheral                                    | 351 (90.7)     | 27 (93.1)     | 93 (90.3)            | 293 (86.4)              | 35 (87.5)        | 31 (77.5)               | 56 (87.5)                    |
| Intraosseous                                  | 1 (0.3)        | 1 (3.4)       | 0                    | 0                       | 0                | 0                       | 0                            |
| Central                                       | 0              | 0             | 1 (1.0)              | 1 (0.3)                 | 0                | 0                       | 0                            |
| Mechanical CPR device in ED                   | 180 (46.5)     | 8 (27.6)      | 10 (9.7)             | 170 (50.1)              | 26 (65.0)        | 19 (47.5)               | 3 (4.7)                      |
| USG use during CPR in ED                      | 129 (33.3)     | 19 (65.5)     | 75 (72.8)            | 229 (67.6)              | 22 (55.0)        | 5 (12.5)                | 43 (67.2)                    |
| EtCO <sub>2</sub> device use during CPR in ED | 96 (24.8)      | 6 (20.7)      | 46 (44.7)            | 136 (40.1)              | 16 (40.0)        | 2 (5.0)                 | 0                            |
| Survived event                                | 15 (3.9)       | 19 (5.6)      | 17 (16.5)            | 5 (7.8)                 | 2 (5.0)          | 3 (7.5)                 | 8 (27.6)                     |
| Sustained ROSC                                | 102 (26.4)     | 114 (33.7)    | 59 (57.8)            | 21 (32.8)               | 12 (30.0)        | 12 (30.0)               | 14 (48.3)                    |
| Survival to hospital discharge                | 13 (3.4)       | 10 (2.9)      | 14 (13.6)            | 2 (3.1)                 | 3 (7.5)          | 1 (2.5)                 | 1 (3.4)                      |
| Good neurological outcome at discharge        | 11 (2.9)       | 4 (1.2)       | 11 (10.7)            | 0                       | 0                | 1 (2.5)                 | 0                            |

EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, BVM: Bag-valve mask, ETI: Endotracheal intubation, ED: Emergency department, USG: Ultrasonography, EtCO<sub>2</sub>: End-tidal carbon dioxide, ROSC: Return of the spontaneous circulation